



# Full wwPDB X-ray Structure Validation Report ⓘ

Aug 27, 2025 – 12:11 PM JST

PDB ID : 8JSU / pdb\_00008jsu  
Title : The crystal structure of TvaE in complex with TvaLP  
Authors : Li, M.; Pan, L.F.  
Deposited on : 2023-06-20  
Resolution : 1.44 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0rc1  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 2.0rc1  
EDS : 3.0  
buster-report : 1.1.7 (2018)  
Percentile statistics : 20231227.v01 (using entries in the PDB archive December 27th 2023)  
CCP4 : 9.0.006 (Gargrove)  
Density-Fitness : 1.0.12  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.45.1


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*





The reported resolution of this entry is 1.44 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	164625	2809 (1.46-1.42)
Clashscore	180529	3008 (1.46-1.42)
Ramachandran outliers	177936	2971 (1.46-1.42)
Sidechain outliers	177891	2971 (1.46-1.42)
RSRZ outliers	164620	2809 (1.46-1.42)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	42	
1	C	42	
2	B	315	
2	D	315	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit crite-

ria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
4	BR	B	402	-	-	X	-
6	PEG	B	405	-	-	X	-
7	GOL	B	406	-	-	X	-

## 2 Entry composition i

There are 8 unique types of molecules in this entry. The entry contains 6155 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called TvaLP.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	39	292	180	45	64	3	0	2	0
1	C	42	318	195	49	70	4	0	4	0

- Molecule 2 is a protein called TvaE.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	B	310	2418	1521	459	431	7	0	16	0
2	D	311	2404	1504	456	437	7	0	14	0

- Molecule 3 is SODIUM ION (CCD ID: NA) (formula: Na).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	B	1	Total	Na	0	0
			1	1		
3	D	1	Total	Na	0	0
			1	1		


- Molecule 4 is BROMIDE ION (CCD ID: BR) (formula: Br).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	B	2	Total	Br	0	0
			2	2		
4	D	1	Total	Br	0	0
			1	1		

- Molecule 5 is CHLORIDE ION (CCD ID: CL) (formula: Cl).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	B	1	Total Cl 1 1	0	0

- Molecule 6 is DI(HYDROXYETHYL)ETHER (CCD ID: PEG) (formula:  $C_4H_{10}O_3$ ) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	B	1	Total C O 7 4 3	0	0
6	D	1	Total C O 7 4 3	0	0

- Molecule 7 is GLYCEROL (CCD ID: GOL) (formula:  $C_3H_8O_3$ ).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
7	B	1	Total C O 6 3 3	0	0
7	B	1	Total C O 6 3 3	0	0
7	D	1	Total C O 6 3 3	0	0


- Molecule 8 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
8	A	24	Total O 24 24	0	0
8	B	300	Total O 300 300	0	0
8	C	39	Total O 39 39	0	0
8	D	322	Total O 322 322	0	0



### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.




- Molecule 1: TvaLP






- Molecule 1: TvaLP



- Molecule 2: TvaE



- Molecule 2: TvaE



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	45.22Å 52.50Å 73.67Å 94.72° 90.55° 112.55°	Depositor
Resolution (Å)	40.14 – 1.44 40.14 – 1.44	Depositor EDS
% Data completeness (in resolution range)	92.3 (40.14-1.44) 92.3 (40.14-1.44)	Depositor EDS
$R_{merge}$	0.04	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.16 (at 1.43Å)	Xtrriage
Refinement program	PHENIX (1.20.1_4487)	Depositor
R, $R_{free}$	0.173 , 0.197 0.173 , 0.198	Depositor DCC
$R_{free}$ test set	5267 reflections (4.63%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	10.7	Xtrriage
Anisotropy	0.677	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.38 , 41.8	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.50$ , $\langle L^2 \rangle = 0.33$	Xtrriage
Estimated twinning fraction	0.020 for h,-h-k,-l	Xtrriage
$F_o, F_c$ correlation	0.96	EDS
Total number of atoms	6155	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	19.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 7.30% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: GOL, NA, CL, PEG, BR

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.19	0/300	0.38	0/405
1	C	0.30	0/329	0.51	0/443
2	B	0.29	0/2521	0.53	0/3434
2	D	0.30	0/2501	0.53	0/3410
All	All	0.29	0/5651	0.52	0/7692

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
2	D	0	3

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (3) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
2	D	220[A]	ARG	Sidechain
2	D	220[B]	ARG	Sidechain
2	D	220[C]	ARG	Sidechain

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within

the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	292	0	282	5	0
1	C	318	0	308	9	0
2	B	2418	0	2504	35	0
2	D	2404	0	2461	23	0
3	B	1	0	0	0	0
3	D	1	0	0	0	0
4	B	2	0	0	3	0
4	D	1	0	0	0	0
5	B	1	0	0	0	0
6	B	7	0	10	5	0
6	D	7	0	10	0	0
7	B	12	0	16	6	0
7	D	6	0	8	0	0
8	A	24	0	0	1	0
8	B	300	0	0	10	2
8	C	39	0	0	1	1
8	D	322	0	0	13	2
All	All	6155	0	5599	70	3

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

All (70) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:D:35:GLU:OE2	8:D:501:HOH:O	1.76	1.02
2:B:55[A]:GLU:OE2	8:B:501:HOH:O	1.79	0.99
2:B:21:ASN:ND2	8:B:502:HOH:O	1.97	0.97
2:D:21[A]:ASN:OD1	8:D:502:HOH:O	1.90	0.89
2:B:188[A]:SER:OG	8:B:503:HOH:O	1.98	0.81
2:B:68:ILE:HB	7:B:407:GOL:H12	1.63	0.80
2:D:7[B]:ARG:NH2	8:D:507:HOH:O	2.14	0.79
4:B:403:BR:BR	8:B:748:HOH:O	2.58	0.76
2:B:115:LEU:HD21	2:B:187:LEU:HD22	1.68	0.75
2:B:306:GLY:HA3	7:B:406:GOL:H11	1.74	0.69
2:B:232[B]:GLN:NE2	8:B:507:HOH:O	2.26	0.68
2:D:224:GLN:OE1	8:D:505:HOH:O	2.12	0.68
2:B:17:GLY:O	8:B:502:HOH:O	2.13	0.67
2:D:35:GLU:OE1	8:D:506:HOH:O	2.12	0.67
2:D:156:ASP:OD2	8:D:508:HOH:O	2.15	0.64

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:D:39:VAL:HG22	8:D:749:HOH:O	1.97	0.64
2:B:39:VAL:HG22	8:B:727:HOH:O	1.97	0.64
2:D:95:SER:OG	2:D:229[A]:MET:HE2	1.99	0.63
2:D:309:ARG:O	8:D:509:HOH:O	2.16	0.62
1:A:18:VAL:HG22	2:B:23:VAL:HG21	1.84	0.60
2:D:274[A]:VAL:HG13	2:D:287:ALA:HB1	1.84	0.60
1:C:28[A]:MET:SD	1:C:33:ARG:HG2	2.44	0.57
1:C:17:LEU:HD23	2:D:23[B]:VAL:HG23	1.86	0.57
1:C:28[B]:MET:HE3	1:C:33:ARG:HG2	1.87	0.57
4:B:402:BR:BR	8:B:638:HOH:O	2.73	0.56
1:C:7:SER:O	1:C:11:GLU:HG3	2.08	0.54
2:B:94:GLU:OE1	2:B:98:SER:OG	2.20	0.54
2:D:91:THR:O	2:D:94:GLU:HB2	2.11	0.51
2:D:7[A]:ARG:NH2	8:D:521:HOH:O	2.43	0.51
2:B:219:LEU:HD12	2:B:262[B]:LEU:HD21	1.93	0.50
2:D:95:SER:OG	2:D:229[B]:MET:SD	2.68	0.50
2:B:199[B]:ARG:NE	6:B:405:PEG:H11	2.28	0.49
2:B:199[A]:ARG:HG3	6:B:405:PEG:H22	1.94	0.49
1:A:29:THR:OG1	1:A:32:GLN:HG3	2.12	0.49
2:B:233:GLN:HG3	2:B:237:GLN:OE1	2.13	0.48
2:D:7[B]:ARG:NH2	2:D:28[B]:ASP:OD1	2.47	0.47
2:B:68:ILE:O	7:B:407:GOL:H32	2.14	0.47
2:B:69:PRO:O	4:B:402:BR:BR	2.88	0.47
2:D:17:GLY:O	8:D:502:HOH:O	2.20	0.47
2:B:290[A]:LEU:HD11	8:B:549:HOH:O	2.14	0.46
2:B:103:GLN:OE1	2:B:196:PRO:HG3	2.16	0.45
2:B:229[B]:MET:HE3	2:B:229[B]:MET:HB2	1.71	0.45
1:A:32:GLN:O	1:A:35:VAL:HG22	2.16	0.45
2:B:199[B]:ARG:HE	6:B:405:PEG:H11	1.81	0.45
1:C:18:VAL:HG22	2:D:23[B]:VAL:HG11	1.98	0.45
2:B:7[B]:ARG:HD2	2:B:7[B]:ARG:H	1.81	0.45
2:D:232:GLN:HG3	8:D:619:HOH:O	2.17	0.45
2:D:277:ARG:NH1	8:D:514:HOH:O	2.26	0.44
2:B:150:ALA:O	2:B:154:ARG:HG3	2.17	0.44
2:B:115:LEU:HD11	2:B:187:LEU:HD23	1.99	0.44
2:B:197:ALA:O	6:B:405:PEG:O4	2.30	0.44
2:B:199[B]:ARG:HG3	6:B:405:PEG:H22	2.00	0.43
1:C:25:GLU:CD	1:C:33:ARG:HH22	2.25	0.43
2:B:122:PRO:HD2	2:B:125[A]:ARG:HE	1.83	0.43
7:B:406:GOL:H12	2:D:84:THR:HB	1.99	0.43
1:C:5:MET:HB2	1:C:5:MET:HE2	1.78	0.43

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:D:95:SER:CB	2:D:229[A]:MET:HE2	2.49	0.43
2:B:309:ARG:O	2:B:313:GLY:N	2.47	0.43
2:D:169:ALA:O	2:D:173:GLU:HG3	2.19	0.43
2:D:125[B]:ARG:NE	8:D:517:HOH:O	2.35	0.42
1:A:14[B]:PHE:HD2	8:A:115:HOH:O	2.01	0.42
2:B:305:PHE:C	7:B:406:GOL:H31	2.45	0.41
1:A:14[B]:PHE:HE2	2:B:16:PRO:HB3	1.85	0.41
2:B:277:ARG:HD2	2:B:277:ARG:O	2.20	0.41
2:B:293:ARG:HD2	8:B:539:HOH:O	2.20	0.40
1:C:5:MET:HE3	8:C:134:HOH:O	2.22	0.40
1:C:36:ILE:C	1:C:38:PRO:HD3	2.46	0.40
2:B:306:GLY:CA	7:B:406:GOL:H11	2.46	0.40
2:B:287:ALA:HA	2:B:290[C]:LEU:HD12	2.02	0.40
2:B:64:GLN:H	2:B:64:GLN:HG2	1.78	0.40

All (3) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
8:D:549:HOH:O	8:D:785:HOH:O[1_655]	1.93	0.27
8:B:643:HOH:O	8:C:123:HOH:O[1_445]	1.99	0.21
8:B:510:HOH:O	8:D:507:HOH:O[1_455]	2.19	0.01

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	39/42 (93%)	39 (100%)	0	0	100	100
1	C	44/42 (105%)	43 (98%)	1 (2%)	0	100	100
2	B	326/315 (104%)	322 (99%)	4 (1%)	0	100	100
2	D	325/315 (103%)	319 (98%)	6 (2%)	0	100	100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
All	All	734/714 (103%)	723 (98%)	11 (2%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	30/29 (103%)	30 (100%)	0	100	100
1	C	33/29 (114%)	33 (100%)	0	100	100
2	B	247/233 (106%)	247 (100%)	0	100	100
2	D	246/233 (106%)	245 (100%)	1 (0%)	89	78
All	All	556/524 (106%)	555 (100%)	1 (0%)	92	82

All (1) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
2	D	311	VAL

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (5) such sidechains are listed below:

Mol	Chain	Res	Type
2	B	10	GLN
2	B	21	ASN
2	B	233	GLN
2	B	253	GLN
2	D	61	GLN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

Of 11 ligands modelled in this entry, 6 are monoatomic - leaving 5 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
6	PEG	D	404	-	6,6,6	0.11	0	5,5,5	0.11	0
7	GOL	B	406	-	5,5,5	0.88	0	5,5,5	0.91	0
7	GOL	B	407	-	5,5,5	1.01	0	5,5,5	0.92	0
6	PEG	B	405	-	6,6,6	0.10	0	5,5,5	0.06	0
7	GOL	D	403	-	5,5,5	0.87	0	5,5,5	1.26	1 (20%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
6	PEG	D	404	-	-	4/4/4/4	-
7	GOL	B	406	-	-	2/4/4/4	-
7	GOL	B	407	-	-	2/4/4/4	-
6	PEG	B	405	-	-	4/4/4/4	-
7	GOL	D	403	-	-	2/4/4/4	-

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
7	D	403	GOL	C3-C2-C1	-2.15	103.35	111.70

There are no chirality outliers.

All (14) torsion outliers are listed below:



Mol	Chain	Res	Type	Atoms
7	B	406	GOL	O1-C1-C2-C3
7	B	407	GOL	C1-C2-C3-O3
7	D	403	GOL	C1-C2-C3-O3
6	B	405	PEG	O1-C1-C2-O2
6	D	404	PEG	O1-C1-C2-O2
7	B	407	GOL	O2-C2-C3-O3
6	B	405	PEG	O2-C3-C4-O4
7	B	406	GOL	O1-C1-C2-O2
7	D	403	GOL	O2-C2-C3-O3
6	D	404	PEG	O2-C3-C4-O4
6	B	405	PEG	C4-C3-O2-C2
6	D	404	PEG	C1-C2-O2-C3
6	B	405	PEG	C1-C2-O2-C3
6	D	404	PEG	C4-C3-O2-C2

There are no ring outliers.

3 monomers are involved in 11 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
7	B	406	GOL	4	0
7	B	407	GOL	2	0
6	B	405	PEG	5	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



## 5.7 Other polymers [\(i\)](#)

There are no such residues in this entry.



## 5.8 Polymer linkage issues

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	39/42 (92%)	1.05	7 (17%) <b>4</b> <b>3</b>	12, 26, 48, 64	2 (5%)
1	C	42/42 (100%)	1.11	8 (19%) <b>4</b> <b>3</b>	8, 20, 54, 67	4 (9%)
2	B	310/315 (98%)	0.27	25 (8%) <b>19</b> <b>18</b>	5, 13, 36, 63	16 (5%)
2	D	311/315 (98%)	0.14	20 (6%) <b>27</b> <b>26</b>	6, 14, 37, 79	14 (4%)
All	All	702/714 (98%)	0.30	60 (8%) <b>18</b> <b>17</b>	5, 15, 40, 79	36 (5%)

All (60) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	B	315	LEU	6.5
1	C	-1	GLY	5.8
1	C	0	SER	4.7
1	C	39	ALA	4.4
2	B	6	VAL	4.4
2	B	279	TRP	4.4
1	C	40	ALA	4.3
1	A	39	ALA	4.2
2	D	312	THR	4.1
2	D	311	VAL	4.1
2	D	95	SER	4.0
2	B	313	GLY	4.0
1	C	1	MET	3.9
1	A	38	PRO	3.8
1	A	29	THR	3.5
2	B	280	SER	3.5
2	D	279	TRP	3.4
2	D	310	GLU	3.3
2	D	96	PRO	3.2
2	D	6	VAL	3.2
2	D	99	ALA	3.1

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
2	D	281	ALA	3.1
2	B	8	THR	3.0
2	D	309	ARG	3.0
2	B	12	VAL	3.0
2	B	286	GLY	2.9
2	B	312	THR	2.9
2	D	282	SER	2.9
1	A	14[A]	PHE	2.8
2	B	65	TYR	2.8
2	B	100	VAL	2.7
2	D	98	SER	2.7
2	B	284	PRO	2.7
2	B	283	ASP	2.7
2	B	281	ALA	2.7
2	B	290[A]	LEU	2.6
1	A	31	GLU	2.6
2	D	100	VAL	2.6
2	B	311	VAL	2.6
1	A	35	VAL	2.5
2	B	314[A]	ARG	2.5
2	B	7[A]	ARG	2.5
1	C	38	PRO	2.5
1	A	1	MET	2.5
2	B	309	ARG	2.4
2	B	133	THR	2.4
2	B	101	VAL	2.4
1	C	6	ALA	2.4
2	D	97	ALA	2.4
2	B	310	GLU	2.4
2	B	64	GLN	2.3
2	D	158	ASP	2.3
1	C	2	SER	2.2
2	D	286	GLY	2.2
2	D	5	ALA	2.2
2	B	142	PRO	2.1
2	D	284	PRO	2.1
2	B	97	ALA	2.1
2	D	157	LYS	2.1
2	D	285	GLU	2.1

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

## 6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
6	PEG	B	405	7/7	0.77	0.17	32,38,41,54	0
6	PEG	D	404	7/7	0.77	0.15	30,38,41,41	0
7	GOL	B	406	6/6	0.84	0.14	28,32,36,41	0
7	GOL	B	407	6/6	0.86	0.12	21,31,35,37	0
4	BR	B	402	1/1	0.90	0.25	54,54,54,54	0
7	GOL	D	403	6/6	0.91	0.13	13,33,35,37	0
4	BR	B	403	1/1	0.93	0.13	31,31,31,31	0
3	NA	B	401	1/1	0.93	0.10	42,42,42,42	0
5	CL	B	404	1/1	0.98	0.04	11,11,11,11	0
3	NA	D	401	1/1	0.99	0.04	16,16,16,16	0
4	BR	D	402	1/1	0.99	0.06	20,20,20,20	0




The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

**Electron density around PEG B 405:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)

**Electron density around PEG D 404:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)



## 6.5 Other polymers [i](#)

There are no such residues in this entry.