



# Full wwPDB X-ray Structure Validation Report ⓘ

Nov 29, 2023 – 03:27 pm GMT

PDB ID : 8ORK  
Title : cyclic 2,3-diphosphoglycerate synthetase from the hyperthermophilic archaeon  
Methanothermus fervidus  
Authors : De Rose, S.A.; Isupov, M.  
Deposited on : 2023-04-14  
Resolution : 1.64 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.4, CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.36  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.36


# 1 Overall quality at a glance i

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 1.64 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	3122 (1.66-1.62)
Clashscore	141614	3268 (1.66-1.62)
Ramachandran outliers	138981	3215 (1.66-1.62)
Sidechain outliers	138945	3215 (1.66-1.62)
RSRZ outliers	127900	3079 (1.66-1.62)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	AAA	460	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	EDO	AAA	504	-	-	-	X

*Continued on next page...*

*Continued from previous page...*

<b>Mol</b>	<b>Type</b>	<b>Chain</b>	<b>Res</b>	<b>Chirality</b>	<b>Geometry</b>	<b>Clashes</b>	<b>Electron density</b>
4	PEG	AAA	527	-	-	-	X

## 2 Entry composition [i](#)


There are 7 unique types of molecules in this entry. The entry contains 8415 atoms, of which 4183 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Cyclic 2,3-diphosphoglycerate synthetase.

Mol	Chain	Residues	Atoms								ZeroOcc	AltConf	Trace
			Total	C	H	N	O	S	Se				
1	AAA	460	7824	2464	4008	625	700	10	17	78	45	0	

- Molecule 2 is 3[N-MORPHOLINO]PROPANE SULFONIC ACID (three-letter code: MPO) (formula:  $C_7H_{15}NO_4S$ ).



Mol	Chain	Residues	Atoms						ZeroOcc	AltConf
			Total	C	H	N	O	S		
2	AAA	1	28	7	15	1	4	1	1	0

- Molecule 3 is 1,2-ETHANEDIOL (three-letter code: EDO) (formula:  $C_2H_6O_2$ ).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	AAA	1	Total	C	H	O	1	0
			10	2	6	2		
3	AAA	1	Total	C	H	O	1	0
			10	2	6	2		
3	AAA	1	Total	C	H	O	1	0
			10	2	6	2		
3	AAA	1	Total	C	H	O	1	0
			10	2	6	2		
3	AAA	1	Total	C	H	O	1	0
			10	2	6	2		
3	AAA	1	Total	C	H	O	1	0
			10	2	6	2		
3	AAA	1	Total	C	H	O	1	0
			10	2	6	2		
3	AAA	1	Total	C	H	O	1	0
			10	2	6	2		
3	AAA	1	Total	C	H	O	1	0
			10	2	6	2		
3	AAA	1	Total	C	H	O	1	0
			10	2	6	2		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	AAA	1	Total	C	H	O	1	0
			10	2	6	2		
3	AAA	1	Total	C	H	O	1	0
			10	2	6	2		
3	AAA	1	Total	C	H	O	1	0
			10	2	6	2		
3	AAA	1	Total	C	H	O	1	0
			10	2	6	2		
3	AAA	1	Total	C	H	O	1	0
			10	2	6	2		
3	AAA	1	Total	C	H	O	1	0
			10	2	6	2		
3	AAA	1	Total	C	H	O	1	0
			10	2	6	2		
3	AAA	1	Total	C	H	O	1	0
			10	2	6	2		
3	AAA	1	Total	C	H	O	1	0
			10	2	6	2		

- Molecule 4 is DI(HYDROXYETHYL)ETHER (three-letter code: PEG) (formula: C<sub>4</sub>H<sub>10</sub>O<sub>3</sub>).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
4	AAA	1	Total	C	H	O	1	0
			17	4	10	3		

- Molecule 5 is SODIUM ION (three-letter code: NA) (formula: Na).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	AAA	1	Total	Na	0	0
			1	1		

- Molecule 6 is CHLORIDE ION (three-letter code: CL) (formula: Cl).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	AAA	1	Total	Cl	0	0
			1	1		


- Molecule 7 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	AAA	294	Total	O	0	0
			294	294		

### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Cyclic 2,3-diphosphoglycerate synthetase





## 4 Data and refinement statistics

Property	Value	Source
Space group	I 2 2 2	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	74.26Å 105.61Å 156.62Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	42.71 – 1.64 42.71 – 1.64	Depositor EDS
% Data completeness (in resolution range)	96.0 (42.71-1.64) 96.0 (42.71-1.64)	Depositor EDS
$R_{merge}$	0.08	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	0.98 (at 1.64Å)	Xtrriage
Refinement program	REFMAC 5.8.0267	Depositor
R, $R_{free}$	0.177 , 0.213 0.190 , 0.224	Depositor DCC
$R_{free}$ test set	3573 reflections (4.93%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	28.0	Xtrriage
Anisotropy	0.150	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.42 , 53.9	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.48$ , $\langle L^2 \rangle = 0.32$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
$F_o, F_c$ correlation	0.97	EDS
Total number of atoms	8415	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	38.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.25% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: MPO, NA, CL, PEG, EDO

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	AAA	0.68	0/3991	0.82	3/5361 (0.1%)

There are no bond length outliers.

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	AAA	451	MSE	CG-SE-CE	7.09	114.50	98.90
1	AAA	217	ARG	NE-CZ-NH1	-5.77	117.42	120.30
1	AAA	1	MSE	CG-SE-CE	5.04	110.00	98.90

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	AAA	3816	4008	4049	42	0
2	AAA	13	15	15	0	0
3	AAA	100	150	150	2	0
4	AAA	7	10	10	1	0
5	AAA	1	0	0	0	0
6	AAA	1	0	0	0	0
7	AAA	294	0	0	8	0
All	All	4232	4183	4224	43	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (43) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:AAA:32[B]:HIS:ND1	7:AAA:603:HOH:O	2.13	0.81
1:AAA:169:VAL:HG23	1:AAA:218:ILE:HD11	1.69	0.74
1:AAA:1:MSE:H1	1:AAA:1:MSE:SE	2.27	0.68
1:AAA:32[B]:HIS:HE1	7:AAA:656:HOH:O	1.77	0.66
1:AAA:337:PRO:HB2	1:AAA:427[A]:CYS:SG	2.36	0.66
1:AAA:32[A]:HIS:CD2	7:AAA:603:HOH:O	2.49	0.65
1:AAA:197:GLU:OE1	1:AAA:197:GLU:N	2.29	0.65
1:AAA:32[B]:HIS:CE1	7:AAA:656:HOH:O	2.50	0.64
1:AAA:256[B]:MSE:HG3	1:AAA:268:THR:HG21	1.80	0.62
1:AAA:190:ILE:HD12	1:AAA:405:LEU:HD21	1.81	0.62
1:AAA:447[A]:GLU:HG2	1:AAA:451:MSE:HE2	1.80	0.62
1:AAA:150:THR:HG22	1:AAA:215[A]:MSE:HE1	1.82	0.61
1:AAA:239:VAL:HG11	1:AAA:265:PRO:HD2	1.81	0.61
1:AAA:323[B]:GLU:HG2	1:AAA:324:ILE:HG23	1.84	0.59
1:AAA:178[B]:ARG:HH11	1:AAA:178[B]:ARG:HG3	1.70	0.57
1:AAA:32[B]:HIS:CE1	7:AAA:603:HOH:O	2.54	0.56
1:AAA:351:ALA:HB1	1:AAA:381:LEU:HD21	1.88	0.55
1:AAA:198:GLN:O	1:AAA:201[A]:LYS:HG3	2.06	0.54
1:AAA:261:ALA:HB1	1:AAA:292[B]:ARG:NH2	2.23	0.53
1:AAA:1:MSE:H1	1:AAA:1:MSE:HE3	1.73	0.53
1:AAA:337:PRO:CB	1:AAA:427[A]:CYS:SG	2.98	0.52
3:AAA:503:EDO:H22	3:AAA:521:EDO:O1	2.10	0.52
1:AAA:195:LEU:O	1:AAA:196:LEU:C	2.48	0.51
1:AAA:1:MSE:SE	1:AAA:1:MSE:N	2.94	0.49
1:AAA:439:GLY:HA3	3:AAA:525:EDO:O2	2.12	0.49
1:AAA:178[A]:ARG:NE	7:AAA:614:HOH:O	2.46	0.48
1:AAA:199:ALA:CB	1:AAA:410:VAL:HB	2.44	0.48
1:AAA:275:VAL:O	1:AAA:301:ILE:HA	2.14	0.48
1:AAA:1:MSE:H1	1:AAA:1:MSE:CE	2.28	0.47
1:AAA:196:LEU:HD23	1:AAA:196:LEU:HA	1.75	0.47
1:AAA:217:ARG:NH1	1:AAA:405:LEU:HD23	2.31	0.46
1:AAA:256[B]:MSE:HG3	1:AAA:268:THR:CG2	2.44	0.46
1:AAA:261:ALA:HB1	1:AAA:292[A]:ARG:NH2	2.32	0.45
1:AAA:178[B]:ARG:HG3	1:AAA:178[B]:ARG:NH1	2.29	0.45
1:AAA:105:ARG:NH1	7:AAA:621:HOH:O	2.50	0.44
1:AAA:256[B]:MSE:SE	1:AAA:266:VAL:HB	2.67	0.44
1:AAA:45[B]:GLU:HG2	7:AAA:718:HOH:O	2.18	0.44

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:AAA:199:ALA:HA	1:AAA:210:TRP:CZ3	2.52	0.44
1:AAA:260:GLY:H	4:AAA:527:PEG:H12	1.82	0.44
1:AAA:150:THR:CG2	1:AAA:215[B]:MSE:HE1	2.49	0.43
1:AAA:138[A]:ILE:HA	1:AAA:270:ARG:O	2.19	0.42
1:AAA:351:ALA:HB1	1:AAA:381:LEU:CD2	2.51	0.41
1:AAA:137:SER:OG	1:AAA:268:THR:HA	2.21	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	AAA	503/460 (109%)	494 (98%)	9 (2%)	0	100 100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	AAA	436/380 (115%)	419 (96%)	17 (4%)	32 8

All (17) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	AAA	1	MSE
1	AAA	149[A]	LYS
1	AAA	149[B]	LYS
1	AAA	178[A]	ARG
1	AAA	178[B]	ARG
1	AAA	198	GLN
1	AAA	201[A]	LYS
1	AAA	201[B]	LYS
1	AAA	217	ARG
1	AAA	225	ARG
1	AAA	303[A]	MSE
1	AAA	303[B]	MSE
1	AAA	391	LEU
1	AAA	403	THR
1	AAA	416	VAL
1	AAA	434	ILE
1	AAA	460	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

### 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

### 5.6 Ligand geometry [i](#)

Of 29 ligands modelled in this entry, 2 are monoatomic - leaving 27 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond

length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
3	EDO	AAA	516	-	3,3,3	0.19	0	2,2,2	0.38	0
3	EDO	AAA	511	-	3,3,3	0.10	0	2,2,2	0.22	0
3	EDO	AAA	515	-	3,3,3	0.07	0	2,2,2	0.12	0
3	EDO	AAA	504	-	3,3,3	0.27	0	2,2,2	0.29	0
3	EDO	AAA	524	-	3,3,3	0.06	0	2,2,2	0.24	0
3	EDO	AAA	514	-	3,3,3	0.06	0	2,2,2	0.12	0
3	EDO	AAA	518	-	3,3,3	0.20	0	2,2,2	0.32	0
2	MPO	AAA	501	-	13,13,13	0.58	0	17,17,17	1.23	2 (11%)
3	EDO	AAA	521	-	3,3,3	0.05	0	2,2,2	0.11	0
3	EDO	AAA	503	-	3,3,3	0.23	0	2,2,2	0.32	0
3	EDO	AAA	502	-	3,3,3	0.08	0	2,2,2	0.14	0
3	EDO	AAA	520	-	3,3,3	0.22	0	2,2,2	0.21	0
3	EDO	AAA	505	-	3,3,3	0.26	0	2,2,2	0.21	0
3	EDO	AAA	522	-	3,3,3	0.09	0	2,2,2	0.14	0
3	EDO	AAA	526	-	3,3,3	0.07	0	2,2,2	0.27	0
4	PEG	AAA	527	-	6,6,6	0.30	0	5,5,5	0.16	0
3	EDO	AAA	509	-	3,3,3	0.27	0	2,2,2	0.35	0
3	EDO	AAA	519	-	3,3,3	0.15	0	2,2,2	0.04	0
3	EDO	AAA	525	-	3,3,3	0.21	0	2,2,2	0.23	0
3	EDO	AAA	508	-	3,3,3	0.16	0	2,2,2	0.20	0
3	EDO	AAA	512	-	3,3,3	0.04	0	2,2,2	0.20	0
3	EDO	AAA	507	-	3,3,3	0.13	0	2,2,2	0.27	0
3	EDO	AAA	506	-	3,3,3	0.13	0	2,2,2	0.07	0
3	EDO	AAA	513	-	3,3,3	0.26	0	2,2,2	0.10	0
3	EDO	AAA	517	-	3,3,3	0.38	0	2,2,2	0.29	0
3	EDO	AAA	510	-	3,3,3	0.06	0	2,2,2	0.31	0
3	EDO	AAA	523	-	3,3,3	0.16	0	2,2,2	0.22	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	EDO	AAA	516	-	-	1/1/1/1	-
3	EDO	AAA	511	-	-	1/1/1/1	-
3	EDO	AAA	515	-	-	1/1/1/1	-
3	EDO	AAA	504	-	-	1/1/1/1	-
3	EDO	AAA	524	-	-	0/1/1/1	-

*Continued on next page...*

*Continued from previous page...*

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	EDO	AAA	514	-	-	1/1/1/1	-
3	EDO	AAA	518	-	-	1/1/1/1	-
2	MPO	AAA	501	-	-	0/7/15/15	0/1/1/1
3	EDO	AAA	521	-	-	1/1/1/1	-
3	EDO	AAA	503	-	-	1/1/1/1	-
3	EDO	AAA	502	-	-	0/1/1/1	-
3	EDO	AAA	520	-	-	1/1/1/1	-
3	EDO	AAA	505	-	-	1/1/1/1	-
3	EDO	AAA	522	-	-	1/1/1/1	-
3	EDO	AAA	526	-	-	1/1/1/1	-
4	PEG	AAA	527	-	-	1/4/4/4	-
3	EDO	AAA	509	-	-	0/1/1/1	-
3	EDO	AAA	519	-	-	1/1/1/1	-
3	EDO	AAA	525	-	-	0/1/1/1	-
3	EDO	AAA	508	-	-	0/1/1/1	-
3	EDO	AAA	512	-	-	0/1/1/1	-
3	EDO	AAA	507	-	-	1/1/1/1	-
3	EDO	AAA	506	-	-	1/1/1/1	-
3	EDO	AAA	513	-	-	1/1/1/1	-
3	EDO	AAA	517	-	-	1/1/1/1	-
3	EDO	AAA	510	-	-	1/1/1/1	-
3	EDO	AAA	523	-	-	0/1/1/1	-

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	AAA	501	MPO	O3-S1-O1	-3.29	103.23	111.27
2	AAA	501	MPO	O2-S1-O1	3.22	125.10	113.95

There are no chirality outliers.

All (19) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	AAA	510	EDO	O1-C1-C2-O2
3	AAA	511	EDO	O1-C1-C2-O2
3	AAA	513	EDO	O1-C1-C2-O2
3	AAA	514	EDO	O1-C1-C2-O2
3	AAA	515	EDO	O1-C1-C2-O2
3	AAA	517	EDO	O1-C1-C2-O2
3	AAA	518	EDO	O1-C1-C2-O2

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms
3	AAA	519	EDO	O1-C1-C2-O2
3	AAA	520	EDO	O1-C1-C2-O2
3	AAA	526	EDO	O1-C1-C2-O2
4	AAA	527	PEG	O2-C3-C4-O4
3	AAA	516	EDO	O1-C1-C2-O2
3	AAA	521	EDO	O1-C1-C2-O2
3	AAA	522	EDO	O1-C1-C2-O2
3	AAA	505	EDO	O1-C1-C2-O2
3	AAA	503	EDO	O1-C1-C2-O2
3	AAA	507	EDO	O1-C1-C2-O2
3	AAA	506	EDO	O1-C1-C2-O2
3	AAA	504	EDO	O1-C1-C2-O2

There are no ring outliers.

4 monomers are involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	AAA	521	EDO	1	0
3	AAA	503	EDO	1	0
4	AAA	527	PEG	1	0
3	AAA	525	EDO	1	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.



## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	AAA	448/460 (97%)	0.29	37 (8%) <b>11</b> <b>9</b>	17, 31, 69, 108	0

All (37) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	AAA	200	ASP	8.0
1	AAA	49	ILE	6.3
1	AAA	460	ARG	5.9
1	AAA	204	HIS	5.6
1	AAA	196	LEU	5.5
1	AAA	203	VAL	5.5
1	AAA	50	GLU	4.7
1	AAA	201[A]	LYS	4.7
1	AAA	194	TYR	4.5
1	AAA	358	VAL	4.4
1	AAA	411[A]	ASP	4.3
1	AAA	459[A]	ASN	4.3
1	AAA	199	ALA	4.2
1	AAA	458[A]	ASN	4.1
1	AAA	198	GLN	3.8
1	AAA	47	LEU	3.7
1	AAA	202	GLY	3.6
1	AAA	357	VAL	3.4
1	AAA	48	GLN	3.4
1	AAA	53[A]	LYS	3.3
1	AAA	193[A]	GLU	3.3
1	AAA	44	THR	3.1
1	AAA	51	ASP	2.9
1	AAA	65	PHE	2.9
1	AAA	250[A]	ASP	2.9
1	AAA	382	SER	2.8
1	AAA	54	GLU	2.7

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	AAA	52	PRO	2.7
1	AAA	124	PHE	2.5
1	AAA	412	VAL	2.3
1	AAA	395	ILE	2.3
1	AAA	236	ILE	2.2
1	AAA	456	PHE	2.2
1	AAA	55	TYR	2.2
1	AAA	58	LYS	2.2
1	AAA	178[A]	ARG	2.1
1	AAA	195	LEU	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
3	EDO	AAA	507	4/4	0.59	0.19	46,59,61,67	10
3	EDO	AAA	523	4/4	0.63	0.16	37,54,57,57	10
3	EDO	AAA	502	4/4	0.78	0.18	46,54,62,65	10
3	EDO	AAA	514	4/4	0.78	0.23	48,56,59,60	10
3	EDO	AAA	516	4/4	0.78	0.14	31,42,62,62	10
3	EDO	AAA	504	4/4	0.78	0.84	38,53,74,80	10
3	EDO	AAA	525	4/4	0.79	0.21	46,54,59,66	10
4	PEG	AAA	527	7/7	0.79	0.43	35,49,81,87	17
3	EDO	AAA	509	4/4	0.80	0.31	39,54,70,73	1
3	EDO	AAA	517	4/4	0.81	0.14	47,50,66,71	10
3	EDO	AAA	526	4/4	0.84	0.65	42,45,59,60	10
3	EDO	AAA	515	4/4	0.84	0.13	48,55,71,71	10
3	EDO	AAA	508	4/4	0.85	0.11	54,60,63,67	1

*Continued on next page...*

*Continued from previous page...*

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
3	EDO	AAA	524	4/4	0.85	0.34	49,54,60,60	10
3	EDO	AAA	510	4/4	0.85	0.25	38,49,55,56	10
3	EDO	AAA	519	4/4	0.85	0.27	53,55,63,63	10
3	EDO	AAA	522	4/4	0.85	0.23	49,54,63,64	10
3	EDO	AAA	520	4/4	0.86	0.27	32,48,61,62	10
3	EDO	AAA	511	4/4	0.89	0.16	39,48,64,73	10
3	EDO	AAA	506	4/4	0.90	0.11	30,46,63,65	10
3	EDO	AAA	505	4/4	0.91	0.13	48,51,53,57	1
3	EDO	AAA	512	4/4	0.92	0.07	52,57,68,68	10
3	EDO	AAA	513	4/4	0.94	0.21	43,49,61,63	1
3	EDO	AAA	521	4/4	0.94	0.31	32,44,68,72	10
3	EDO	AAA	518	4/4	0.95	0.14	27,40,44,49	10
3	EDO	AAA	503	4/4	0.97	0.12	27,43,54,64	1
6	CL	AAA	529	1/1	0.97	0.14	47,47,47,47	0
2	MPO	AAA	501	13/13	0.98	0.14	24,40,49,54	1
5	NA	AAA	528	1/1	1.00	0.17	25,25,25,25	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.