

Full wwPDB EM Validation Report (i)

Nov 3, 2024 – 09:19 pm GMT

PDB ID : 8PNQ

EMDB ID : EMD-17783

Title: Influenza A/H7N9 polymerase in elongation state with continuous Pol II pS5

CTD peptide mimic bound in site 1A/2A

Authors : Arragain, B.; Cusack, S.

Deposited on : 2023-06-30

Resolution : 2.88 Å(reported)

Based on initial model : 7QTL

This is a Full wwPDB EM Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
https://www.wwpdb.org/validation/2017/EMValidationReportHelp
with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (1)) were used in the production of this report:

EMDB validation analysis : 0.0.1.dev113

Mogul : 1.8.4, CSD as541be (2020)

MolProbity : 4.02b-467 buster-report : 1.1.7 (2018)

Percentile statistics : 20231227.v01 (using entries in the PDB archive December 27th 2023)

 $MapQ \quad : \quad 1.9.13$

Ideal geometry (proteins) : Engh & Huber (2001) Ideal geometry (DNA, RNA) : Parkinson et al. (1996)


Validation Pipeline (wwPDB-VP) : 2.39

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: $ELECTRON\ MICROSCOPY$

The reported resolution of this entry is 2.88 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive $(\# \mathrm{Entries})$	${ m EM~structures} \ (\#{ m Entries})$
Clashscore	210492	15764
Ramachandran outliers	207382	16835
Sidechain outliers	206894	16415
RNA backbone	6643	2191

The table below summarises the geometric issues observed across the polymeric chains and their fit to the map. The red, orange, yellow and green segments of the bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the EM map (all-atom inclusion <40%). The numeric value is given above the bar.

Mol	Chain	Length		Qualit	ty of chai	n			
1	A	729	6%	84%				12%	•
2	В	757	<u>•</u>	87%				11%	•••
3	С	788	28%	·		68%			_
4	V	51	22%		20%	•	29%		_
5	M	33	27%		7.	3%			_
6	G	42	19%		12% •		40%		_



2 Entry composition (i)

There are 8 unique types of molecules in this entry. The entry contains 14913 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

• Molecule 1 is a protein called Polymerase acidic protein.

\mathbf{Mol}	Chain	Residues		\mathbf{A}	toms			AltConf	Trace
1	A	708	Total 5748	C 3641	N 976	O 1090	S 41	0	0

There are 15 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-12	MET	-	initiating methionine	UNP M9TI86
A	-11	HIS	-	expression tag	UNP M9TI86
A	-10	HIS	-	expression tag	UNP M9TI86
A	-9	HIS	-	expression tag	UNP M9TI86
A	-8	HIS	-	expression tag	UNP M9TI86
A	-7	HIS	-	expression tag	UNP M9TI86
A	-6	HIS	-	expression tag	UNP M9TI86
A	-5	HIS	-	expression tag	UNP M9TI86
A	-4	HIS	-	expression tag	UNP M9TI86
A	-3	GLY	-	expression tag	UNP M9TI86
A	-2	SER	-	expression tag	UNP M9TI86
A	-1	GLY	-	expression tag	UNP M9TI86
A	0	SER	-	expression tag	UNP M9TI86
A	349	LYS	GLU	engineered mutation	UNP M9TI86
A	490	ILE	ARG	engineered mutation	UNP M9TI86

• Molecule 2 is a protein called RNA-directed RNA polymerase catalytic subunit.

Mol (Chain	Residues		Α	toms			AltConf	Trace
2	В	744	Total 5958	C 3743	N 1049	O 1119	S 47	1	0

There is a discrepancy between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
В	577	GLY	LYS	engineered mutation	UNP S5ME50



• Molecule 3 is a protein called Polymerase basic protein 2.

Mo	Chain	Residues		At	oms			AltConf	Trace
3	С	249	Total 2024	C 1276	N 365	O 367	S 16	0	0

There are 30 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
С	74	ARG	GLY	engineered mutation	UNP X5F427
С	760	GLY	-	expression tag	UNP X5F427
С	761	TRP	-	expression tag	UNP X5F427
С	762	SER	-	expression tag	UNP X5F427
С	763	HIS	-	expression tag	UNP X5F427
С	764	PRO	-	expression tag	UNP X5F427
С	765	GLN	-	expression tag	UNP X5F427
С	766	PHE	-	expression tag	UNP X5F427
С	767	GLU	-	expression tag	UNP X5F427
С	768	LYS	-	expression tag	UNP X5F427
С	769	GLY	-	expression tag	UNP X5F427
С	770	GLY	-	expression tag	UNP X5F427
С	771	GLY	_	expression tag	UNP X5F427
С	772	SER	-	expression tag	UNP X5F427
С	773	GLY	_	expression tag	UNP X5F427
С	774	GLY	-	expression tag	UNP X5F427
С	775	GLY	_	expression tag	UNP X5F427
С	776	SER	-	expression tag	UNP X5F427
С	777	GLY	-	expression tag	UNP X5F427
С	778	GLY	_	expression tag	UNP X5F427
С	779	SER	-	expression tag	UNP X5F427
С	780	ALA	-	expression tag	UNP X5F427
С	781	TRP	-	expression tag	UNP X5F427
С	782	SER	-	expression tag	UNP X5F427
С	783	HIS	-	expression tag	UNP X5F427
С	784	PRO	-	expression tag	UNP X5F427
С	785	GLN	-	expression tag	UNP X5F427
С	786	PHE	-	expression tag	UNP X5F427
С	787	GLU	-	expression tag	UNP X5F427
С	788	LYS	-	expression tag	UNP X5F427

 \bullet Molecule 4 is a RNA chain called 51-mer vRNA loop (v51_mut_S).



Mol	Chain	Residues		A	toms			AltConf	Trace
4	V	36	Total 758	C 339	N 124	O 259	P 36	0	0


• Molecule 5 is a RNA chain called Replication product.

Mol	Chain	Residues		\mathbf{At}	oms			AltConf	Trace
5	M	9	Total 202	C 90		O 58	P o	0	0

• Molecule 6 is a protein called RNA Pol II CTD 6 repeats (site 1A/2A).

Mo	Chain	Residues		Ato	oms			AltConf	Trace
6	C	25	Total	С	N	О	Р	0	0
0	G	25	192	116	25	49	2	U	U

• Molecule 7 is 5'-O-[(S)-hydroxy{[(S)-hydroxy(phosphonooxy)phosphoryl]amino}phosphoryl] uridine (three-letter code: 2KH) (formula: $C_9H_{16}N_3O_{14}P_3$) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues		Atoms				
7	D	1	Total	С	N	О	Р	0
(D	1	29	9	3	14	3	U

• Molecule 8 is MAGNESIUM ION (three-letter code: MG) (formula: Mg) (labeled as "Ligand of Interest" by depositor).




Mol	Chain	Residues	Atoms	AltConf
8	В	2	Total Mg 2 2	0




3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and atom inclusion in map density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red diamond above a residue indicates a poor fit to the EM map for this residue (all-atom inclusion < 40%). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Polymerase acidic protein




• Molecule 2: RNA-directed RNA polymerase catalytic subunit







• Molecule 3: Polymerase basic protein 2



• Molecule 5: Replication product

Chain M: 27% 73%





 \bullet Molecule 6: RNA Pol II CTD 6 repeats (site 1A/2A)







4 Experimental information (i)

Property	Value	Source
EM reconstruction method	SINGLE PARTICLE	Depositor
Imposed symmetry	POINT, C1	Depositor
Number of particles used	14878	Depositor
Resolution determination method	FSC 0.143 CUT-OFF	Depositor
CTF correction method	PHASE FLIPPING AND AMPLITUDE	Depositor
	CORRECTION	
Microscope	FEI TITAN KRIOS	Depositor
Voltage (kV)	300	Depositor
Electron dose $(e^-/\text{Å}^2)$	40	Depositor
Minimum defocus (nm)	800	Depositor
Maximum defocus (nm)	2000	Depositor
Magnification	Not provided	
Image detector	GATAN K3 BIOQUANTUM (6k x 4k)	Depositor
Maximum map value	0.045	Depositor
Minimum map value	-0.021	Depositor
Average map value	0.000	Depositor
Map value standard deviation	0.002	Depositor
Recommended contour level	0.008	Depositor
Map size (Å)	302.4, 302.4, 302.4	wwPDB
Map dimensions	360, 360, 360	wwPDB
Map angles (°)	90.0, 90.0, 90.0	wwPDB
Pixel spacing (Å)	0.84, 0.84, 0.84	Depositor



5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: MG, 2KH, SEP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol Chain		Bo	nd lengths	Bond angles		
IVIOI	Chain	RMSZ	# Z > 5	RMSZ	# Z >5	
1	A	0.25	0/5870	0.48	0/7911	
2	В	0.25	0/6071	0.50	0/8180	
3	С	0.25	0/2064	0.51	0/2782	
4	V	0.43	1/843 (0.1%)	0.74	0/1305	
5	M	0.24	0/228	0.68	0/355	
6	G	0.28	0/180	0.40	0/247	
All	All	0.26	$1/15256 \ (0.0\%)$	0.52	0/20780	

All (1) bond length outliers are listed below:

\mathbf{Mol}	Chain	Res	Type	Atoms	\mathbf{Z}	$\operatorname{Observed}(\text{\AA})$	$\operatorname{Ideal}(ext{\AA})$
4	V	1	A	OP3-P	-10.76	1.48	1.61

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	5748	0	5689	54	0
2	В	5958	0	5957	54	0
3	С	2024	0	2075	14	0
4	V	758	0	384	3	0



Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
5	M	202	0	99	0	0
6	G	192	0	156	1	0
7	В	29	0	15	0	0
8	В	2	0	0	0	0
All	All	14913	0	14375	106	0

The all-atom clash score is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clash score for this structure is 4.

All (106) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

A 4 a 1	A + 9	Interatomic	Clash	
Atom-1	Atom-2	${\rm distance}(\mathring{\rm A})$	overlap (Å)	
1:A:243:GLU:N	1:A:243:GLU:OE2	2.10	0.83	
4:V:5:G:N2	4:V:5:G:OP2	2.15	0.80	
2:B:423:ILE:HD11	2:B:473:VAL:HG21	1.66	0.78	
2:B:460:GLN:NE2	2:B:464:ASP:OD1	2.19	0.75	
1:A:358:LYS:NZ	1:A:359:ASN:O	2.19	0.73	
2:B:493:THR:HG22	2:B:493:THR:O	1.88	0.72	
2:B:226:THR:OG1	2:B:350:ARG:O	2.05	0.72	
1:A:2:GLU:N	1:A:2:GLU:OE1	2.26	0.69	
1:A:43:GLU:OE2	1:A:174:ARG:NH1	2.27	0.67	
1:A:141:GLU:N	1:A:141:GLU:OE2	2.27	0.67	
1:A:110:TYR:OH	1:A:115:ASN:OD1	2.12	0.67	
1:A:146:HIS:NE2	1:A:154:GLU:OE2	2.28	0.66	
2:B:629:ASN:ND2	2:B:661:ALA:O	2.28	0.65	
1:A:344:GLU:OE1	1:A:358:LYS:NZ	2.24	0.65	
2:B:583:THR:O	2:B:586:LYS:NZ	2.31	0.64	
3:C:17:ARG:O	3:C:21:THR:HG22	1.98	0.64	
2:B:626:ASN:ND2	3:C:104:PRO:O	2.30	0.64	
2:B:158:ASN:ND2	2:B:169:ASP:OD2	2.31	0.64	
1:A:239:ASN:O	1:A:239:ASN:OD1	2.15	0.63	
2:B:288:LYS:NZ	2:B:439:ASP:OD2	2.31	0.63	
1:A:424:GLU:O	2:B:553:LYS:NZ	2.32	0.63	
4:V:7:A:O2'	4:V:8:A:OP2	2.17	0.62	
1:A:593:GLU:OE1	2:B:543:THR:OG1	2.17	0.62	
2:B:715:VAL:HG21	3:C:11:MET:HE1	1.84	0.60	
1:A:427:GLU:N	1:A:427:GLU:OE1	2.35	0.58	
2:B:158:ASN:N	2:B:169:ASP:OD2	2.36	0.57	
1:A:623:GLU:OE1	1:A:713:HIS:NE2	2.38	0.56	
1:A:309:LYS:O	1:A:313:THR:HG23	2.06	0.56	
2:B:86:ASP:OD1	2:B:87:CYS:N	2.40	0.55	



Continued from previous page...

Continued from prev		Interatomic	Clash
Atom-1	Atom-2	${\rm distance}\ ({\rm \AA})$	overlap (Å)
3:C:10:LEU:CD1	3:C:16:THR:HG21	2.38	0.54
1:A:292:ILE:HG22	1:A:313:THR:HG21	1.89	0.54
1:A:30:ILE:HD12	1:A:31:GLU:N	2.22	0.54
1:A:203:GLU:OE2	2:B:162:ARG:NH2	2.40	0.54
2:B:575:GLU:OE1	3:C:101:ARG:NE	2.41	0.54
2:B:727:ARG:NH1	2:B:731:GLU:OE1	2.41	0.53
1:A:77:GLU:OE2	1:A:78:ILE:N	2.42	0.53
2:B:131:TRP:O	2:B:220:ARG:NH1	2.39	0.53
3:C:60:ASP:OD1	3:C:61:LYS:N	2.44	0.51
1:A:165:GLU:OE2	1:A:168:ARG:NH1	2.43	0.50
1:A:470:LEU:HD22	1:A:579:MET:CE	2.41	0.49
3:C:10:LEU:HG	3:C:16:THR:HG21	1.95	0.49
2:B:684:GLU:O	2:B:688:MET:HG2	2.13	0.49
1:A:382:GLU:OE1	1:A:382:GLU:N	2.42	0.49
3:C:21:THR:HG23	3:C:22:LYS:HG2	1.95	0.49
1:A:153:GLU:OE1	1:A:154:GLU:N	2.46	0.48
2:B:572:ARG:HG2	2:B:576:LEU:HD13	1.95	0.48
2:B:182:THR:O	2:B:182:THR:HG23	2.12	0.48
1:A:141:GLU:HG2	1:A:141:GLU:O	2.13	0.48
2:B:340:ALA:HB3	2:B:341:PRO:HD3	1.96	0.48
1:A:90:VAL:HG22	2:B:724:ILE:HG12	1.94	0.47
1:A:256:ARG:NH1	1:A:257:ILE:O	2.47	0.47
1:A:473:SER:O	1:A:509:SER:OG	2.33	0.47
3:C:146:ASP:OD1	3:C:146:ASP:C	2.52	0.47
2:B:36:THR:HG23	2:B:351:LEU:HB3	1.96	0.47
2:B:302:ILE:HG23	2:B:449:LEU:HB3	1.97	0.47
2:B:584:ARG:HH11	2:B:584:ARG:HG3	1.79	0.47
2:B:561:CYS:HA	2:B:576:LEU:HD21	1.97	0.46
2:B:655:MET:CE	2:B:655:MET:HA	2.45	0.46
1:A:30:ILE:HD12	1:A:30:ILE:C	2.36	0.46
3:C:55:TYR:OH	3:C:76:THR:HG21	2.15	0.45
1:A:85:THR:HG21	2:B:709:VAL:O	2.16	0.45
1:A:674:ASP:OD2	2:B:498:ARG:NH2	2.48	0.45
1:A:181:GLU:OE1	1:A:181:GLU:O	2.34	0.45
2:B:584:ARG:HG3	2:B:584:ARG:NH1	2.30	0.45
1:A:587:GLN:HB2	2:B:509:LEU:HD11	1.98	0.45
1:A:189:ASP:N	1:A:189:ASP:OD1	2.49	0.45
2:B:130:ASP:O	2:B:134:ASN:N	2.50	0.45
1:A:181:GLU:OE1	1:A:185:ARG:NE	2.50	0.45
2:B:636:GLU:HA	2:B:636:GLU:OE2	2.17	0.44
2:B:131:TRP:CH2	2:B:149:VAL:HG11	2.53	0.44



Continued from previous page...

Continued from press		Interatomic	Clash
Atom-1	Atom-2	${ m distance} \; ({ m \AA})$	overlap (Å)
2:B:423:ILE:HD11	2:B:473:VAL:CG2	2.44	0.44
3:C:231:VAL:HG22	3:C:234:LEU:HB3	1.99	0.44
1:A:638:ARG:NH1	6:G:7:SEP:O2P	2.48	0.44
1:A:111:ASP:O	1:A:115:ASN:N	2.50	0.44
1:A:166:GLU:O	1:A:170:ARG:HG3	2.17	0.44
2:B:110:THR:HG21	2:B:258:LEU:HB2	2.01	0.43
2:B:724:ILE:HD12	2:B:724:ILE:HA	1.90	0.43
1:A:169:ALA:O	1:A:173:THR:HG23	2.19	0.43
1:A:322:ILE:HD13	1:A:543:LEU:CD2	2.49	0.43
1:A:341:VAL:CG2	1:A:505:ILE:HD11	2.49	0.43
1:A:21:MET:SD	1:A:35:PHE:HA	2.59	0.43
2:B:576:LEU:HD12	3:C:97:THR:HG21	2.01	0.43
1:A:324:LYS:NZ	1:A:537:TRP:O	2.36	0.42
1:A:439:ALA:HB2	2:B:541:PRO:HB2	2.01	0.42
2:B:335:ASN:O	2:B:339:ILE:HG12	2.19	0.42
2:B:368:VAL:HG11	2:B:376:ILE:CD1	2.49	0.42
2:B:493:THR:O	2:B:493:THR:CG2	2.57	0.42
1:A:322:ILE:HD13	1:A:543:LEU:HD23	2.01	0.42
1:A:83:ASP:OD1	2:B:711:ILE:HG22	2.20	0.42
1:A:509:SER:OG	1:A:509:SER:O	2.36	0.42
1:A:12:MET:HG3	1:A:13:ILE:N	2.34	0.41
3:C:14:SER:OG	3:C:15:ARG:N	2.53	0.41
1:A:50:ASP:O	1:A:52:HIS:ND1	2.48	0.41
1:A:97:THR:OG1	2:B:728:ILE:HG23	2.20	0.41
1:A:123:THR:HB	1:A:130:TYR:CD2	2.56	0.41
2:B:241:ILE:HD12	4:V:18:A:C8	2.56	0.41
2:B:660:VAL:HG12	3:C:202:MET:HE1	2.01	0.41
1:A:245:LYS:HB3	2:B:472:LEU:HD21	2.02	0.41
2:B:134:ASN:O	2:B:134:ASN:ND2	2.52	0.41
1:A:41:HIS:NE2	1:A:120:ILE:O	2.53	0.41
2:B:213:ASN:OD1	2:B:213:ASN:N	2.53	0.41
2:B:248:ILE:O	2:B:248:ILE:HG22	2.20	0.41
1:A:559:ARG:HH11	1:A:559:ARG:HG3	1.87	0.40
1:A:583:ARG:NH1	2:B:512:PHE:O	2.50	0.40
2:B:251:PHE:CE1	2:B:339:ILE:HG21	2.56	0.40
1:A:376:PRO:O	2:B:365:ARG:NE	2.55	0.40

There are no symmetry-related clashes.



5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percer	ntiles
1	A	704/729~(97%)	692 (98%)	12 (2%)	0	100	100
2	В	741/757~(98%)	726 (98%)	15 (2%)	0	100	100
3	\mathbf{C}	$247/788 \ (31\%)$	244 (99%)	3 (1%)	0	100	100
6	G	21/42~(50%)	21 (100%)	0	0	100	100
All	All	1713/2316 (74%)	1683 (98%)	30 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percen	tiles
1	A	638/656 (97%)	613 (96%)	25 (4%)	27	59
2	В	658/667 (99%)	632 (96%)	26 (4%)	27	58
3	С	224/686 (33%)	212 (95%)	12 (5%)	18	46
6	G	23/37~(62%)	19 (83%)	4 (17%)	1	4
All	All	1543/2046 (75%)	1476 (96%)	67 (4%)	27	55

All (67) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	4	PHE
1	A	8	CYS



Continued from previous page...

	nued fron				
Mol	Chain	Res	Type		
1	A	15	GLU		
1	A	37	SER		
1	A	57	ARG		
1	A A	108	ASP		
1	A	130	TYR		
1	A	195	GLU		
1	A A A	225	SER		
1	A	245	LYS		
1	A	250	SER		
1	A A	262	ARG		
1	A	272	ASP		
1	A	383	ASP		
1	A A	391	LYS		
1	A A A	405	LYS CYS		
1	A	441	MET		
1	A	499	ASN		
1	A	512	ARG		
1	A	559	ARG		
1	A	589	LEU		
1	A	599	GLU		
1	A	603	LYS		
1	A	624	SER		
1	A	648	SER		
2	В	41	ASP		
2	В	116	GLN		
2	В	134	ASN		
2 2	В	180	GLU		
	В	187	ARG		
2 2	В	190	ARG		
2	В	246	MET		
2	В	284	ASN		
2	В	290	MET		
2	В	302	ILE		
2	В	307	THR		
2	В	356	MET		
2 2	В	404	SER		
2 2 2 2	В	506	SER		
2	В	592	SER		
2	В	610	CYS		
2	В	617	ASP		
2	В	626	ASN		
2	В	628	MET		



Continued from previous page...

Mol	Chain	Res	Type
2	В	633	SER
2	В	636	GLU
2	В	640	VAL
2	В	658	ASP
2	В	678	SER
2	В	727	ARG
2	В	754	ARG
3	С	36	SER
3	С	48	LYS
3	С	53	MET
3	С	73	GLN
3	С	79	SER
3	С	116	LYS
3	С	117	THR
3	С	146	ASP
3	С	153	ASP
3	C C C C C C G G G G	155	SER
3	С	192	GLU
3	С	217	PHE
6	G	4	SER
6	G	9	SER
6	G	11	SER
6	G	16	SER

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA (i)

Mol	Chain	Analysed	Backbone Outliers	Pucker Outliers
4	V	34/51 (66%)	9 (26%)	0
5	M	8/33 (24%)	0	0
All	All	42/84 (50%)	9 (21%)	0

All (9) RNA backbone outliers are listed below:

Mol	Chain	Res	Type
4	V	7	A
4	V	8	A
4	V	11	A
4	V	13	G



Continued from previous page...

Mol	Chain	Res	Type
4	V	14	G
4	V	28	С
4	V	29	U
4	V	47	U
4	V	51	U

There are no RNA pucker outliers to report.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

2 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mal	Type Chair		Chain Res	Dog Link	В	Bond lengths			Bond angles		
Mol	Type	Chain	nes	Link	Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2	
6	SEP	G	7	6	8,9,10	1.53	1 (12%)	8,12,14	1.59	2 (25%)	
6	SEP	G	21	6	8,9,10	1.51	1 (12%)	8,12,14	1.55	2 (25%)	

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
6	SEP	G	7	6	-	1/5/8/10	-
6	SEP	G	21	6	-	0/5/8/10	-

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	\mathbf{Z}	$\operatorname{Observed}(\mathring{\mathbf{A}})$	$\operatorname{Ideal}(\text{\AA})$
6	G	7	SEP	P-O1P	3.34	1.61	1.50
6	G	21	SEP	P-O1P	3.30	1.61	1.50

All (4) bond angle outliers are listed below:



Mol	Chain	Res	Type	Atoms	\mathbf{Z}	$Observed(^o)$	$\operatorname{Ideal}({}^{o})$
6	G	7	SEP	P-OG-CB	-3.00	110.02	118.30
6	G	21	SEP	P-OG-CB	-2.85	110.45	118.30
6	G	7	SEP	OG-CB-CA	2.80	110.87	108.14
6	G	21	SEP	OG-CB-CA	2.70	110.78	108.14

There are no chirality outliers.

All (1) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
6	G	7	SEP	N-CA-CB-OG

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
6	G	7	SEP	1	0

5.5 Carbohydrates (i)

There are no oligosaccharides in this entry.

5.6 Ligand geometry (i)

Of 3 ligands modelled in this entry, 2 are monoatomic - leaving 1 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
WIOI	Туре		rtes		Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
7	2KH	В	801	8	29,30,30	3.58	17 (58%)	42,47,47	1.53	7 (16%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.



Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
7	2KH	В	801	8	-	3/19/38/38	0/2/2/2

All (17) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	\mathbf{Z}	Observed(A)	$\operatorname{Ideal}(\text{\AA})$
7	В	801	2KH	C3'-C2'	-8.00	1.31	1.53
7	В	801	2KH	C2-N1	7.79	1.50	1.38
7	В	801	2KH	C2-N3	7.08	1.50	1.38
7	В	801	2KH	C6-C5	6.94	1.51	1.35
7	В	801	2KH	PB-O3B	4.24	1.64	1.59
7	В	801	2KH	O4'-C1'	-4.01	1.32	1.42
7	В	801	2KH	O4'-C4'	3.86	1.53	1.45
7	В	801	2KH	O3'-C3'	3.85	1.52	1.43
7	В	801	2KH	PB-O1B	-3.28	1.47	1.56
7	В	801	2KH	C5-C4	3.09	1.50	1.43
7	В	801	2KH	C4-N3	3.07	1.44	1.38
7	В	801	2KH	C5'-C4'	-2.63	1.43	1.51
7	В	801	2KH	C6-N1	2.61	1.44	1.38
7	В	801	2KH	PA-O5'	2.60	1.67	1.57
7	В	801	2KH	PB-N3A	2.59	1.70	1.63
7	В	801	2KH	C2'-C1'	2.34	1.61	1.53
7	В	801	2KH	O2-C2	-2.33	1.18	1.23

All (7) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	\mathbf{Z}	$Observed(^o)$	$\mathrm{Ideal}(^{o})$
7	В	801	2KH	C4-N3-C2	-4.63	120.47	126.58
7	В	801	2KH	N3-C2-N1	3.85	120.00	114.89
7	В	801	2KH	C5-C4-N3	3.26	119.72	114.84
7	В	801	2KH	PB-O3B-PG	-2.87	122.50	132.62
7	В	801	2KH	O4-C4-C5	-2.73	120.36	125.16
7	В	801	2KH	C3'-C2'-C1'	2.25	105.70	101.43
7	В	801	2KH	O2-C2-N1	-2.04	120.08	122.79

There are no chirality outliers.

All (3) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
7	В	801	2KH	PB-N3A-PA-O1A
7	В	801	2KH	C5'-O5'-PA-O2A
7	В	801	2KH	PA-N3A-PB-O2B



There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less then 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers (i)

There are no such residues in this entry.



5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.




6 Map visualisation (i)

This section contains visualisations of the EMDB entry EMD-17783. These allow visual inspection of the internal detail of the map and identification of artifacts.


Images derived from a raw map, generated by summing the deposited half-maps, are presented below the corresponding image components of the primary map to allow further visual inspection and comparison with those of the primary map.

6.1 Orthogonal projections (i)

6.1.1 Primary map



6.1.2 Raw map





The images above show the map projected in three orthogonal directions.




6.2 Central slices (i)

6.2.1 Primary map









X Index: 180


Y Index: 180

Z Index: 180

6.2.2 Raw map







X Index: 180

Y Index: 180


Z Index: 180

The images above show central slices of the map in three orthogonal directions.




6.3 Largest variance slices (i)


6.3.1 Primary map








Y Index: 173




Z Index: 183


6.3.2 Raw map



X Index: 172



Y Index: 173




Z Index: 183

The images above show the largest variance slices of the map in three orthogonal directions.




6.4 Orthogonal standard-deviation projections (False-color) (i)

6.4.1 Primary map



6.4.2 Raw map




The images above show the map standard deviation projections with false color in three orthogonal directions. Minimum values are shown in green, max in blue, and dark to light orange shades represent small to large values respectively.




6.5 Orthogonal surface views (i)

6.5.1 Primary map



The images above show the 3D surface view of the map at the recommended contour level 0.008. These images, in conjunction with the slice images, may facilitate assessment of whether an appropriate contour level has been provided.

6.5.2 Raw map



These images show the 3D surface of the raw map. The raw map's contour level was selected so that its surface encloses the same volume as the primary map does at its recommended contour level.




6.6 Mask visualisation (i)

This section shows the 3D surface view of the primary map at 50% transparency overlaid with the specified mask at 0% transparency

A mask typically either:

- Encompasses the whole structure
- Separates out a domain, a functional unit, a monomer or an area of interest from a larger structure


6.6.1 emd_17783_msk_1.map (i)



7 Map analysis (i)

This section contains the results of statistical analysis of the map.


7.1 Map-value distribution (i)



The map-value distribution is plotted in 128 intervals along the x-axis. The y-axis is logarithmic. A spike in this graph at zero usually indicates that the volume has been masked.



7.2 Volume estimate (i)




The volume at the recommended contour level is $73~\mathrm{nm}^3$; this corresponds to an approximate mass of $66~\mathrm{kDa}$.

The volume estimate graph shows how the enclosed volume varies with the contour level. The recommended contour level is shown as a vertical line and the intersection between the line and the curve gives the volume of the enclosed surface at the given level.



7.3 Rotationally averaged power spectrum (i)




^{*}Reported resolution corresponds to spatial frequency of 0.347 $\rm \mathring{A}^{-1}$



8 Fourier-Shell correlation (i)

Fourier-Shell Correlation (FSC) is the most commonly used method to estimate the resolution of single-particle and subtomogram-averaged maps. The shape of the curve depends on the imposed symmetry, mask and whether or not the two 3D reconstructions used were processed from a common reference. The reported resolution is shown as a black line. A curve is displayed for the half-bit criterion in addition to lines showing the 0.143 gold standard cut-off and 0.5 cut-off.

8.1 FSC (i)



*Reported resolution corresponds to spatial frequency of 0.347 $\mbox{Å}^{-1}$



8.2 Resolution estimates (i)

Resolution estimate (Å)	Estimation criterion (FSC cut-off)			
rtesolution estimate (A)	0.143	0.5	Half-bit	
Reported by author	2.88	-	-	
Author-provided FSC curve	2.87	3.39	2.93	
Unmasked-calculated*	3.78	7.97	3.91	


^{*}Resolution estimate based on FSC curve calculated by comparison of deposited half-maps. The value from deposited half-maps intersecting FSC 0.143 CUT-OFF 3.78 differs from the reported value 2.88 by more than 10 %



9 Map-model fit (i)

This section contains information regarding the fit between EMDB map EMD-17783 and PDB model 8PNQ. Per-residue inclusion information can be found in section 3 on page 7.


9.1 Map-model overlay (i)



The images above show the 3D surface view of the map at the recommended contour level 0.008 at 50% transparency in yellow overlaid with a ribbon representation of the model coloured in blue. These images allow for the visual assessment of the quality of fit between the atomic model and the map.




9.2 Q-score mapped to coordinate model (i)



The images above show the model with each residue coloured according its Q-score. This shows their resolvability in the map with higher Q-score values reflecting better resolvability. Please note: Q-score is calculating the resolvability of atoms, and thus high values are only expected at resolutions at which atoms can be resolved. Low Q-score values may therefore be expected for many entries.


9.3 Atom inclusion mapped to coordinate model (i)



The images above show the model with each residue coloured according to its atom inclusion. This shows to what extent they are inside the map at the recommended contour level (0.008).



9.4 Atom inclusion (i)



At the recommended contour level, 90% of all backbone atoms, 81% of all non-hydrogen atoms, are inside the map.



9.5 Map-model fit summary (i)

The table lists the average atom inclusion at the recommended contour level (0.008) and Q-score for the entire model and for each chain.

Chain	Atom inclusion	Q-score
All	0.8110	0.5860
A	0.8060	0.5850
В	0.8490	0.6000
С	0.7690	0.5780
G	0.6220	0.5180
M	0.9210	0.5730
V	0.6850	0.5270



