

Nov 10, 2024 - 07:25 am GMT

PDB ID	:	8RO0
EMDB ID	:	EMD-19397
Title	:	Structure of the C. elegans Intron Lariat Spliceosome primed for disassembly
		(ILS')
Authors	:	Vorlaender, M.K.; Rothe, P.; Plaschka, C.
Deposited on	:	2024-01-11
Resolution	:	2.90 Å(reported)
Based on initial model	:	

This is a Full wwPDB EM Validation Report for a publicly released PDB entry.

We welcome your comments at *validation@mail.wwpdb.org* A user guide is available at https://www.wwpdb.org/validation/2017/EMValidationReportHelp with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (1)) were used in the production of this report:

EMDB validation analysis	:	0.0.1.dev113
Mogul	:	1.8.4, CSD as541be (2020)
MolProbity	:	4.02b-467
buster-report	:	1.1.7(2018)
Percentile statistics	:	20231227.v01 (using entries in the PDB archive December 27th 2023)
MapQ	:	1.9.13
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.39

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: $ELECTRON\ MICROSCOPY$

The reported resolution of this entry is 2.90 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	$egin{array}{c} { m Whole \ archive} \ (\#{ m Entries}) \end{array}$	${f EM\ structures}\ (\#{ m Entries})$		
Ramachandran outliers	207382	16835		
Sidechain outliers	206894	16415		
RNA backbone	6643	2191		

The table below summarises the geometric issues observed across the polymeric chains and their fit to the map. The red, orange, yellow and green segments of the bar indicate the fraction of residues that contain outliers for $\geq=3, 2, 1$ and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq=5\%$ The upper red bar (where present) indicates the fraction of residues that have poor fit to the EM map (all-atom inclusion < 40%). The numeric value is given above the bar.

Mol	Chain	Length	(Quality of chain	
1	2	191	5% 28% •	68%	
2	5	121	• 54%	31%	7% • 8%
3	6	101	31% 48%	36%	6% 11%
4	А	2329	12%	93%	• •
5	В	2145	<u>.</u>	83%	• 16%
6	С	974	i i i i i i i i i i i i i i i i i i i	90%	• 8%
7	D	267	13% 32%	67%	
8	Е	331	<u>-</u>	93%	• 6%



Chain Length Quality of chain Mol i 9 Ι 855 82% 15% • 9% J 10 74475% 23% 12% 11 Κ 23885% 15% 23% 12L 75579% 19% . 16% 13М 23482% 16% • 14Ν 14795% .. . 8% 15Ο 40882% 16% • 38% Р 2301662% 35% . 24% 17РХ 809 58% 42% 7% 18Q 146793% • 6% 6% 19R 53549% 47% • 9% \mathbf{S} 2016998% i 21Т 49476% • 22% TF 22830 59% 40% . 26% W 2356784% 13% . 15% 24Ζ 16939% 59% 25136 \mathbf{a} 58% 40% • 10% 13625h 57% 40% • 26b 16059% 39% • 22% 26i 16048% 51% . 27127 \mathbf{c} 37% 61% 26% 27127j 63% 37% 10% 28d 11877% 20% • 29% 28k 11866% 33% • 2990 • 11% е 88%



Mol	Chain	Length		Quality o	of chain	
29	1	90	28%	86%		• 11%
30	f	85		84%		• 15%
30	m	85	16%	81%		• 15%
31	g	77	8%	97%		•
31	n	77	17%	96%		•
32	0	253	26%	63%	•	36%
33	р	217	35%		65%	
34	q	492	24%		75%	
34	r	492	26% •		73%	
34	s	492	55	5% 93%		• 5%
34	t	492	26%		74%	
35	У	331	12% 23%		76%	
36	In	19	21%	100%)	



2 Entry composition (i)

There are 40 unique types of molecules in this entry. The entry contains 121370 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

• Molecule 1 is a RNA chain called U2 snRNA.

Mol	Chain	Residues		At	\mathbf{oms}	AltConf	Trace		
1	2	61	Total 947	C 410	N 82	0 394	Р 61	0	0

• Molecule 2 is a RNA chain called U5 snRNA.

Mol	Chain	Residues		A	AltConf	Trace			
2	5	111	Total 2349	C 1052	N 405	0 781	Р 111	0	0

• Molecule 3 is a RNA chain called U6 snRNA.

Mol	Chain	Residues		A	toms	AltConf	Trace		
3	6	90	Total 1927	C 863	N 358	O 616	Р 90	0	0

• Molecule 4 is a protein called Pre-mRNA-splicing factor 8 homolog.

Mol	Chain	Residues		At	AltConf	Trace			
4	А	2231	Total 18444	C 11870	N 3216	O 3272	S 86	0	0

• Molecule 5 is a protein called U5 small nuclear ribonucleoprotein 200 kDa helicase.

Mol	Chain	Residues		A	AltConf	Trace			
5	В	1806	Total 14404	C 9175	N 2476	O 2669	S 84	0	0

• Molecule 6 is a protein called Tr-type G domain-containing protein.

Mol	Chain	Residues		Α	AltConf	Trace			
6	С	898	Total	С	N	0	S	0	0
, , , , , , , , , , , , , , , , , , ,	-		7152	4558	1211	1337	46		, i i i i i i i i i i i i i i i i i i i



• Molecule 7 is a protein called Protein isy-1.

Mol	Chain	Residues		At	oms	AltConf	Trace		
7	D	88	Total 726	C 458	N 122	0 143	${ m S} { m 3}$	0	0

• Molecule 8 is a protein called WD_REPEATS_REGION domain-containing protein.

Mol	Chain	Residues		At		AltConf	Trace		
8	Е	312	Total 2443	C 1528	N 429	0 466	S 20	0	0

• Molecule 9 is a protein called Pre-mRNA-splicing factor SYF1.

Mol	Chain	Residues		Α	toms			AltConf	Trace
9	Ι	724	Total	C 2012	N 1047	0	S 42	0	0
			5995	3813	1047	1092	43		

• Molecule 10 is a protein called TPR_REGION domain-containing protein.

Mol	Chain	Residues		At	oms			AltConf	Trace
10	J	574	Total 4893	C 3122	N 855	O 896	S 20	0	0

• Molecule 11 is a protein called Pre-mRNA-splicing factor SPF27.

Mol	Chain	Residues		At	oms			AltConf	Trace
11	K	203	Total 1665	C 1041	N 298	O 309	S 17	0	0

• Molecule 12 is a protein called Cell division cycle 5-like protein.

Mol	Chain	Residues		At	oms			AltConf	Trace
12	L	609	Total 4898	C 3039	N 902	0 934	S 23	0	0

• Molecule 13 is a protein called Pre-mRNA-splicing factor syf-2.

Mol	Chain	Residues		Ate	oms			AltConf	Trace
13	М	196	Total 1652	C 1021	N 308	0 317	S 6	0	0

• Molecule 14 is a protein called Protein BUD31 homolog.



Mol	Chain	Residues		\mathbf{A}	toms	AltConf	Trace		
14	Ν	142	Total 1163	С 731	N 212	O 208	S 12	0	0

• Molecule 15 is a protein called Pre-mRNA-splicing factor RBM22.

Mol	Chain	Residues		At		AltConf	Trace		
15	О	342	Total 2717	C 1703	N 493	O 502	S 19	0	0

• Molecule 16 is a protein called Spliceosome-associated protein CWC15 homolog.

Mol	Chain	Residues		At	oms	AltConf	Trace		
16	Р	150	Total 1204	C 729	N 232	0 237	S 6	0	0

• Molecule 17 is a protein called GCF C-terminal domain-containing protein.

Mol	Chain	Residues		At	oms			AltConf	Trace
17	РХ	472	Total 3838	C 2396	N 695	0 720	S 27	0	0

• Molecule 18 is a protein called Intron-binding protein aquarius.

Mol	Chain	Residues		Α	toms			AltConf	Trace
18	Q	1378	Total 11292	C 7218	N 1964	O 2063	S 47	0	0

• Molecule 19 is a protein called Uncharacterized protein T27F2.1.

Mol	Chain	Residues		At	oms			AltConf	Trace
19	R	282	Total 2207	C 1379	N 404	0 414	S 10	0	0

• Molecule 20 is a protein called Peptidyl-prolyl cis-trans isomerase.

Mol	Chain	Residues		At	oms	AltConf	Trace		
20	S	169	Total 1301	C 818	N 233	0 243	${f S}{7}$	0	0

• Molecule 21 is a protein called WD_REPEATS_REGION domain-containing protein.



Mol	Chain	Residues		At	AltConf	Trace			
21	Т	387	Total 3066	C 1937	N 555	O 555	S 19	0	0

• Molecule 22 is a protein called Septin and tuftelin-interacting protein 1 homolog.

Mol	Chain	Residues		At	AltConf	Trace			
22	TF	497	Total 4094	C 2637	N 685	O 750	S 22	0	0

• Molecule 23 is a protein called WD_REPEATS_REGION domain-containing protein.

Mol	Chain	Residues		At	AltConf	Trace			
23	W	496	Total 4071	C 2584	N 726	0 746	S 15	0	0

• Molecule 24 is a protein called Coiled-coil domain-containing protein 12.

Mol	Chain	Residues		At	oms	AltConf	Trace		
24	Z	69	Total 567	C 356	N 104	0 105	${S \over 2}$	0	0

• Molecule 25 is a protein called Small nuclear ribonucleoprotein Sm D3.

Mol	Chain	Residues		Atoms					Trace
25	0	81	Total	С	Ν	0	\mathbf{S}	0	0
2.0	a	01	634	396	113	119	6	0	0
25	h	81	Total	С	Ν	Ο	\mathbf{S}	0	0
	11	01	634	396	113	119	6		

• Molecule 26 is a protein called Probable small nuclear ribonucleoprotein-associated protein B.

Mol	Chain	Residues		At	oms		AltConf	Trace	
26	h	08	Total	С	Ν	0	\mathbf{S}	0	0
20	U	90	754	475	141	130	8	0	0
26	i	70	Total	С	Ν	0	\mathbf{S}	0	0
20	1	19	638	405	117	110	6		0

• Molecule 27 is a protein called Small nuclear ribonucleoprotein Sm D1.



Mol	Chain	Residues		At	oms		AltConf	Trace	
27	С	80	Total 621	C 396	N 109	0 112	$\frac{S}{4}$	0	0
27	j	80	Total 621	C 396	N 109	0 112	$\frac{S}{4}$	0	0

• Molecule 28 is a protein called Probable small nuclear ribonucleoprotein Sm D2.

Mol	Chain	Residues		At	oms		AltConf	Trace	
28	d	04	Total	С	Ν	Ο	S	0	0
20	u	94	747	469	135	138	5	0	0
<u> </u>	l.	70	Total	С	Ν	0	S	0	0
20	K	19	630	398	118	109	5	0	0

• Molecule 29 is a protein called Probable small nuclear ribonucleoprotein E.

Mol	Chain	Residues	Atoms					AltConf	Trace
29	е	80	Total 663	C 424	N 118	0 119	${S \over 2}$	0	0
29	1	80	Total 663	C 424	N 118	0 119	$\begin{array}{c} \mathrm{S} \\ \mathrm{2} \end{array}$	0	0

• Molecule 30 is a protein called Probable small nuclear ribonucleoprotein F.

Mol	Chain	Residues		At	oms		AltConf	Trace	
30	f	79	Total	С	Ν	Ο	\mathbf{S}	0	0
- 50	1	12	557	359	93	101	4	0	0
30	m	79	Total	С	Ν	0	S	0	0
- 50	111	12	557	359	93	101	4	0	0

• Molecule 31 is a protein called Probable small nuclear ribonucleoprotein G.

Mol	Chain	Residues		At	oms		AltConf	Trace	
21	a c	77	Total	С	Ν	0	S	0	0
	g	11	606	379	107	113	7	0	0
21	n	77	Total	С	Ν	0	S	0	0
	11	11	606	379	107	113	7	0	0

• Molecule 32 is a protein called Probable U2 small nuclear ribonucleoprotein A'.

Mol	Chain	Residues		At	oms			AltConf	Trace
32	О	162	Total 1335	C 849	N 236	0 243	${f S}{7}$	0	0



• Molecule 33 is a protein called RRM domain-containing protein.

Mol	Chain	Residues	Atoms				AltConf	Trace	
33	р	76	Total 626	C 402	N 114	O 106	${f S}$ 4	0	0

• Molecule 34 is a protein called Pre-mRNA-processing factor 19.

Mol	Chain	Residues	Atoms					AltConf	Trace
24	a	191	Total	С	Ν	0	S	0	0
34 q	q	121	939	585	165	184	5	0	0
24	r	121	Total	С	Ν	0	S	0	0
34	1	101	1003	621	179	198	5	0	
24	G	460	Total	С	Ν	0	S	0	0
34	s 409	409	3570	2239	620	702	9		
24	+	199	Total	С	Ν	0	S	0	0
34	t	t 128	991	620	173	193	5	U	U

• Molecule 35 is a protein called Peptidyl-prolyl cis-trans isomerase E.

Mol	Chain	Residues		At	oms			AltConf	Trace
35	У	79	Total 618	C 396	N 100	0 117	${f S}{5}$	0	0

• Molecule 36 is a DNA chain called Intron lariat RNA.

Mol	Chain	Residues		Ato	\mathbf{ms}		AltConf	Trace
36	In	19	Total 225	C 95	O 112	Р 18	0	0

• Molecule 37 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms	AltConf
37	6	6	Total Mg 6 6	0
37	С	1	Total Mg 1 1	0

• Molecule 38 is INOSITOL HEXAKISPHOSPHATE (three-letter code: IHP) (formula: $C_6H_{18}O_{24}P_6$).





Mol	Chain	Residues	Atoms				AltConf	
90	Δ	1	Total	С	Ο	Р	0	
30	A	1	36	6	24	6	0	
90	J	т	1	Total	С	Ο	Р	0
38		JI		6	24	6	0	

• Molecule 39 is GUANOSINE-5'-TRIPHOSPHATE (three-letter code: GTP) (formula: $C_{10}H_{16}N_5O_{14}P_3$).



Mol	Chain	Residues	Atoms				AltConf	
39	С	1	Total 32	C 10	N 5	0 14	Р 3	0



 $\bullet\,$ Molecule 40 is ZINC ION (three-letter code: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms	AltConf
40	Ν	3	Total Zn 3 3	0
40	О	3	Total Zn 3 3	0



3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and atom inclusion in map density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red diamond above a residue indicates a poor fit to the EM map for this residue (all-atom inclusion < 40%). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.



93%

Chain A:

















THR PRO LEU ALA

• Molecule 20: Peptidyl-prolyl cis-trans isomerase

Chain S:	9%	98%		
*****	• • •• •			
M1 P2 A3 P4 15 N6	D7 K42 G77 D78 D78 D82	NIO TIL		
• Molecule	21: WD_RI	EPEATS_REGION do	omain-containing pro	otein
Chain T:		76%	•	22%
ET LLA AL ER SP RD	HR C L C C C C C C C C C C C C C C C C C		. ↓ ↓ ↓ ↓ ↓ ↓ ↓ ↓ ↓ ↓ ↓ ↓ ↓ ↓ ↓ ↓ ↓ ↓ ↓	E E E E E E E E E E E E E E E E E E E

• Molecule 22: Septin and tuftelin-interacting protein 1 homolog







 \bullet Molecule 25: Small nuclear ribonucleoprotein Sm D3





• Molecule 28: Probable small nuclear ribonucleoprotein Sm D2



_	10%			
Chain d:	77%		20%	
MET SER ALA GLN ALA LYS PRO ARG	SER GLU MET A13 A13 A17 A17 A17 A17 A18 A18 A18 A18 A18 A18 A18 A18 A18 A18	E76 VAL VAL LYS CLYS CLY CLYS CLYS CLYS CLYS CLYS	L114 A115 Q116 A117 E118	
• Molecule	28: Probable small nuclear rib	oonucleoprotein Sm	D2	
Chain lu	29%		220/	
Chain K:	66%	•	33%	
MET SER ALA GLN ALA LYS PRO ARG	SER MET THR MET THR ALA ALA ALA ALA ALA ALA ALA ALA ALA AL	V42 V42 143 143 N45 N45 N49 N49 K50 K51 L52 L53	N64 M65 V66 L67 E68 E68 K71	M73 W74 W75 E76 VAL PR0 LV8 LV8 THR CLY
LYS GLY LYS LYS ALA SER SER	V90 A91 K92 K92 F93 F95 F95 K98 F95 F95 F95 G03 F100 F100 F100 F100 V109	A117		
• Molecule	29: Probable small nuclear rib	oonucleoprotein E		
Chain e:	88%		• 11%	
MET SER THR ARG LYS LEU ASN LYS	MILT MET M11 C R R R R R R R R R R R R R R R R R R			
• Molecule	29: Probable small nuclear rib	oonucleoprotein E		
	28%			
Chain I:	86%		• 11%	
MET SER THR ARG LYS LYS LEU ASN LYS	VAL WAL V11 012 117 117 133 133 133 133 133 133 134 143 143 143	A57 A57 M61 M62 K63 K65 K65 K65 C66	472 671 885 485 485 486 487 988 988	
• Molecule	30: Probable small nuclear rib	oonucleoprotein F		
Chain f:	84%		• 15%	
MET SER ALA VAL Q5 M29 M29	676 VAL ASP ALV ASU ALU ASU ALA ALA			
• Molecule	30: Probable small nuclear rib	oonucleoprotein F		
Chain m:	16% 81%		• 15%	I
MET SER ALA VAL Q5 V21	L25 M29 F33 F33 F33 F33 F33 F33 F33 F33 F33 F3	R67 C68 C68 C68 C68 C78 C77 C75 C75 C75 C75 C75 C75 C75 C75 C75	GLU THR SER ALA	
• Molecule	31: Probable small nuclear rib	oonucleoprotein G		
Cha:	8%			
Unam g:	97	%	•	
		PROTEIN DATA BANK		



• Molecule 31: Probable small nuclear ribonucleoprotein G















4 Experimental information (i)

Property	Value	Source
EM reconstruction method	SINGLE PARTICLE	Depositor
Imposed symmetry	POINT, Not provided	
Number of particles used	879523	Depositor
Resolution determination method	FSC 0.143 CUT-OFF	Depositor
CTF correction method	PHASE FLIPPING AND AMPLITUDE	Depositor
	CORRECTION	
Microscope	FEI TITAN KRIOS	Depositor
Voltage (kV)	300	Depositor
Electron dose $(e^-/\text{\AA}^2)$	60	Depositor
Minimum defocus (nm)	700	Depositor
Maximum defocus (nm)	2000	Depositor
Magnification	Not provided	
Image detector	GATAN K3 BIOQUANTUM (6k x 4k)	Depositor
Maximum map value	8.985	Depositor
Minimum map value	0.000	Depositor
Average map value	0.009	Depositor
Map value standard deviation	0.094	Depositor
Recommended contour level	0.5	Depositor
Map size (Å)	582.9824, 582.9824, 582.9824	wwPDB
Map dimensions	448, 448, 448	wwPDB
Map angles ($^{\circ}$)	90.0, 90.0, 90.0	wwPDB
Pixel spacing (Å)	1.3013, 1.3013, 1.3013	Depositor



5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: GTP, IHP, ZN, MG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mal	Chain	B	ond lengths	Bond angles	
	Chain	RMSZ	# Z > 5	RMSZ	# Z > 5
1	2	0.56	0/558	1.16	1/863~(0.1%)
2	5	1.19	5/2622~(0.2%)	1.47	28/4077~(0.7%)
3	6	0.54	0/2159	1.21	7/3363~(0.2%)
4	А	0.50	2/18943~(0.0%)	0.70	1/25695~(0.0%)
5	В	0.31	0/14697	0.61	1/19895~(0.0%)
6	С	0.60	1/7309~(0.0%)	0.72	1/9906~(0.0%)
7	D	0.37	0/734	0.70	0/984
8	Ε	0.44	0/2500	0.71	2/3384~(0.1%)
9	Ι	0.45	0/6133	0.68	0/8286
10	J	0.45	0/5006	0.68	0/6735
11	Κ	0.33	0/1688	0.62	0/2260
12	L	0.32	0/4970	0.68	1/6673~(0.0%)
13	М	0.35	0/1678	0.67	0/2238
14	Ν	0.57	1/1190~(0.1%)	0.70	0/1597
15	0	0.49	0/2779	0.71	0/3764
16	Р	0.33	0/1220	0.74	0/1624
17	РХ	0.34	0/3893	0.71	0/5223
18	Q	0.41	0/11554	0.65	1/15626~(0.0%)
19	R	0.40	0/2252	0.74	1/3040~(0.0%)
20	S	0.33	0/1330	0.70	0/1800
21	Т	0.51	1/3144~(0.0%)	0.75	0/4259
22	TF	0.53	0/4202	0.72	4/5711~(0.1%)
23	W	0.41	0/4179	0.77	0/5638
24	Ζ	0.27	0/571	0.66	0/765
25	a	0.44	0/642	0.76	0/864
25	h	0.37	0/642	0.76	0/864
26	b	0.42	0/766	0.76	$0/1\overline{021}$
26	i	0.34	$0/\overline{648}$	0.75	$0/\overline{865}$
27	с	0.36	0/627	0.77	1/848~(0.1%)
27	j	0.34	$0/\overline{627}$	0.69	0/848
28	d	0.36	0/755	0.69	0/1013
28	k	0.37	$0/\overline{637}$	0.79	0/854



Mal	Chain	В	ond lengths	E	Bond angles
	Chain	RMSZ	# Z > 5	RMSZ	# Z > 5
29	е	0.39	0/674	0.70	0/909
29	1	0.36	0/674	0.77	0/909
30	f	0.41	0/568	0.67	0/769
30	m	0.42	0/568	0.79	0/769
31	g	0.39	0/614	0.76	1/820~(0.1%)
31	n	0.36	0/614	0.73	0/820
32	0	0.33	0/1358	0.70	0/1837
33	р	0.37	0/638	0.72	0/850
34	q	0.32	0/951	0.65	0/1282
34	r	0.30	0/1017	0.67	0/1373
34	s	0.32	0/3632	0.67	0/4913
34	t	0.30	0/1004	0.62	0/1355
35	У	0.42	0/630	0.76	0/846
All	All	0.46	10/123597~(0.0%)	0.73	50/168035~(0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
2	5	0	1
4	А	0	5
5	В	0	3
6	С	0	1
9	Ι	0	1
10	J	0	2
17	PX	0	1
19	R	0	1
22	TF	0	1
26	b	0	1
34	q	0	1
All	All	0	18

All (10) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	5	26	С	N1-C6	-7.06	1.32	1.37
21	Т	286	CYS	CB-SG	-6.26	1.71	1.82
14	Ν	142	CYS	CB-SG	-5.94	1.72	1.81
6	С	552	CYS	CB-SG	-5.92	1.72	1.81
2	5	60	G	C6-N1	-5.78	1.35	1.39



Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	5	15	C	N1-C6	-5.27	1.33	1.37
4	А	198	TYR	CD1-CE1	-5.15	1.31	1.39
2	5	26	С	C5-C6	-5.14	1.30	1.34
4	А	214	TYR	CE1-CZ	-5.09	1.31	1.38
2	5	13	С	C4-C5	-5.01	1.39	1.43

All (50) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	$Observed(^{o})$	$Ideal(^{o})$
2	5	109	C	C6-N1-C2	-11.45	115.72	120.30
2	5	5	U	C5-C4-O4	-11.34	119.10	125.90
2	5	5	U	N3-C4-O4	11.19	127.23	119.40
2	5	27	G	O4'-C1'-N9	-10.79	99.57	108.20
2	5	25	С	C6-N1-C2	-10.04	116.28	120.30
3	6	89	С	C6-N1-C2	-7.98	117.11	120.30
2	5	115	А	O4'-C1'-N9	7.56	114.25	108.20
2	5	26	С	C4-C5-C6	7.44	121.12	117.40
2	5	13	С	C6-N1-C2	-7.37	117.35	120.30
19	R	292	ASP	CB-CG-OD1	7.21	124.79	118.30
22	TF	392	ARG	C-N-CA	6.92	139.00	121.70
2	5	5	U	C6-N1-C2	-6.81	116.91	121.00
2	5	41	С	C6-N1-C2	-6.80	117.58	120.30
3	6	37	С	C6-N1-C2	-6.70	117.62	120.30
3	6	89	С	C2-N1-C1'	6.70	126.16	118.80
2	5	109	C	O4'-C1'-N1	6.69	113.55	108.20
4	А	511	ASP	CB-CG-OD1	6.68	124.32	118.30
27	с	33	ASP	CB-CG-OD1	6.64	124.27	118.30
2	5	18	С	C6-N1-C2	-6.21	117.82	120.30
12	L	361	LEU	CA-CB-CG	6.18	129.51	115.30
22	TF	360	ASP	CB-CG-OD1	6.18	123.86	118.30
2	5	25	C	C2'-C3'-O3'	6.08	123.42	113.70
2	5	82	U	C6-N1-C2	-6.04	117.38	121.00
2	5	35	С	C6-N1-C2	-5.94	117.93	120.30
31	g	35	ASP	CB-CG-OD1	5.91	123.62	118.30
1	2	7	U	C5-C6-N1	5.89	125.64	122.70
2	5	114	G	O4'-C1'-N9	5.86	112.89	108.20
5	В	1150	ASP	CB-CG-OD1	5.85	123.57	118.30
22	TF	334	ASP	CB-CG-OD1	5.82	123.54	118.30
2	5	33	A	C8-N9-C4	-5.78	103.49	105.80
2	5	116	G	O4'-C1'-N9	5.69	112.75	108.20
3	6	8	C	C6-N1-C2	-5.69	118.03	120.30
18	Q	364	ASP	CB-CG-OD1	5.67	123.40	118.30



Mol	Chain	Res	Type	Atoms	Z	$Observed(^{o})$	$Ideal(^{o})$
2	5	109	С	C5-C6-N1	5.57	123.78	121.00
2	5	80	А	O4'-C1'-N9	5.56	112.65	108.20
3	6	81	U	P-O3'-C3'	5.53	126.33	119.70
2	5	100	G	C1'-O4'-C4'	-5.50	105.50	109.90
2	5	5	U	O4'-C1'-N1	5.47	112.57	108.20
2	5	13	С	C5-C6-N1	5.41	123.70	121.00
8	Е	87	LEU	CA-CB-CG	5.41	127.73	115.30
3	6	74	С	C6-N1-C2	-5.24	118.20	120.30
2	5	15	С	C5-C6-N1	5.24	123.62	121.00
2	5	25	С	C5-C6-N1	5.24	123.62	121.00
8	Е	228	ASP	CB-CG-OD1	5.22	123.00	118.30
3	6	28	G	P-O3'-C3'	5.16	125.89	119.70
2	5	33	А	N7-C8-N9	5.15	116.38	113.80
2	5	94	U	O4'-C1'-N1	5.14	112.31	108.20
6	С	233	ASP	CB-CG-OD1	5.05	122.84	118.30
2	5	9	G	C8-N9-C1'	-5.03	120.46	127.00
22	TF	786	ARG	NE-CZ-NH1	5.01	122.81	120.30

There are no chirality outliers.

All (18)	planarity	outliers	are	listed	below:
------------	-----------	----------	-----	--------	--------

Mol	Chain	Res	Type	Group
2	5	109	С	Sidechain
4	А	1409	ILE	Peptide
4	А	1432	ARG	Sidechain
4	А	229	ARG	Sidechain
4	А	655	ARG	Sidechain
4	А	933	LYS	Peptide
5	В	1513	ARG	Sidechain
5	В	1692	ARG	Sidechain
5	В	857	ARG	Sidechain
6	С	162	ARG	Sidechain
9	Ι	689	ARG	Sidechain
10	J	152	ARG	Sidechain
10	J	174	ARG	Sidechain
17	PX	469	ARG	Sidechain
19	R	190	ASN	Peptide
22	TF	392	ARG	Peptide
26	b	25	ARG	Sidechain
34	q	28	ARG	Sidechain



5.2 Too-close contacts (i)

Due to software issues we are unable to calculate clashes - this section is therefore empty.

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Perce	entiles
4	А	2223/2329~(95%)	2118 (95%)	91 (4%)	14 (1%)	22	52
5	В	1802/2145~(84%)	1780 (99%)	20 (1%)	2 (0%)	48	77
6	С	896/974~(92%)	857 (96%)	36 (4%)	3~(0%)	37	66
7	D	84/267~(32%)	82 (98%)	2 (2%)	0	100	100
8	Е	310/331~(94%)	294 (95%)	15 (5%)	1 (0%)	37	66
9	Ι	722/855~(84%)	702 (97%)	19 (3%)	1 (0%)	48	77
10	J	570/744 (77%)	547 (96%)	23 (4%)	0	100	100
11	К	201/238 (84%)	195 (97%)	5 (2%)	1 (0%)	25	56
12	L	599/755~(79%)	562 (94%)	33 (6%)	4 (1%)	19	49
13	М	190/234 (81%)	185 (97%)	5 (3%)	0	100	100
14	N	140/147~(95%)	130 (93%)	10 (7%)	0	100	100
15	Ο	334/408~(82%)	317 (95%)	15 (4%)	2 (1%)	22	52
16	Р	146/230~(64%)	137 (94%)	5 (3%)	4 (3%)	4	17
17	PX	470/809~(58%)	464 (99%)	4 (1%)	2 (0%)	30	60
18	Q	1376/1467~(94%)	1330 (97%)	44 (3%)	2 (0%)	48	77
19	R	274/535~(51%)	249 (91%)	23 (8%)	2 (1%)	19	49
20	S	167/169~(99%)	156 (93%)	10 (6%)	1 (1%)	22	52
21	Т	383/494~(78%)	361 (94%)	20 (5%)	2 (0%)	25	56
22	TF	491/830~(59%)	473 (96%)	16 (3%)	2(0%)	30	60
23	W	490/567~(86%)	445 (91%)	40 (8%)	5 (1%)	13	40
24	Z	67/169~(40%)	62 (92%)	5 (8%)	0	100	100



Mol	Chain	Analysed	Favoured	Allowed	Outliers	Perce	entiles
25	a	79/136~(58%)	78~(99%)	1 (1%)	0	100	100
25	h	79/136~(58%)	68~(86%)	9 (11%)	2 (2%)	4	18
26	b	92/160~(58%)	83 (90%)	8 (9%)	1 (1%)	12	37
26	i	75/160~(47%)	69~(92%)	5 (7%)	1 (1%)	10	33
27	с	78/127~(61%)	76~(97%)	2 (3%)	0	100	100
27	j	78/127~(61%)	72 (92%)	6 (8%)	0	100	100
28	d	90/118 (76%)	84 (93%)	6 (7%)	0	100	100
28	k	75/118~(64%)	72 (96%)	3 (4%)	0	100	100
29	е	78/90~(87%)	75~(96%)	2 (3%)	1 (1%)	10	33
29	1	78/90~(87%)	73 (94%)	3 (4%)	2 (3%)	4	17
30	f	70/85~(82%)	68 (97%)	2 (3%)	0	100	100
30	m	70/85~(82%)	66 (94%)	3 (4%)	1 (1%)	9	31
31	g	75/77~(97%)	71 (95%)	4 (5%)	0	100	100
31	n	75/77~(97%)	68 (91%)	5 (7%)	2 (3%)	4	17
32	0	160/253~(63%)	154 (96%)	5 (3%)	1 (1%)	22	52
33	р	74/217~(34%)	70 (95%)	4 (5%)	0	100	100
34	q	117/492~(24%)	112 (96%)	5 (4%)	0	100	100
34	r	129/492~(26%)	125 (97%)	4 (3%)	0	100	100
34	S	465/492~(94%)	443 (95%)	20 (4%)	2 (0%)	30	60
34	t	124/492~(25%)	119 (96%)	5 (4%)	0	100	100
35	У	77/331 (23%)	73 (95%)	4 (5%)	0	100	100
All	All	14173/19052 (74%)	13565 (96%)	547 (4%)	61 (0%)	32	60

All (61) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
4	А	934	PRO
6	С	97	VAL
12	L	350	LEU
12	L	507	ALA
16	Р	57	LYS
16	Р	58	ASP
18	Q	349	ASN
22	TF	392	ARG
22	TF	393	SER



Mol	Chain	Res	Type
23	W	508	GLN
4	А	1084	ILE
4	А	1542	VAL
4	А	2095	TYR
4	А	2275	ASN
17	РХ	411	VAL
29	1	72	ARG
4	А	1243	SER
4	А	2200	ASP
4	А	2285	MET
12	L	353	THR
15	0	295	PHE
16	Р	59	LEU
19	R	113	VAL
25	h	46	GLU
31	n	59	MET
4	А	1881	GLU
6	С	95	PRO
6	С	523	GLU
8	Е	120	ARG
9	Ι	242	ILE
11	K	71	PHE
15	0	303	LYS
21	Т	483	VAL
23	W	317	GLY
29	е	72	ARG
25	h	54	GLY
26	i	4	SER
4	А	454	ARG
5	В	2045	ILE
12	L	352	ASP
17	PX	482	ARG
18	Q	774	ILE
20	S	78	ASP
23	W	360	GLY
23	W	484	PHE
31	n	68	VAL
34	S	171	LYS
5	В	1664	ASP
16	Р	48	GLN
34	S	442	GLY
4	А	2303	ARG



Mol	Chain	Res	Type
23	W	531	VAL
4	А	2167	PRO
30	m	74	VAL
4	А	620	GLY
19	R	117	VAL
21	Т	224	GLY
29	1	44	ILE
32	0	32	PRO
4	А	417	PRO
26	b	100	GLY

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent side chain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Perce	\mathbf{ntiles}
4	А	2018/2100~(96%)	1972~(98%)	46 (2%)	45	77
5	В	1593/1886~(84%)	1583~(99%)	10 (1%)	84	95
6	\mathbf{C}	793/861~(92%)	780~(98%)	13 (2%)	58	84
7	D	80/230~(35%)	77~(96%)	3(4%)	28	63
8	Ε	275/291~(94%)	273~(99%)	2(1%)	81	94
9	Ι	641/754~(85%)	624~(97%)	17 (3%)	40	73
10	J	508/650~(78%)	494 (97%)	14 (3%)	38	73
11	Κ	183/214~(86%)	183 (100%)	0	100	100
12	L	529/645~(82%)	517 (98%)	12 (2%)	45	77
13	М	178/212~(84%)	175~(98%)	3 (2%)	56	83
14	Ν	125/129~(97%)	122 (98%)	3 (2%)	44	76
15	О	295/351~(84%)	291 (99%)	4 (1%)	62	86
16	Р	129/197~(66%)	125 (97%)	4 (3%)	35	70
17	PX	423/724 (58%)	420 (99%)	3 (1%)	81	94
18	Q	1235/1311~(94%)	1223 (99%)	12 (1%)	73	91
19	R	235/447~(53%)	221 (94%)	14 (6%)	16	44



Mol	Chain	Analysed	Rotameric	Outliers	Perce	ntiles
20	\mathbf{S}	137/137~(100%)	134~(98%)	3~(2%)	47	78
21	Т	333/421~(79%)	322~(97%)	11 (3%)	33	68
22	TF	451/718~(63%)	444 (98%)	7(2%)	58	84
23	W	444/502~(88%)	429~(97%)	15 (3%)	32	67
24	Ζ	63/145~(43%)	60~(95%)	3~(5%)	21	54
25	a	71/106~(67%)	69~(97%)	2(3%)	38	73
25	h	71/106~(67%)	70~(99%)	1 (1%)	62	86
26	b	79/116~(68%)	78~(99%)	1 (1%)	65	88
26	i	70/116~(60%)	68~(97%)	2(3%)	37	72
27	с	73/98 (74%)	71~(97%)	2(3%)	40	73
27	j	73/98 (74%)	73 (100%)	0	100	100
28	d	84/103 (82%)	81 (96%)	3 (4%)	30	65
28	k	72/103~(70%)	71~(99%)	1 (1%)	62	86
29	е	71/81~(88%)	71~(100%)	0	100	100
29	1	71/81~(88%)	70~(99%)	1 (1%)	62	86
30	f	61/71~(86%)	60~(98%)	1 (2%)	58	84
30	m	61/71~(86%)	59~(97%)	2(3%)	33	68
31	g	69/69~(100%)	68~(99%)	1 (1%)	62	86
31	n	69/69~(100%)	68~(99%)	1 (1%)	62	86
32	0	151/225~(67%)	150 (99%)	1 (1%)	81	94
33	р	68/192~(35%)	67~(98%)	1 (2%)	60	85
34	q	108/417~(26%)	107~(99%)	1 (1%)	75	92
34	r	114/417~(27%)	109~(96%)	5 (4%)	24	57
34	S	$\overline{396/417}\ (95\%)$	$\overline{389}\ (98\%)$	7 (2%)	54	82
34	t	115/417 (28%)	113 (98%)	2 (2%)	56	83
35	У	64/270~(24%)	62~(97%)	2 (3%)	35	70
All	All	12679/16568~(76%)	12443 (98%)	236 (2%)	52	81

All (236) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
4	А	87	HIS
4	А	103	ILE
	a	7	

Mol	Chain	Res	Type
4	А	105	ASP
4	А	178	LEU
4	А	311	LEU
4	А	443	LEU
4	А	519	VAL
4	А	560	ASN
4	А	743	THR
4	А	781	GLU
4	А	934	PRO
4	А	961	SER
4	А	969	MET
4	А	987	ARG
4	А	1012	LYS
4	А	1138	ASP
4	А	1174	ASN
4	А	1177	MET
4	А	1203	ASP
4	А	1274	GLN
4	А	1347	ARG
4	А	1361	LEU
4	А	1369	GLU
4	А	1377	ARG
4	А	1397	THR
4	А	1542	VAL
4	А	1564	ILE
4	А	1627	GLN
4	А	1671	ARG
4	А	1715	MET
4	А	1997	GLN
4	А	2106	LYS
4	А	2131	GLN
4	А	2133	LYS
4	А	2149	GLN
4	Α	2168	LEU
4	А	2170	TRP
4	A	2171	MET
4	А	2177	GLU
4	A	2178	LEU
4	А	2182	SER
4	А	2185	ASP
4	А	2189	HIS
4	А	2245	MET



Mol	Chain	Res	Type
4	А	2258	ASP
4	А	2276	PHE
5	В	162	ASP
5	В	513	GLN
5	В	534	TYR
5	В	720	SER
5	В	878	GLN
5	В	1070	LEU
5	В	1461	PHE
5	В	1609	ARG
5	В	2012	HIS
5	В	2084	LEU
6	С	76	TYR
6	С	94	GLU
6	C	206	ILE
6	С	243	GLU
6	С	290	ASN
6	С	326	PHE
6	С	381	MET
6	С	392	VAL
6	С	417	ARG
6	С	667	ASN
6	С	760	LYS
6	С	802	THR
6	С	870	SER
7	D	155	ARG
7	D	243	LEU
7	D	247	GLN
8	E	121	THR
8	E	314	MET
9	I	49	HIS
9	I	54	LYS
9	I	135	GLU
9	I	176	TYR
9	I	320	GLU
9	I	323	GLN
9	I	336	MET
9	Ι	351	MET
9	I	372	ILE
9	Ι	388	VAL
9	Ι	395	ILE
9	I	399	LYS



Mol	Chain	Res	Type
9	Ι	414	ASN
9	Ι	467	VAL
9	Ι	663	LEU
9	Ι	681	THR
9	Ι	735	GLU
10	J	34	THR
10	J	36	GLU
10	J	37	GLN
10	J	83	ARG
10	J	89	TRP
10	J	224	TRP
10	J	254	GLU
10	J	297	PHE
10	J	302	GLN
10	J	458	TYR
10	J	545	TYR
10	J	583	GLN
10	J	615	ARG
10	J	617	GLU
12	L	13	ASN
12	L	52	GLU
12	L	234	TYR
12	L	316	ARG
12	L	351	LYS
12	L	355	GLN
12	L	361	LEU
12	L	436	ARG
12	L	452	GLN
12	L	504	GLU
12	L	508	LYS
12	L	585	LYS
13	М	55	LYS
13	М	63	ARG
13	М	206	ASN
14	Ν	19	ASP
14	Ν	48	ILE
14	N	142	CYS
15	0	27	THR
15	0	78	VAL
15	Ο	115	ARG
15	Ο	181	GLU
16	Р	31	SER



Mol	Chain	Res	Type
16	Р	33	LYS
16	Р	59	LEU
16	Р	89	ARG
17	PX	396	LEU
17	PX	406	CYS
17	PX	443	GLN
18	Q	237	GLN
18	Q	318	ASP
18	Q	357	VAL
18	Q	602	LEU
18	Q	780	ARG
18	Q	825	VAL
18	Q	979	LYS
18	Q	1117	MET
18	Q	1118	GLU
18	Q	1285	HIS
18	Q	1291	ARG
18	Q	1308	ARG
19	R	6	ARG
19	R	16	ASP
19	R	53	PHE
19	R	91	THR
19	R	130	TRP
19	R	200	ILE
19	R	202	MET
19	R	210	MET
19	R	241	THR
19	R	243	LYS
19	R	259	ASN
19	R	267	LEU
19	R	276	ARG
19	R	298	ASP
20	S	100	ASN
20	S	117	THR
20	S	158	LEU
21	Т	85	LYS
21	Т	190	VAL
21	Т	220	LEU
21	Т	322	ASP
21	Т	343	ARG
21	Т	345	MET
21	Т	392	HIS



Mol	Chain	Res	Type
21	Т	397	ASN
21	Т	435	GLN
21	Т	451	ASP
21	Т	473	ASP
22	TF	326	ILE
22	TF	341	ARG
22	TF	360	ASP
22	TF	449	GLU
22	TF	548	MET
22	TF	550	ASP
22	TF	638	GLU
23	W	58	LYS
23	W	158	LYS
23	W	282	GLN
23	W	294	CYS
23	W	299	LYS
23	W	312	VAL
23	W	343	VAL
23	W	354	LYS
23	W	369	HIS
23	W	385	LYS
23	W	410	SER
23	W	417	ASN
23	W	434	TRP
23	W	454	MET
23	W	504	PHE
24	Ζ	120	ASP
24	Ζ	135	GLU
24	Ζ	142	VAL
25	a	46	GLU
25	a	77	MET
26	b	38	MET
27	с	72	ASP
27	с	77	ASP
28	d	48	ASN
28	d	65	MET
28	d	73	MET
30	f	29	MET
31	g	50	LYS
25	h	77	MET
26	i	35	ASP
26	i	88	ARG



Mol	Chain	Res	Type
28	k	49	ASN
29	1	54	PHE
30	m	72	LEU
30	m	73	TYR
31	n	32	ARG
32	0	106	ILE
33	р	34	GLN
34	q	129	ARG
34	r	28	ARG
34	r	32	LYS
34	r	46	GLU
34	r	115	CYS
34	r	121	LEU
34	S	199	GLN
34	S	289	ILE
34	S	306	ASP
34	S	342	HIS
34	S	352	ILE
34	s	395	LEU
34	s	455	GLU
34	t	36	GLU
34	t	110	GLN
35	У	33	PRO
35	у	36	ASP

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (7) such sidechains are listed below:

Mol	Chain	Res	Type
4	А	1997	GLN
5	В	681	GLN
17	РХ	394	GLN
22	TF	346	GLN
22	TF	532	GLN
22	TF	819	ASN
30	f	70	ASN

5.3.3 RNA (i)

Mol	Chain	Analysed	Backbone Outliers	Pucker Outliers
1	2	22/191~(11%)	6 (27%)	0



Mol	Chain	Analysed	Backbone Outliers	Pucker Outliers
2	5	110/121 (90%)	28 (25%)	13 (11%)
3	6	89/101 (88%)	39~(43%)	10 (11%)
All	All	221/413~(53%)	73~(33%)	23 (10%)

All (73) RNA backbone outliers are listed below:

Mol	Chain	Res	Type
1	2	10	U
1	2	19	U
1	2	20	А
1	2	21	G
1	2	25	А
1	2	26	G
2	5	4	С
2	5	6	С
2	5	8	G
2	5	10	U
2	5	20	U
2	5	23	A
2	5	24	А
2	5	26	С
2	5	27	G
2	5	28	U
2	5	43	U
2	5	72	А
2	5	74	А
2	5	76	А
2	5	77	U
2	5	79	G
2	5	80	А
2	5	81	G
2	5	82	U
2	5	96	U
2	5	99	G
2	5	100	G
2	5	101	А
2	5	110	G
2	5	112	А
2	5	113	А
2	5	114	G
2	5	117	С
3	6	4	С



Mol	Chain	Res	Type
3	6	5	U
3	6	6	U
3	6	7	С
3	6	8	С
3	6	9	G
3	6	10	А
3	6	20	С
3	6	21	U
3	6	22	А
3	6	23	А
3	6	24	А
3	6	29	G
3	6	32	С
3	6	34	А
3	6	37	С
3	6	39	G
3	6	41	G
3	6	43	А
3	6	44	G
3	6	46	U
3	6	48	А
3	6	49	G
3	6	54	G
3	6	55	С
3	6	56	С
3	6	62	G
3	6	64	А
3	6	69	U
3	6	74	С
3	6	75	G
3	6	79	A
3	6	80	U
3	6	81	U
3	6	82	С
3	6	84	U
3	6	85	G
3	6	86	А
3	6	89	С

Continued from previous page...

All (23) RNA pucker outliers are listed below: Continued on next page...



Mol	Chain	Res	Type
Mol	Chain	\mathbf{Res}	Type
2	5	3	A
2	5	5	U
2	5	19	А
2	5	23	А
2	5	25	С
2	5	75	С
2	5	79	G
2	5	81	G
2	5	100	G
2	5	109	С
2	5	111	А
2	5	112	А
2	5	113	А
3	6	23	А
3	6	28	G
3	6	37	С
3	6	41	G
3	6	45	А
3	6	48	А
3	6	53	G
3	6	79	А
3	6	81	U
3	6	85	G

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

There are no oligosaccharides in this entry.

5.6 Ligand geometry (i)

Of 16 ligands modelled in this entry, 13 are monoatomic - leaving 3 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The



Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mal	Tuno	Chain	Dog	Tink	B	Bond lengths			Bond angles		
WIOI	туре	Unain	nes		Counts	RMSZ	# Z >2	Counts	RMSZ	# Z > 2	
38	IHP	J	3000	-	36,36,36	2.23	10 (27%)	54,60,60	0.89	1 (1%)	
39	GTP	С	1101	37	26,34,34	1.89	5 (19%)	32,54,54	1.82	6 (18%)	
38	IHP	А	3000	-	36,36,36	2.22	10 (27%)	54,60,60	1.44	7 (12%)	

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
38	IHP	J	3000	-	-	0/30/54/54	0/1/1/1
39	GTP	С	1101	37	-	0/18/38/38	0/3/3/3
38	IHP	А	3000	-	-	4/30/54/54	0/1/1/1

All (25) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	$\mathrm{Ideal}(\mathrm{\AA})$
39	С	1101	GTP	C5-C6	-6.23	1.34	1.47
38	J	3000	IHP	C3-C2	-5.31	1.41	1.52
38	J	3000	IHP	P3-O13	4.91	1.68	1.59
38	А	3000	IHP	P5-O15	4.83	1.68	1.59
38	А	3000	IHP	P2-O12	4.64	1.68	1.59
38	А	3000	IHP	P6-O16	4.48	1.67	1.59
38	J	3000	IHP	P6-O16	4.37	1.67	1.59
38	А	3000	IHP	P3-O13	4.31	1.67	1.59
38	J	3000	IHP	P2-O12	4.30	1.67	1.59
38	А	3000	IHP	C3-C2	-4.24	1.43	1.52
38	J	3000	IHP	P5-O15	4.14	1.67	1.59
38	А	3000	IHP	P4-014	4.13	1.67	1.59
38	А	3000	IHP	P1-011	4.06	1.67	1.59
38	J	3000	IHP	P1-011	4.01	1.66	1.59
38	J	3000	IHP	P4-014	3.98	1.66	1.59
38	А	3000	IHP	C5-C4	3.25	1.59	1.52
39	С	1101	GTP	C5-C4	-3.04	1.35	1.43
38	J	3000	IHP	O12-C2	-2.85	1.33	1.44
38	J	3000	IHP	O13-C3	-2.82	1.33	1.44
38	А	3000	IHP	O12-C2	-2.71	1.34	1.44



Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)		
38	А	3000	IHP	O13-C3	-2.58	1.34	1.44		
38	J	3000	IHP	C2-C1	-2.53	1.47	1.52		
39	С	1101	GTP	C6-N1	-2.34	1.34	1.37		
39	С	1101	GTP	PG-O3G	-2.08	1.46	1.54		
39	С	1101	GTP	O4'-C4'	-2.01	1.40	1.45		

All (14) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms		$Observed(^{o})$	$Ideal(^{o})$
39	С	1101	GTP	PB-O3B-PG	-5.24	114.86	132.83
38	А	3000	IHP	O15-C5-C4	5.13	120.78	108.69
39	С	1101	GTP	C5-C6-N1	4.51	121.91	113.95
38	А	3000	IHP	O15-C5-C6	4.13	118.42	108.69
39	С	1101	GTP	C2-N1-C6	-3.55	118.56	125.10
39	С	1101	GTP	O6-C6-C5	-3.19	118.15	124.37
39	С	1101	GTP	PA-O3A-PB	-2.86	123.00	132.83
38	А	3000	IHP	C6-C5-C4	-2.76	104.37	110.41
39	С	1101	GTP	C8-N7-C5	2.35	107.48	102.99
38	А	3000	IHP	O11-C1-C2	2.34	114.21	108.69
38	J	3000	IHP	C5-C4-C3	2.32	115.48	110.41
38	А	3000	IHP	O13-C3-C2	2.25	114.00	108.69
38	A	3000	IHP	C5-C6-C1	-2.13	105.74	110.41
38	А	3000	IHP	C3-C2-C1	2.11	115.04	110.41

There are no chirality outliers.

All (4) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
38	А	3000	IHP	C4-C3-O13-P3
38	А	3000	IHP	C2-C3-O13-P3
38	А	3000	IHP	C3-O13-P3-O23
38	А	3000	IHP	C2-O12-P2-O32

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less then 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be



highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.











5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



6 Map visualisation (i)

This section contains visualisations of the EMDB entry EMD-19397. These allow visual inspection of the internal detail of the map and identification of artifacts.

No raw map or half-maps were deposited for this entry and therefore no images, graphs, etc. pertaining to the raw map can be shown.

6.1 Orthogonal projections (i)

6.1.1 Primary map



The images above show the map projected in three orthogonal directions.

6.2 Central slices (i)

6.2.1 Primary map



X Index: 224

Y Index: 224



Z Index: 224

The images above show central slices of the map in three orthogonal directions.

6.3 Largest variance slices (i)

6.3.1 Primary map



X Index: 221

Y Index: 228

Z Index: 282

The images above show the largest variance slices of the map in three orthogonal directions.

6.4 Orthogonal standard-deviation projections (False-color) (i)

6.4.1 Primary map



The images above show the map standard deviation projections with false color in three orthogonal directions. Minimum values are shown in green, max in blue, and dark to light orange shades represent small to large values respectively.



6.5 Orthogonal surface views (i)

6.5.1 Primary map



The images above show the 3D surface view of the map at the recommended contour level 0.5. These images, in conjunction with the slice images, may facilitate assessment of whether an appropriate contour level has been provided.

6.6 Mask visualisation (i)

This section was not generated. No masks/segmentation were deposited.



7 Map analysis (i)

This section contains the results of statistical analysis of the map.

7.1 Map-value distribution (i)



The map-value distribution is plotted in 128 intervals along the x-axis. The y-axis is logarithmic. A spike in this graph at zero usually indicates that the volume has been masked.



7.2 Volume estimate (i)



The volume at the recommended contour level is 1151 nm^3 ; this corresponds to an approximate mass of 1040 kDa.

The volume estimate graph shows how the enclosed volume varies with the contour level. The recommended contour level is shown as a vertical line and the intersection between the line and the curve gives the volume of the enclosed surface at the given level.



7.3 Rotationally averaged power spectrum (i)



*Reported resolution corresponds to spatial frequency of 0.345 ${\rm \AA^{-1}}$



8 Fourier-Shell correlation (i)

This section was not generated. No FSC curve or half-maps provided.



9 Map-model fit (i)

This section contains information regarding the fit between EMDB map EMD-19397 and PDB model 8RO0. Per-residue inclusion information can be found in section 3 on page 13.

9.1 Map-model overlay (i)



The images above show the 3D surface view of the map at the recommended contour level 0.5 at 50% transparency in yellow overlaid with a ribbon representation of the model coloured in blue. These images allow for the visual assessment of the quality of fit between the atomic model and the map.



9.2 Q-score mapped to coordinate model (i)



The images above show the model with each residue coloured according its Q-score. This shows their resolvability in the map with higher Q-score values reflecting better resolvability. Please note: Q-score is calculating the resolvability of atoms, and thus high values are only expected at resolutions at which atoms can be resolved. Low Q-score values may therefore be expected for many entries.

9.3 Atom inclusion mapped to coordinate model (i)



The images above show the model with each residue coloured according to its atom inclusion. This shows to what extent they are inside the map at the recommended contour level (0.5).



9.4 Atom inclusion (i)



At the recommended contour level, 85% of all backbone atoms, 76% of all non-hydrogen atoms, are inside the map.



1.0

0.0 <0.0

9.5 Map-model fit summary (i)

The table lists the average atom inclusion at the recommended contour level (0.5) and Q-score for the entire model and for each chain.

Chain	Atom inclusion	$\mathbf{Q} extsf{-score}$
All	0.7580	0.3720
2	0.7730	0.1980
5	0.9440	0.4570
6	0.6260	0.3220
А	0.8000	0.4350
В	0.8460	0.2000
С	0.9420	0.5700
D	0.4670	0.3640
E	0.8940	0.5330
Ι	0.8460	0.4750
In	0.6220	0.2310
J	0.7620	0.4580
K	0.6820	0.4230
L	0.5830	0.3790
М	0.7150	0.3100
N	0.9410	0.5520
0	0.8020	0.4680
Р	0.3610	0.2980
PX	0.4710	0.0840
Q	0.8220	0.5150
R	0.7750	0.4450
S	0.7380	0.4350
T	0.9210	0.5490
TF	0.8490	0.3840
W	0.6190	0.1620
Z	0.4980	0.4050
a	0.9130	0.5030
b	0.8860	0.4550
С	0.9100	0.4790
d	0.7980	0.3850
e	0.8640	0.4680
f	0.8600	0.4670
g	0.8250	0.4430
h	0.7500	0.0960
i	0.4880	0.0510



Chain	Atom inclusion	Q-score
j	0.5000	0.0490
k	0.4860	0.0630
1	0.6030	0.0520
m	0.6560	0.0740
n	0.6890	0.0690
0	0.5080	0.0310
р	0.5550	0.0610
q	0.7370	0.4590
r	0.4870	0.3880
s	0.3600	0.1630
t	0.5380	0.3930
У	0.4010	0.1490

