

Full wwPDB X-ray Structure Validation Report (i)

Jul 2, 2025 – 05:25 pm BST

PDB ID : 8S0V / pdb 00008s0v

Title : Crystal structure of Cryptosporidium parvum - Trypanosoma cruzi mutant

lysyl tRNA synthetase in complex with inhibitor

Authors : Dawson, A.; Wyllie, S.

Deposited on : 2024-02-14

Resolution : 1.60 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org
A user guide is available at

https://www.wwpdb.org/validation/2017/XrayValidationReportHelp with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (1)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0rc1

Mogul : 1.8.4, CSD as 541 be (2020)

Xtriage (Phenix) : 2.0rc1

EDS : 3.0 buster-report : 1.1.7 (2018)

Percentile statistics : 20231227.v01 (using entries in the PDB archive December 27th 2023)

CCP4 : 9.0.003 (Gargrove)

Density-Fitness : 1.0.11

Ideal geometry (proteins) : Engh & Huber (2001) Ideal geometry (DNA, RNA) : Parkinson et al. (1996)


Validation Pipeline (wwPDB-VP) : 2.44

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: X- $RAY\ DIFFRACTION$

The reported resolution of this entry is 1.60 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive	Similar resolution
Metric	$(\# ext{Entries})$	$(\# ext{Entries}, ext{ resolution range}(ext{Å}))$
R_{free}	164625	4274 (1.60-1.60)
Clashscore	180529	4682 (1.60-1.60)
Ramachandran outliers	177936	4583 (1.60-1.60)
Sidechain outliers	177891	4582 (1.60-1.60)
RSRZ outliers	164620	4272 (1.60-1.60)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain		
1	A	536	85%	7%	7%
1	В	536	84%	8%	8%



2 Entry composition (i)

There are 7 unique types of molecules in this entry. The entry contains 8912 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

• Molecule 1 is a protein called Lysine–tRNA ligase.

	\mathbf{Mol}	Chain	Residues	Atoms			ZeroOcc	AltConf	Trace		
	1	Λ	496	Total	С	Ν	О	S	0	9	0
	1	Λ	490	4029	2584	670	747	28	U	2	0
ſ	1	B	495	Total	С	N	О	S	0	2	0
	1	D	490	4020	2579	669	744	28	0		

There are 52 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	24	MET	-	initiating methionine	UNP Q5CR27
A	25	GLY	-	expression tag	UNP Q5CR27
A	26	SER	-	expression tag	UNP Q5CR27
A	27	SER	-	expression tag	UNP Q5CR27
A	28	HIS	-	expression tag	UNP Q5CR27
A	29	HIS	-	expression tag	UNP Q5CR27
A	30	HIS	-	expression tag	UNP Q5CR27
A	31	HIS	-	expression tag	UNP Q5CR27
A	32	HIS	-	expression tag	UNP Q5CR27
A	33	HIS	-	expression tag	UNP Q5CR27
A	34	SER	-	expression tag	UNP Q5CR27
A	35	SER	-	expression tag	UNP Q5CR27
A	36	GLY	-	expression tag	UNP Q5CR27
A	37	GLU	-	expression tag	UNP Q5CR27
A	38	ASN	-	expression tag	UNP Q5CR27
A	39	LEU	-	expression tag	UNP Q5CR27
A	40	TYR	-	expression tag	UNP Q5CR27
A	41	PHE	-	expression tag	UNP Q5CR27
A	42	GLN	-	expression tag	UNP Q5CR27
A	43	GLY	-	expression tag	UNP Q5CR27
A	44	HIS	-	expression tag	UNP Q5CR27
A	45	MET	-	expression tag	UNP Q5CR27
A	290	LEU	ILE	engineered mutation	UNP Q5CR27
A	309	SER	ALA	engineered mutation	UNP Q5CR27
A	310	VAL	MET	engineered mutation	UNP Q5CR27




Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
A	538	LEU	ILE	engineered mutation	UNP Q5CR27
В	24	MET	_	initiating methionine	UNP Q5CR27
В	25	GLY	-	expression tag	UNP Q5CR27
В	26	SER	-	expression tag	UNP Q5CR27
В	27	SER	-	expression tag	UNP Q5CR27
В	28	HIS	-	expression tag	UNP Q5CR27
В	29	HIS	-	expression tag	UNP Q5CR27
В	30	HIS	-	expression tag	UNP Q5CR27
В	31	HIS	-	expression tag	UNP Q5CR27
В	32	HIS	-	expression tag	UNP Q5CR27
В	33	HIS	-	expression tag	UNP Q5CR27
В	34	SER	-	expression tag	UNP Q5CR27
В	35	SER	-	expression tag	UNP Q5CR27
В	36	GLY	-	expression tag	UNP Q5CR27
В	37	GLU	-	expression tag	UNP Q5CR27
В	38	ASN	-	expression tag	UNP Q5CR27
В	39	LEU	-	expression tag	UNP Q5CR27
В	40	TYR	_	expression tag	UNP Q5CR27
В	41	PHE	-	expression tag	UNP Q5CR27
В	42	GLN	-	expression tag	UNP Q5CR27
В	43	GLY	-	expression tag	UNP Q5CR27
В	44	HIS	-	expression tag	UNP Q5CR27
В	45	MET	-	expression tag	UNP Q5CR27
В	290	LEU	ILE	engineered mutation	UNP Q5CR27
В	309	SER	ALA	engineered mutation	UNP Q5CR27
В	310	VAL	MET	engineered mutation	UNP Q5CR27
В	538	LEU	ILE	engineered mutation	UNP Q5CR27


 \bullet Molecule 2 is LYSINE (CCD ID: LYS) (formula: $\mathrm{C_6H_{15}N_2O_2}).$





Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	A	1	Total 10				0	0
2	В	1	Total 10		N 2	O 2	0	0

 $\bullet \ \, \text{Molecule 3 is} \ \, \{N\} \text{7-cyclopentyl-5,6,8-tris} \\ \text{(fluoranyl)-2-methyl-quinazoline-4,7-diamine} \\ \text{(CCD ID: A1H4X)} \\ \text{(formula: $C_{14}H_{15}F_3N_4$)} \\ \text{(labeled as "Ligand of Interest" by depositor)}.$




\mathbf{Mol}	Chain	Residues	Atoms				ZeroOcc	AltConf
3	A	1	Total 21	C 14	_	N 4	0	0



Continued from previous page...

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	В	1	Total C 21 14	F 3	N 4	0	0


• Molecule 4 is GLYCEROL (CCD ID: GOL) (formula: $C_3H_8O_3$).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	1	Total C O 6 3 3	0	0
4	A	1	Total C O 6 3 3	0	0
4	В	1	Total C O 6 3 3	0	0


• Molecule 5 is 2-AMINO-2-HYDROXYMETHYL-PROPANE-1,3-DIOL (CCD ID: TRS) (formula: $C_4H_{12}NO_3$).





Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	A	1	Total C N O 8 4 1 3	0	0
5	В	1	Total C N O 8 4 1 3	0	0
5	В	1	Total C N O 8 4 1 3	0	0

 \bullet Molecule 6 is SULFATE ION (CCD ID: SO4) (formula: $\mathrm{O_4S}).$



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
6	A	1	Total 5	O 4	S 1	0	0



• Molecule 7 is water.


Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
7	A	377	Total O 377 377	0	0
7	В	377	Total O 377 377	0	0



3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Lysine–tRNA ligase





4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 2 21 21	Depositor
Cell constants	73.14Å 116.70Å 142.88Å	Depositor
a, b, c, α , β , γ	90.00° 90.00° 90.00°	Depositor
Resolution (Å)	58.35 - 1.60	Depositor
resolution (A)	58.35 - 1.60	EDS
% Data completeness	99.7 (58.35-1.60)	Depositor
(in resolution range)	99.7 (58.35-1.60)	EDS
R_{merge}	0.08	Depositor
R_{sym}	(Not available)	Depositor
$< I/\sigma(I) > 1$	1.79 (at 1.60Å)	Xtriage
Refinement program	REFMAC 5.8.0425	Depositor
P.P.	0.184 , 0.208	Depositor
R, R_{free}	0.193 , 0.217	DCC
R_{free} test set	8166 reflections (5.06%)	wwPDB-VP
Wilson B-factor (Å ²)	20.3	Xtriage
Anisotropy	0.241	Xtriage
Bulk solvent $k_{sol}(e/Å^3)$, $B_{sol}(Å^2)$	0.35, 34.6	EDS
L-test for twinning ²	$ < L > = 0.49, < L^2> = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	8912	wwPDB-VP
Average B, all atoms $(Å^2)$	25.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: The largest off-origin peak in the Patterson function is 4.13% of the height of the origin peak. No significant pseudotranslation is detected.

²Theoretical values of <|L|>, $<L^2>$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



¹Intensities estimated from amplitudes.

5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: GOL, A1H4X, TRS, SO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol Chain		Bond lengths		Bond angles	
IVIOI	Chain	RMSZ	# Z > 5	RMSZ	# Z > 5
1	A	0.79	0/4133	1.12	6/5574 (0.1%)
1	В	0.80	0/4124	1.13	7/5562 (0.1%)
All	All	0.79	0/8257	1.12	13/11136 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a maintenain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1
1	В	0	3
All	All	0	4

There are no bond length outliers.

All (13) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	\mathbf{Z}	$Observed(^o)$	$Ideal(^{o})$
1	A	406	HIS	CA-CB-CG	-8.33	105.47	113.80
1	A	219	ARG	CG-CD-NE	-7.33	95.87	112.00
1	A	244	ASN	CA-CB-CG	6.99	119.59	112.60
1	A	216	PHE	CA-CB-CG	-6.27	107.53	113.80
1	В	216	PHE	CA-CB-CG	-6.10	107.70	113.80
1	A	196	GLU	N-CA-CB	5.59	119.67	110.39
1	В	304	ASN	O-C-N	-5.43	116.74	121.35
1	В	264	THR	CA-CB-OG1	-5.34	101.59	109.60
1	В	286	LYS	CG-CD-CE	5.24	123.34	111.30
1	A	338	HIS	CB-CG-CD2	-5.21	124.43	131.20
1	В	244	ASN	CA-CB-CG	5.15	117.75	112.60
1	В	408	ARG	N-CA-CB	-5.12	104.29	110.53



Continued from previous page...

Mol	Chain	Res	Type	Atoms	\mathbf{Z}	$\mathbf{Observed}(^{o})$	$\operatorname{Ideal}(^{o})$
1	В	477	GLN	CB-CA-C	-5.11	102.31	110.79

There are no chirality outliers.

All (4) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	198	ARG	Sidechain
1	В	150	ARG	Sidechain
1	В	198	ARG	Sidechain
1	В	219	ARG	Sidechain

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4029	0	3988	29	0
1	В	4020	0	3982	28	0
2	A	10	0	12	0	0
2	В	10	0	12	0	0
3	A	21	0	0	0	0
3	В	21	0	0	0	0
4	A	12	0	16	0	0
4	В	6	0	8	0	0
5	A	8	0	12	0	0
5	В	16	0	24	0	0
6	A	5	0	0	1	0
7	A	377	0	0	4	0
7	В	377	0	0	8	0
All	All	8912	0	8054	51	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (51) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.



		Interatomic	Clash
Atom-1	Atom-2	distance (Å)	overlap (Å)
1:A:268[A]:MET:HE3	1:B:268[A]:MET:HE3	1.49	0.94
1:B:219:ARG:HG3	1:B:525:ILE:HD13	1.74	0.70
1:B:51:ARG:HH21	1:B:54:MET:HE1	1.58	0.69
1:A:246:ILE:HD11	1:A:498:PRO:HG2	1.80	0.62
1:A:297:GLU:O	1:B:259:HIS:HE1	1.82	0.62
1:A:338:HIS:HD2	7:A:1036:HOH:O	1.82	0.60
1:A:246:ILE:HD11	1:A:498:PRO:CG	2.33	0.59
1:A:51:ARG:HH21	1:A:54:MET:HE1	1.69	0.56
1:A:246:ILE:CD1	1:A:498:PRO:HG2	2.35	0.56
1:B:338:HIS:HD2	7:B:1048:HOH:O	1.88	0.55
1:A:246:ILE:HD11	1:A:498:PRO:HB2	1.87	0.55
1:A:183:HIS:HD2	7:A:1043:HOH:O	1.90	0.54
1:A:420:HIS:HE1	7:A:1029:HOH:O	1.89	0.54
1:B:219:ARG:HG3	1:B:525:ILE:CD1	2.36	0.54
1:A:131:ILE:HG23	1:A:132:MET:HE2	1.90	0.53
1:A:140:GLU:OE1	1:A:142:SER:OG	2.25	0.53
1:A:58:ILE:HG23	1:A:63:ARG:HB2	1.91	0.53
1:A:198:ARG:NH2	7:A:706:HOH:O	2.41	0.53
1:B:117:ASP:OD2	1:B:124:LYS:HE3	2.09	0.51
1:A:296:ASN:HD21	1:B:260:ASN:HD22	1.59	0.50
1:B:299:ILE:HG23	7:B:858:HOH:O	2.11	0.50
1:B:54:MET:HE2	1:B:55:MET:SD	2.51	0.50
1:A:246:ILE:HD11	1:A:498:PRO:CB	2.44	0.48
1:A:246:ILE:O	1:A:246:ILE:HG23	2.13	0.47
1:A:389:ASN:O	1:A:393:MET:HG3	2.15	0.46
1:B:176:VAL:HG23	7:B:811:HOH:O	2.15	0.46
1:B:400:HIS:HE1	7:B:728:HOH:O	1.98	0.45
1:B:117:ASP:CG	1:B:124:LYS:HE3	2.41	0.45
1:A:296:ASN:HD21	1:B:260:ASN:ND2	2.15	0.45
1:A:207:MET:HE3	1:A:207:MET:HB2	1.78	0.45
1:A:355:ILE:HD11	1:A:461:LEU:HD11	2.00	0.45
1:B:265:GLN:OE1	7:B:701:HOH:O	2.21	0.45
1:A:243:LEU:HD21	1:B:268[A]:MET:HE1	2.00	0.44
1:A:246:ILE:CD1	1:A:498:PRO:CG	2.95	0.43
1:B:420:HIS:HE1	7:B:1040:HOH:O	2.01	0.43
1:B:98:SER:OG	1:B:157:THR:HG22	2.17	0.43
1:B:382:ARG:HA	1:B:383:PRO:C	2.44	0.43
1:A:126:GLN:O	1:A:169:SER:HA	2.19	0.42
1:A:184:MET:HE2	1:A:184:MET:HB3	2.00	0.42
1:B:218:LEU:HD23	1:B:218:LEU:C	2.45	0.42
1:A:98:SER:OG	1:A:157:THR:HG22	2.20	0.42
1:A:481:PHE:O	1:A:485[A]:GLN:HG2	2.19	0.42



I 'ontinued	trom	mromonie	naaa
Continued	110110	DICULUUS	pauc

Atom-1	Atom-2	Interatomic	Clash
Atom-1	Atom-2	$\operatorname{distance}\ (ext{Å})$	overlap (Å)
1:B:299:ILE:CG2	7:B:858:HOH:O	2.67	0.42
1:B:481:PHE:O	1:B:485[A]:GLN:HG2	2.20	0.42
6:A:606:SO4:O3	1:B:260:ASN:HB3	2.20	0.41
1:B:512:PRO:O	1:B:513:PRO:C	2.61	0.41
1:B:259:HIS:HD2	7:B:1037:HOH:O	2.02	0.41
1:A:474:GLU:HG2	1:B:66:TYR:CE2	2.55	0.41
1:A:382:ARG:HA	1:A:383:PRO:C	2.46	0.41
1:B:126:GLN:O	1:B:169:SER:HA	2.21	0.41
1:B:366:PHE:HB3	1:B:371:GLU:HG3	2.03	0.40

There are no symmetry-related clashes.

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Perce	ntiles
1	A	494/536~(92%)	484 (98%)	10 (2%)	0	100	100
1	В	493/536~(92%)	483 (98%)	10 (2%)	0	100	100
All	All	987/1072~(92%)	967 (98%)	20 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.



Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	443/475 (93%)	442 (100%)	1 (0%)	92 86
1	В	442/475 (93%)	436 (99%)	6 (1%)	62 43
All	All	885/950 (93%)	878 (99%)	7 (1%)	79 66

All (7) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	319	TYR
1	В	71	LYS
1	В	85	ASN
1	В	187	THR
1	В	319	TYR
1	В	461	LEU
1	В	522	ASP

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (15) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	123	GLN
1	A	183	HIS
1	A	228	ASN
1	A	244	ASN
1	A	338	HIS
1	A	400	HIS
1	A	420	HIS
1	В	50	ASN
1	В	244	ASN
1	В	259	HIS
1	В	260	ASN
1	В	338	HIS
1	В	400	HIS
1	В	420	HIS
1	В	545	ASN

5.3.3 RNA (i)

There are no RNA molecules in this entry.



5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

There are no oligosaccharides in this entry.

5.6 Ligand geometry (i)

11 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mal	Trens	Chain	Dag	Link	Во	ond leng	ths	В	ond ang	eles
Mol	Type	Chain	Res	Lilik	Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
4	GOL	В	603	-	5,5,5	0.32	0	5,5,5	1.62	1 (20%)
6	SO4	A	606	-	4,4,4	0.45	0	6,6,6	0.60	0
2	LYS	A	601	-	8,9,9	0.77	0	9,10,10	1.04	0
5	TRS	В	605	-	7,7,7	0.32	0	9,9,9	0.38	0
5	TRS	В	604	_	7,7,7	0.21	0	9,9,9	0.25	0
3	A1H4X	A	602	-	23,23,23	0.55	0	33,34,34	0.98	1 (3%)
3	A1H4X	В	602	-	23,23,23	0.52	0	33,34,34	1.04	3 (9%)
4	GOL	A	604	-	5,5,5	0.28	0	5,5,5	0.76	0
5	TRS	A	605	-	7,7,7	0.23	0	9,9,9	0.44	0
2	LYS	В	601	-	8,9,9	0.41	0	9,10,10	1.02	0
4	GOL	A	603	-	5,5,5	0.09	0	5,5,5	0.18	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	GOL	В	603	-	-	4/4/4/4	-
2	LYS	A	601	-	-	0/9/9/9	-
5	TRS	В	605	-	-	6/9/9/9	-



 $Continued\ from\ previous\ page...$

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	TRS	В	604	-	-	0/9/9/9	-
3	A1H4X	A	602	-	-	0/4/11/11	0/3/3/3
3	A1H4X	В	602	_	-	0/4/11/11	0/3/3/3
4	GOL	A	604	-	_	4/4/4/4	_
5	TRS	A	605	-	-	6/9/9/9	-
2	LYS	В	601	-	-	0/9/9/9	-
4	GOL	A	603	-	-	1/4/4/4	-

There are no bond length outliers.

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	\mathbf{Z}	$\mathbf{Observed}(^{o})$	$\operatorname{Ideal}({}^{o})$
3	В	602	A1H4X	C12-C13-C9	2.92	116.84	114.86
3	A	602	A1H4X	C10-N2-C9	2.65	117.57	115.45
3	В	602	A1H4X	C10-N2-C9	2.27	117.27	115.45
4	В	603	GOL	O2-C2-C3	2.21	118.85	109.12
3	В	602	A1H4X	C4-C3-N1	2.12	116.27	111.76

There are no chirality outliers.

All (21) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	A	604	GOL	O1-C1-C2-C3
4	A	604	GOL	C1-C2-C3-O3
4	В	603	GOL	O1-C1-C2-C3
5	A	605	TRS	C1-C-C2-O2
5	A	605	TRS	C3-C-C2-O2
5	A	605	TRS	N-C-C2-O2
5	В	605	TRS	C2-C-C1-O1
5	В	605	TRS	C3-C-C1-O1
5	В	605	TRS	N-C-C1-O1
5	В	605	TRS	C1-C-C3-O3
5	В	605	TRS	C2-C-C3-O3
4	A	604	GOL	O1-C1-C2-O2
4	A	603	GOL	O1-C1-C2-C3
4	В	603	GOL	C1-C2-C3-O3
4	A	604	GOL	O2-C2-C3-O3
4	В	603	GOL	O1-C1-C2-O2
4	В	603	GOL	O2-C2-C3-O3
5	A	605	TRS	N-C-C1-O1
5	В	605	TRS	N-C-C3-O3



Continued from previous page...

Mol	Chain	Res	Type	Atoms
5	A	605	TRS	C2-C-C1-O1
5	A	605	TRS	C3-C-C1-O1


There are no ring outliers.


1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
6	A	606	SO4	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less then 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.









5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ>2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95^{th} percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ $>$	$\# \mathrm{RSRZ}{>}2$	$OWAB(A^2)$	Q<0.9
1	A	496/536 (92%)	0.20	19 (3%) 44 44	12, 21, 41, 76	2 (0%)
1	В	495/536~(92%)	0.24	16 (3%) 50 51	11, 22, 46, 82	2 (0%)
All	All	991/1072 (92%)	0.22	35 (3%) 47 48	11, 22, 44, 82	4 (0%)

All (35) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	189	ILE	5.8
1	В	189	ILE	5.6
1	A	187	THR	4.6
1	В	187	THR	4.5
1	A	46	HIS	4.4
1	В	45	MET	3.9
1	A	188	ALA	3.7
1	A	45	MET	3.4
1	A	246	ILE	3.4
1	В	188	ALA	3.3
1	A	544	ARG	3.3
1	A	57	CYS	3.3
1	A	195	GLN	3.2
1	A	165	ARG	3.0
1	В	46	HIS	3.0
1	A	87	GLU	2.5
1	В	81	LEU	2.5
1	A	54	MET	2.5
1	A	136	ILE	2.5
1	В	91	ILE	2.4
1	A	194	ASP	2.4
1	В	545	ASN	2.4
1	A	132	MET	2.3
1	В	186	PRO	2.2



Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	В	84	GLY	2.2
1	A	61	ALA	2.2
1	В	232	ARG	2.2
1	A	50	ASN	2.1
1	В	184	MET	2.1
1	В	95	THR	2.0
1	В	107	ARG	2.0
1	A	499	ILE	2.0
1	В	86	VAL	2.0
1	A	56	GLU	2.0
1	В	299	ILE	2.0

6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates (i)

There are no oligosaccharides in this entry.


6.4 Ligands (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95^{th} percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	$\operatorname{B-factors}(\mathring{\mathrm{A}}^2)$	Q<0.9
5	TRS	A	605	8/8	0.74	0.17	38,43,49,53	0
5	TRS	В	605	8/8	0.78	0.15	40,40,43,49	0
4	GOL	A	603	6/6	0.87	0.13	35,39,44,44	0
4	GOL	A	604	6/6	0.89	0.12	28,30,37,38	0
5	TRS	В	604	8/8	0.93	0.09	24,29,32,34	0
4	GOL	В	603	6/6	0.93	0.09	21,23,23,27	0
3	A1H4X	A	602	21/21	0.94	0.07	17,21,25,27	0
2	LYS	A	601	10/10	0.95	0.07	15,17,20,21	0
3	A1H4X	В	602	21/21	0.95	0.07	17,20,25,27	0
2	LYS	В	601	10/10	0.96	0.06	16,18,20,22	0
6	SO4	A	606	5/5	0.96	0.09	21,22,28,29	5



The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.







6.5 Other polymers (i)

There are no such residues in this entry.

