

Oct 7, 2024 - 02:26 PM JST

PDB ID	:	8W8S
EMDB ID	:	EMD-37358
Title	:	Cryo-EM structure of the AA14-bound GPR101 complex
Authors	:	Sun, J.P.; Yu, X.; Gao, N.; Yang, F.; Wang, J.Y.; Yang, Z.; Guan, Y.; Wang,
		G.P.
Deposited on	:	2023-09-04
Resolution	:	3.30  Å(reported)

This is a Full wwPDB EM Validation Report for a publicly released PDB entry.

We welcome your comments at *validation@mail.wwpdb.org* A user guide is available at https://www.wwpdb.org/validation/2017/EMValidationReportHelp with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (1)) were used in the production of this report:

EMDB validation analysis	:	FAILED
Mogul	:	1.8.5 (274361), CSD as541be (2020)
MolProbity	:	4.02b-467
buster-report	:	1.1.7(2018)
Percentile statistics	:	20231227.v01 (using entries in the PDB archive December 27th 2023)
$\operatorname{MapQ}$	:	FAILED
Ideal geometry (proteins)	:	Engh & Huber $(2001)$
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.39

# 1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure:  $ELECTRON\ MICROSCOPY$ 

The reported resolution of this entry is 3.30 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	(# Entries)	(# Entries)
Clashscore	210492	15764
Ramachandran outliers	207382	16835
Sidechain outliers	206894	16415

The table below summarises the geometric issues observed across the polymeric chains and their fit to the map. The red, orange, yellow and green segments of the bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5%

Mol	Chain	Length	Quality of chain		
1	R	508	47% 7	% •	45%



# 2 Entry composition (i)

There are 2 unique types of molecules in this entry. The entry contains 2265 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

• Molecule 1 is a protein called Probable G-protein coupled receptor 101.

Mol	Chain	Residues	Atoms			AltConf	Trace		
1	R	281	Total 2223	C 1496	N 357	O 357	S 13	0	0

• Molecule 2 is 1-(4-methylpyridin-2-yl)-3-[3-(trifluoromethyl)phenyl]thiourea (three-letter code: U7D) (formula:  $C_{14}H_{12}F_3N_3S$ ).



Mol	Chain	Residues	Atoms				AltConf	
0	D	1	Total	С	F	Ν	$\mathbf{S}$	0
Z R	1	21	14	3	3	1	0	
0	D	1	Total	С	F	Ν	S	0
	n	1	21	14	3	3	1	0



# 3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.



• Molecule 1: Probable G-protein coupled receptor 101



# 4 Experimental information (i)

Property	Value	Source
EM reconstruction method	SINGLE PARTICLE	Depositor
Imposed symmetry	POINT, Not provided	
Number of particles used	3646024	Depositor
Resolution determination method	FSC 0.143 CUT-OFF	Depositor
CTF correction method	PHASE FLIPPING AND AMPLITUDE	Depositor
	CORRECTION	
Microscope	FEI TITAN KRIOS	Depositor
Voltage (kV)	300	Depositor
Electron dose $(e^-/\text{\AA}^2)$	60	Depositor
Minimum defocus (nm)	800	Depositor
Maximum defocus (nm)	1200	Depositor
Magnification	Not provided	
Image detector	GATAN K3 $(6k \ge 4k)$	Depositor



# 5 Model quality (i)

## 5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: U7D

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mal	Chain	Bond	lengths	Bond angles	
		RMSZ	# Z  > 5	RMSZ	# Z  > 5
1	R	0.30	0/2287	0.45	0/3132

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	R	2223	0	2310	40	0
2	R	42	0	0	0	0
All	All	2265	0	2310	40	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 9.

All (40) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:R:99:LEU:CD2	1:R:103:PHE:CD2	2.55	0.90
1:R:173:ALA:HA	1:R:184:MET:HA	1.57	0.84

Continued on next page...



EMD-37358,	8W8S
------------	------

		Interatomic	Clash
Atom-1	Atom-2	distance (Å)	overlap (Å)
1:R:99:LEU:CD2	1:R:103:PHE:HD2	1.91	0.83
1:R:99:LEU:HD23	1:R:103:PHE:CD2	2.14	0.83
1:R:432:PRO:HB2	1:R:434:TRP:CD1	2.19	0.77
1:R:62:LEU:HG	1:R:63:LEU:HD13	1.67	0.76
1:R:176:ASP:O	1:R:180:ALA:HA	1.89	0.72
1:R:99:LEU:HD21	1:R:103:PHE:CD2	2.28	0.68
1:R:175:PHE:CZ	1:R:180:ALA:HB1	2.29	0.68
1:R:99:LEU:HD23	1:R:103:PHE:HD2	1.54	0.66
1:R:80:GLN:O	1:R:84:VAL:HB	1.99	0.62
1:R:432:PRO:HB2	1:R:434:TRP:NE1	2.14	0.61
1:R:65:VAL:O	1:R:65:VAL:HG23	2.01	0.60
1:R:99:LEU:HD21	1:R:103:PHE:CE2	2.39	0.57
1:R:176:ASP:O	1:R:180:ALA:CA	2.52	0.56
1:R:174:ALA:O	1:R:182:CYS:HA	2.06	0.56
1:R:175:PHE:HA	1:R:182:CYS:HA	1.88	0.55
1:R:194:ILE:O	1:R:198:VAL:HG12	2.09	0.53
1:R:53:ALA:HB2	1:R:74:LEU:HD21	1.90	0.53
1:R:37:VAL:HG11	1:R:441:TRP:CZ2	2.44	0.52
1:R:63:LEU:N	1:R:63:LEU:CD1	2.73	0.52
1:R:176:ASP:OD1	1:R:181:LEU:N	2.45	0.50
1:R:165:PRO:HG2	1:R:192:TYR:CE1	2.47	0.50
1:R:432:PRO:CB	1:R:434:TRP:HE1	2.26	0.49
1:R:93:VAL:HG22	1:R:94:PRO:HD2	1.95	0.48
1:R:62:LEU:HB3	1:R:67:ASN:HD21	1.78	0.48
1:R:63:LEU:N	1:R:63:LEU:HD12	2.29	0.48
1:R:432:PRO:CB	1:R:434:TRP:NE1	2.78	0.47
1:R:62:LEU:O	1:R:67:ASN:ND2	2.48	0.46
1:R:184:MET:HE3	1:R:184:MET:HB2	1.81	0.45
1:R:463:LYS:HE2	1:R:463:LYS:HB2	1.68	0.45
1:R:96:PHE:CD1	1:R:96:PHE:O	2.70	0.44
1:R:116:PHE:CD2	1:R:163:SER:HB2	2.52	0.44
1:R:170:TRP:CD1	1:R:192:TYR:HB2	2.53	0.44
1:R:420:VAL:O	1:R:423:VAL:HG12	2.17	0.44
1:R:151:LEU:O	1:R:155:THR:HG22	2.19	0.43
1:R:62:LEU:HB3	1:R:67:ASN:ND2	2.35	0.42
1:R:176:ASP:O	1:R:180:ALA:N	2.53	0.42
1:R:471:LYS:HE3	1:R:471:LYS:HB3	1.90	0.42
1:R:38:LEU:O	1:R:38:LEU:HD23	2.21	0.41

Continued from previous page...

There are no symmetry-related clashes.



### 5.3 Torsion angles (i)

#### 5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percer	ntiles
1	R	277/508~(54%)	266 (96%)	11 (4%)	0	100	100

There are no Ramachandran outliers to report.

#### 5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent side chain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	R	244/451~(54%)	240~(98%)	4 (2%)	58 76	

All (4) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	R	99	LEU
1	R	182	CYS
1	R	200	PHE
1	R	434	TRP

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type	
1	R	67	ASN	



#### 5.3.3 RNA (i)

There are no RNA molecules in this entry.

### 5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates (i)

There are no oligosaccharides in this entry.

### 5.6 Ligand geometry (i)

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mal	Mal Tuna Chain Bag		Dec	Tink	Bo	ond leng	$_{\rm sths}$	Bond angles		
INIOI	туре	Unain	nes		Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
2	U7D	R	601	-	22,22,22	2.09	4 (18%)	31,31,31	1.83	2 (6%)
2	U7D	R	602	-	22,22,22	2.12	4 (18%)	31,31,31	1.92	3 (9%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	U7D	R	601	-	-	4/14/14/14	0/2/2/2
2	U7D	R	602	-	-	2/14/14/14	0/2/2/2

All (8) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	$\mathbf{Z}$	Observed(Å)	Ideal(Å)
2	R	601	U7D	C07-N02	5.99	1.45	1.35

Continued on next page...



Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	R	602	U7D	C07-N03	5.98	1.45	1.35
2	R	602	U7D	C07-N02	5.92	1.45	1.35
2	R	601	U7D	C07-N03	5.79	1.45	1.35
2	R	602	U7D	C07-S01	-3.28	1.60	1.68
2	R	601	U7D	C07-S01	-3.20	1.60	1.68
2	R	601	U7D	C03-N02	2.32	1.45	1.40
2	R	602	U7D	C03-N02	2.30	1.45	1.40

Continued from previous page...

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Ζ	$Observed(^{o})$	$Ideal(^{o})$
2	R	602	U7D	C03-N02-C07	-8.76	120.96	130.75
2	R	601	U7D	C03-N02-C07	-8.10	121.70	130.75
2	R	601	U7D	C08-N03-C07	-3.39	119.10	128.57
2	R	602	U7D	C08-N03-C07	-2.81	120.72	128.57
2	R	602	U7D	N03-C07-N02	2.18	120.23	114.81

There are no chirality outliers.

All (6) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	R	601	U7D	N01-C03-N02-C07
2	R	602	U7D	N01-C03-N02-C07
2	R	602	U7D	C04-C03-N02-C07
2	R	601	U7D	C04-C03-N02-C07
2	R	601	U7D	C09-C08-N03-C07
2	R	601	U7D	C13-C08-N03-C07

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less then 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient





equivalents in the CSD to analyse the geometry.





## 5.7 Other polymers (i)

There are no such residues in this entry.

## 5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.

