



Full wwPDB X-ray Structure Validation Report ⓘ

Nov 21, 2024 – 04:07 PM EST

PDB ID : 9CGS
Title : Structure of human MAIT A-F7 TCR in complex with human MR1-Pyridoxal-5'-phosphate
Authors : Awad, W.; Rossjohn, R.
Deposited on : 2024-06-30
Resolution : 2.00 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 1.21
EDS : 3.0
Percentile statistics : 20231227.v01 (using entries in the PDB archive December 27th 2023)
CCP4 : 9.0.004 (Gargrove)
Density-Fitness : 1.0.11
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.39


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION






The reported resolution of this entry is 2.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.






Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	164625	9409 (2.00-2.00)
Clashscore	180529	10737 (2.00-2.00)
Ramachandran outliers	177936	10628 (2.00-2.00)
Sidechain outliers	177891	10627 (2.00-2.00)
RSRZ outliers	164620	9409 (2.00-2.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	271	 85% 12% ..
1	C	271	 86% 10% ..
2	B	100	 89% 9% .
2	F	100	 94% 6%
3	D	204	 89% 5% 5%

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
3	G	204	 86% 11% •
4	E	246	 2% 87% 11% •
4	H	246	 % 91% 8% •

2 Entry composition [i](#)

There are 8 unique types of molecules in this entry. The entry contains 13983 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Major histocompatibility complex class I-related gene protein.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace	
			Total	C	N	O	P				S
1	A	266	Total	C	N	O	P	S	6	9	0
			2234	1432	383	406	1	12			
1	C	267	Total	C	N	O	P	S	2	5	0
			2236	1438	383	403	1	11			

There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	0	MET	-	initiating methionine	UNP Q95460
A	261	SER	CYS	conflict	UNP Q95460
C	0	MET	-	initiating methionine	UNP Q95460
C	261	SER	CYS	conflict	UNP Q95460

- Molecule 2 is a protein called Beta-2-microglobulin.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	B	98	Total	C	N	O	S	0	0	0
			784	503	134	144	3			
2	F	100	Total	C	N	O	S	0	0	0
			812	518	138	152	4			

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	0	MET	-	initiating methionine	UNP P61769
F	0	MET	-	initiating methionine	UNP P61769

- Molecule 3 is a protein called TRA@ protein.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
3	D	194	Total	C	N	O	S	0	4	0
			1498	957	237	294	10			

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
3	G	200	1600	1016	254	319	11	1	9	0

There are 42 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
D	0	MET	-	initiating methionine	UNP Q6P4G7
D	91	LYS	ARG	conflict	UNP Q6P4G7
D	93	SER	ALA	conflict	UNP Q6P4G7
D	94	ASN	SER	conflict	UNP Q6P4G7
D	95	TYR	ARG	conflict	UNP Q6P4G7
D	96	GLN	ARG	conflict	UNP Q6P4G7
D	?	-	GLY	deletion	UNP Q6P4G7
D	?	-	GLY	deletion	UNP Q6P4G7
D	?	-	GLY	deletion	UNP Q6P4G7
D	?	-	ASN	deletion	UNP Q6P4G7
D	?	-	LYS	deletion	UNP Q6P4G7
D	98	ILE	THR	conflict	UNP Q6P4G7
D	99	TRP	PHE	conflict	UNP Q6P4G7
D	101	ALA	THR	conflict	UNP Q6P4G7
D	104	LYS	GLN	conflict	UNP Q6P4G7
D	106	ILE	LYS	conflict	UNP Q6P4G7
D	107	ILE	VAL	conflict	UNP Q6P4G7
D	108	LYS	GLU	conflict	UNP Q6P4G7
D	109	PRO	LEU	conflict	UNP Q6P4G7
D	110	ASP	ASN	conflict	UNP Q6P4G7
D	157	CYS	THR	conflict	UNP Q6P4G7
G	0	MET	-	initiating methionine	UNP Q6P4G7
G	91	LYS	ARG	conflict	UNP Q6P4G7
G	93	SER	ALA	conflict	UNP Q6P4G7
G	94	ASN	SER	conflict	UNP Q6P4G7
G	95	TYR	ARG	conflict	UNP Q6P4G7
G	96	GLN	ARG	conflict	UNP Q6P4G7
G	?	-	GLY	deletion	UNP Q6P4G7
G	?	-	GLY	deletion	UNP Q6P4G7
G	?	-	GLY	deletion	UNP Q6P4G7
G	?	-	ASN	deletion	UNP Q6P4G7
G	?	-	LYS	deletion	UNP Q6P4G7
G	98	ILE	THR	conflict	UNP Q6P4G7
G	99	TRP	PHE	conflict	UNP Q6P4G7
G	101	ALA	THR	conflict	UNP Q6P4G7
G	104	LYS	GLN	conflict	UNP Q6P4G7

Continued on next page...


Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
G	106	ILE	LYS	conflict	UNP Q6P4G7
G	107	ILE	VAL	conflict	UNP Q6P4G7
G	108	LYS	GLU	conflict	UNP Q6P4G7
G	109	PRO	LEU	conflict	UNP Q6P4G7
G	110	ASP	ASN	conflict	UNP Q6P4G7
G	157	CYS	THR	conflict	UNP Q6P4G7

- Molecule 4 is a protein called TCR TRBV6-1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
4	E	242	Total	C	N	O	S	0	4	0
			1887	1191	325	362	9			
4	H	245	Total	C	N	O	S	0	10	0
			1948	1231	336	368	13			

- Molecule 5 is GLYCEROL (three-letter code: GOL) (formula: C₃H₈O₃).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	A	1	Total	C	O	0	0
			6	3	3		
5	F	1	Total	C	O	0	0
			6	3	3		
5	H	1	Total	C	O	0	0
			6	3	3		

- Molecule 6 is ACETIC ACID (three-letter code: ACY) (formula: C₂H₄O₂).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	A	1	Total C O 4 2 2	0	0
6	F	1	Total C O 4 2 2	0	0

- Molecule 7 is SODIUM ION (three-letter code: NA) (formula: Na).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
7	H	1	Total Na 1 1	0	0

- Molecule 8 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
8	A	164	Total O 164 164	0	0
8	B	53	Total O 53 53	0	0
8	C	156	Total O 156 156	0	0
8	D	77	Total O 77 77	0	0
8	E	68	Total O 68 68	0	0
8	F	82	Total O 82 82	0	0

Continued on next page...


Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
8	G	155	Total 155	O 155	0	0
8	H	202	Total 202	O 202	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Major histocompatibility complex class I-related gene protein




- Molecule 1: Major histocompatibility complex class I-related gene protein




- Molecule 2: Beta-2-microglobulin



- Molecule 2: Beta-2-microglobulin





- Molecule 3: TRA@ protein







• Molecule 3: TRA@ protein



• Molecule 4: TCR TRBV6-1



• Molecule 4: TCR TRBV6-1



4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	216.80Å 70.08Å 142.88Å 90.00° 104.55° 90.00°	Depositor
Resolution (Å)	44.18 – 2.00 44.18 – 2.00	Depositor EDS
% Data completeness (in resolution range)	98.5 (44.18-2.00) 92.0 (44.18-2.00)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.38 (at 2.00Å)	Xtrriage
Refinement program	PHENIX 1.21.1_5286	Depositor
R, R_{free}	0.173 , 0.200 0.173 , 0.200	Depositor DCC
R_{free} test set	136426 reflections (1.45%)	wwPDB-VP
Wilson B-factor (Å ²)	34.5	Xtrriage
Anisotropy	0.296	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.32 , 41.9	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	13983	wwPDB-VP
Average B, all atoms (Å ²)	49.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.72% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: GOL, NA, LLP, ACY

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.26	0/2299	0.50	0/3124
1	C	0.26	0/2295	0.52	0/3121
2	B	0.26	0/807	0.52	0/1100
2	F	0.27	0/835	0.52	0/1135
3	D	0.27	0/1543	0.51	0/2096
3	G	0.29	0/1662	0.54	1/2252 (0.0%)
4	E	0.27	0/1944	0.54	0/2652
4	H	0.28	0/2026	0.55	0/2757
All	All	0.27	0/13411	0.53	1/18237 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
4	H	0	1

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed($^{\circ}$)	Ideal($^{\circ}$)
3	G	157	CYS	CA-CB-SG	5.09	123.16	114.00

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
4	H	99	GLU	Peptide

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2234	0	2126	24	0
1	C	2236	0	2127	22	0
2	B	784	0	723	3	0
2	F	812	0	758	4	0
3	D	1498	0	1399	7	0
3	G	1600	0	1545	17	0
4	E	1887	0	1747	14	0
4	H	1948	0	1848	14	0
5	A	6	0	8	2	0
5	F	6	0	8	0	0
5	H	6	0	8	3	0
6	A	4	0	3	1	0
6	F	4	0	3	1	0
7	H	1	0	0	0	0
8	A	164	0	0	2	0
8	B	53	0	0	0	0
8	C	156	0	0	1	0
8	D	77	0	0	0	0
8	E	68	0	0	1	0
8	F	82	0	0	1	0
8	G	155	0	0	3	0
8	H	202	0	0	3	0
All	All	13983	0	12303	93	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (93) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:F:7:ILE:H	6:F:202:ACY:H3	1.41	0.84
1:A:43:LLP:O3	1:A:43:LLP:NZ	2.12	0.83
1:A:67:ARG:HH12	5:H:301:GOL:H31	1.50	0.74
3:G:19:GLN:NE2	8:G:303:HOH:O	2.21	0.73
1:C:221:ILE:HG12	1:C:223:GLN:H	1.54	0.72

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:64:GLN:HE22	5:H:301:GOL:H32	1.56	0.70
4:E:155:HIS:HB3	4:E:216:TYR:HB2	1.73	0.69
1:A:163:ALA:HB3	5:A:501:GOL:H12	1.76	0.68
1:C:43:LLP:NZ	1:C:43:LLP:O3	2.30	0.64
3:G:95:TYR:OH	8:G:301:HOH:O	2.14	0.64
1:C:99:GLU:HB2	1:C:107:THR:HG22	1.81	0.63
1:A:146:ASN:OD1	8:A:601:HOH:O	2.15	0.63
4:H:13:LEU:HD21	4:H:19[A]:MET:HG3	1.79	0.62
3:D:159:LEU:HD21	4:E:196:ARG:HB2	1.80	0.62
4:H:13:LEU:HD21	4:H:19[B]:MET:HE3	1.84	0.59
1:A:114:TYR:HB3	6:A:502:ACY:H3	1.86	0.58
4:E:111:ARG:NH1	8:E:303:HOH:O	2.35	0.58
2:F:87:LEU:O	8:F:301:HOH:O	2.17	0.57
4:H:65[A]:ASN:ND2	8:H:404:HOH:O	2.38	0.56
1:A:77:LEU:HD13	1:A:92:TYR:HB2	1.86	0.55
4:E:65[A]:ASN:OD1	4:E:77:ARG:NH1	2.37	0.55
1:A:9:ARG:NE	1:A:43:LLP:OP3	2.26	0.55
3:G:91:LYS:HG2	3:G:95:TYR:HA	1.89	0.54
4:H:174:ASP:HB2	4:H:191:LEU:HD12	1.89	0.54
3:D:28:GLY:HA3	3:D:93[B]:SER:HB3	1.90	0.53
1:C:196:THR:HB	1:C:246:LEU:HB2	1.92	0.52
1:A:113:ALA:HB2	2:B:60:TRP:CE2	2.45	0.52
3:G:28:GLY:HA3	3:G:93[A]:SER:OG	2.10	0.52
1:A:151:LEU:HD22	3:G:51:LEU:HD12	1.91	0.51
1:C:222:VAL:HA	4:H:200:THR:HG21	1.91	0.51
4:E:128:VAL:HG23	4:E:238:ALA:HB3	1.92	0.51
3:G:4[A]:ILE:HD11	3:G:90:VAL:HB	1.93	0.51
3:G:159[A]:LEU:HB3	4:H:172[A]:CYS:HB2	1.93	0.51
3:G:28:GLY:HA3	3:G:93[B]:SER:HB3	1.93	0.50
1:A:196:THR:HG22	1:A:246:LEU:HD12	1.93	0.50
1:C:215:MET:HG3	1:C:257:HIS:CD2	2.46	0.50
1:C:28:VAL:HG23	1:C:33:ILE:HD13	1.93	0.49
1:C:221:ILE:HG12	1:C:223:GLN:N	2.26	0.49
1:C:77:LEU:HD13	1:C:92:TYR:HB2	1.93	0.49
1:C:186:VAL:HG11	1:C:269:VAL:HG22	1.94	0.49
1:C:198:LEU:HD13	1:C:269:VAL:HG21	1.93	0.49
4:H:65[B]:ASN:ND2	8:H:409:HOH:O	2.45	0.49
1:A:21:GLU:OE2	1:A:74:LYS:NZ	2.40	0.49
3:G:108[B]:LYS:HB3	3:G:139:SER:HB3	1.94	0.48
1:A:216:LYS:HB2	1:A:221:ILE:HD13	1.95	0.48
1:A:163:ALA:CB	5:A:501:GOL:H12	2.42	0.48

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:F:64:LEU:HD13	2:F:66:TYR:HE1	1.78	0.48
3:G:41:GLU:OE1	8:G:302:HOH:O	2.20	0.47
4:H:120:ASN:HB2	8:H:413:HOH:O	2.13	0.47
4:E:208:HIS:HB2	4:E:241:TRP:CZ3	2.50	0.47
1:A:252:ASN:HB2	1:A:254:TYR:CE2	2.50	0.47
4:H:69:LEU:HD22	4:H:69:LEU:H	1.80	0.47
1:C:152:TYR:CD1	4:E:100:GLY:HA3	2.50	0.47
1:C:113:ALA:HB2	2:F:60:TRP:CE2	2.49	0.47
3:G:4[B]:ILE:HD11	3:G:100:GLY:HA3	1.96	0.47
1:A:67:ARG:NH1	5:H:301:GOL:H31	2.25	0.46
3:G:121:LEU:HD11	3:G:133:LEU:HB2	1.98	0.46
4:E:88:VAL:HG22	4:E:111:ARG:HD3	1.98	0.46
1:C:151:LEU:HD22	3:D:51:LEU:HD12	1.97	0.45
1:C:34:THR:HB	1:C:43:LLP:HE3	1.99	0.45
1:C:43:LLP:H5'2	8:C:1211:HOH:O	2.17	0.45
1:C:253:LEU:HD23	1:C:253:LEU:HA	1.76	0.44
2:B:54:LEU:HD11	2:B:62:PHE:HB3	1.99	0.44
3:G:138[A]:ASP:OD1	3:G:139:SER:N	2.49	0.44
4:H:128:VAL:HG23	4:H:238:ALA:HB3	2.00	0.44
3:D:1:GLY:N	3:D:27:SER:H	2.16	0.44
4:H:86:THR:HG23	4:H:113:THR:HA	1.99	0.44
1:A:154:LYS:HD3	3:G:51:LEU:HD11	1.99	0.44
4:E:83:PRO:HA	4:E:114:VAL:HB	1.99	0.44
1:C:72:MET:HB3	4:E:96:TRP:CZ2	2.53	0.43
3:D:28:GLY:HA3	3:D:93[A]:SER:OG	2.18	0.43
1:C:46:ARG:HA	1:C:46:ARG:HD3	1.77	0.43
1:C:210:ILE:O	4:H:206:ARG:HG2	2.18	0.43
4:H:131:PRO:HD3	4:H:144:LEU:HG	2.01	0.43
1:A:116:GLY:O	2:B:3:ARG:NH2	2.51	0.43
3:G:108[A]:LYS:HB3	3:G:139:SER:HB3	1.99	0.43
3:D:29:PHE:HB3	3:D:66[B]:ARG:HH21	1.83	0.43
1:A:28:VAL:HG23	1:A:33:ILE:HD13	1.99	0.43
1:A:1[A]:ARG:HD2	8:A:729:HOH:O	2.18	0.42
1:A:4:SER:HB3	1:A:99:GLU:HG2	2.02	0.42
1:A:173:LYS:HE3	1:A:173:LYS:HB2	1.68	0.42
1:C:127:LEU:HD11	1:C:154:LYS:HD2	2.02	0.41
1:C:144:GLU:HA	1:C:150:LEU:HD11	2.03	0.41
3:D:66[B]:ARG:HH11	3:D:66[B]:ARG:HB3	1.86	0.41
4:H:31:SER:HA	4:H:49:SER:O	2.21	0.41
1:A:14:ASP:HB3	1:A:89:SER:HB2	2.02	0.41
4:E:131:PRO:HD2	4:E:202:TRP:CZ2	2.56	0.41

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:G:132:CYS:SG	3:G:182[B]:CYS:HB3	2.59	0.41
4:E:46:ILE:HG22	4:E:47:TYR:HD1	1.86	0.40
3:G:125:LYS:HB3	3:G:125:LYS:HE2	1.90	0.40
4:E:181:GLN:O	4:E:187:SER:HB2	2.20	0.40
1:A:188:ARG:HG2	1:A:198:LEU:HD23	2.04	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	270/271 (100%)	265 (98%)	5 (2%)	0	100	100
1	C	267/271 (98%)	261 (98%)	6 (2%)	0	100	100
2	B	96/100 (96%)	96 (100%)	0	0	100	100
2	F	98/100 (98%)	98 (100%)	0	0	100	100
3	D	194/204 (95%)	186 (96%)	8 (4%)	0	100	100
3	G	207/204 (102%)	204 (99%)	3 (1%)	0	100	100
4	E	244/246 (99%)	239 (98%)	5 (2%)	0	100	100
4	H	253/246 (103%)	250 (99%)	3 (1%)	0	100	100
All	All	1629/1642 (99%)	1599 (98%)	30 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	237/240 (99%)	230 (97%)	7 (3%)	36	37
1	C	235/240 (98%)	226 (96%)	9 (4%)	28	28
2	B	83/95 (87%)	78 (94%)	5 (6%)	16	13
2	F	89/95 (94%)	88 (99%)	1 (1%)	70	76
3	D	160/181 (88%)	154 (96%)	6 (4%)	28	28
3	G	183/181 (101%)	177 (97%)	6 (3%)	33	33
4	E	198/212 (93%)	190 (96%)	8 (4%)	27	26
4	H	210/212 (99%)	203 (97%)	7 (3%)	33	33
All	All	1395/1456 (96%)	1346 (96%)	49 (4%)	34	31

All (49) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	1[A]	ARG
1	A	1[B]	ARG
1	A	57	ASP
1	A	73	PHE
1	A	82	ARG
1	A	221	ILE
1	A	244	ILE
2	B	16	GLU
2	B	70	PHE
2	B	88	SER
2	B	89	GLN
2	B	97	ARG
1	C	14	ASP
1	C	72	MET
1	C	73	PHE
1	C	127	LEU
1	C	221	ILE
1	C	246	LEU
1	C	253	LEU
1	C	259	GLU
1	C	261	SER
3	D	68	LYS
3	D	93[A]	SER
3	D	93[B]	SER

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
3	D	96	GLN
3	D	150	ASP
3	D	157	CYS
4	E	19	MET
4	E	79	GLU
4	E	97	THR
4	E	111	ARG
4	E	194	ARG
4	E	220	GLU
4	E	221	ASN
4	E	227	ASP
2	F	70	PHE
3	G	93[A]	SER
3	G	93[B]	SER
3	G	120	GLN
3	G	162	ARG
3	G	164	MET
3	G	199	SER
4	H	19[A]	MET
4	H	19[B]	MET
4	H	65[A]	ASN
4	H	65[B]	ASN
4	H	132	SER
4	H	194	ARG
4	H	212	GLN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (7) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	42	GLN
1	A	64	GLN
1	C	123	ASN
3	D	38	HIS
3	D	96	GLN
3	G	79	GLN
4	H	22	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains

2 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
1	LLP	A	43	1	23,24,25	1.42	3 (13%)	25,32,34	3.47	3 (12%)
1	LLP	C	43	1	23,24,25	1.21	1 (4%)	25,32,34	2.26	4 (16%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	LLP	A	43	1	-	7/16/17/19	0/1/1/1
1	LLP	C	43	1	-	2/16/17/19	0/1/1/1

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	C	43	LLP	C4'-NZ	4.89	1.43	1.27
1	A	43	LLP	C4'-NZ	4.84	1.43	1.27
1	A	43	LLP	C3-C2	2.60	1.43	1.41
1	A	43	LLP	C2'-C2	-2.34	1.46	1.50

All (7) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	43	LLP	CE-NZ-C4'	-14.67	71.72	118.72
1	C	43	LLP	CE-NZ-C4'	-9.73	87.54	118.72
1	A	43	LLP	OP4-C5'-C5	6.59	121.70	109.36
1	A	43	LLP	C4-C4'-NZ	-4.75	102.12	124.04
1	C	43	LLP	C4-C4'-NZ	-3.22	109.17	124.04
1	C	43	LLP	C4-C3-C2	-2.42	118.78	120.14
1	C	43	LLP	OP4-C5'-C5	2.39	113.84	109.36

There are no chirality outliers.

All (9) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
1	A	43	LLP	C5'-OP4-P-OP2
1	A	43	LLP	C5'-OP4-P-OP3
1	A	43	LLP	N-CA-CB-CG
1	A	43	LLP	C-CA-CB-CG
1	A	43	LLP	CA-CB-CG-CD
1	C	43	LLP	CA-CB-CG-CD
1	A	43	LLP	C5'-OP4-P-OP1
1	C	43	LLP	CG-CD-CE-NZ
1	A	43	LLP	C4-C4'-NZ-CE

There are no ring outliers.

2 monomers are involved in 5 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
1	A	43	LLP	2	0
1	C	43	LLP	3	0

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 6 ligands modelled in this entry, 1 is monoatomic - leaving 5 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
5	GOL	A	501	-	5,5,5	0.36	0	5,5,5	0.53	0
6	ACY	F	202	-	3,3,3	1.12	0	3,3,3	1.19	0
5	GOL	H	301	-	5,5,5	0.39	0	5,5,5	0.40	0
5	GOL	F	201	-	5,5,5	0.41	0	5,5,5	0.35	0
6	ACY	A	502	-	3,3,3	1.14	0	3,3,3	1.21	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	GOL	A	501	-	-	3/4/4/4	-
5	GOL	H	301	-	-	0/4/4/4	-
5	GOL	F	201	-	-	0/4/4/4	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (3) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
5	A	501	GOL	C1-C2-C3-O3
5	A	501	GOL	O2-C2-C3-O3
5	A	501	GOL	O1-C1-C2-C3

There are no ring outliers.

4 monomers are involved in 7 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
5	A	501	GOL	2	0
6	F	202	ACY	1	0
5	H	301	GOL	3	0
6	A	502	ACY	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	265/271 (97%)	-0.18	2 (0%) 82 82	18, 41, 78, 100	15 (5%)
1	C	266/271 (98%)	-0.11	5 (1%) 66 65	24, 41, 68, 88	14 (5%)
2	B	98/100 (98%)	0.11	0 100 100	34, 59, 90, 95	2 (2%)
2	F	100/100 (100%)	-0.27	0 100 100	30, 45, 67, 72	1 (1%)
3	D	194/204 (95%)	0.24	3 (1%) 71 70	24, 56, 98, 123	10 (5%)
3	G	200/204 (98%)	-0.22	1 (0%) 87 86	21, 38, 64, 79	16 (8%)
4	E	242/246 (98%)	0.19	5 (2%) 63 62	21, 56, 93, 111	8 (3%)
4	H	245/246 (99%)	-0.23	2 (0%) 82 82	22, 39, 63, 96	15 (6%)
All	All	1610/1642 (98%)	-0.06	18 (1%) 77 76	18, 44, 83, 123	81 (5%)

All (18) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	C	221	ILE	3.2
4	E	96	TRP	3.2
1	A	195	VAL	3.1
4	H	245	ASP	2.8
3	D	131	VAL	2.7
1	C	18	GLY	2.6
4	E	244	ALA	2.6
3	D	159	LEU	2.4
3	D	1	GLY	2.4
1	C	222	VAL	2.4
1	C	72	MET	2.2
3	G	200	PRO	2.2
4	E	136	ILE	2.1
1	C	69[A]	TRP	2.1
1	A	270	PRO	2.1
4	E	200	THR	2.1

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
4	E	167	VAL	2.0
4	H	99	GLU	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
1	LLP	C	43	24/25	0.93	0.10	36,46,61,74	0
1	LLP	A	43	24/25	0.94	0.10	35,38,61,68	1

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
5	GOL	H	301	6/6	0.89	0.10	33,47,54,63	0
6	ACY	F	202	4/4	0.89	0.12	27,45,55,57	0
5	GOL	A	501	6/6	0.90	0.12	47,54,61,63	0
6	ACY	A	502	4/4	0.92	0.10	40,58,60,66	0
5	GOL	F	201	6/6	0.96	0.08	35,35,38,39	0
7	NA	H	302	1/1	0.98	0.05	42,42,42,42	0

6.5 Other polymers [i](#)

There are no such residues in this entry.