



# Full wwPDB X-ray Structure Validation Report ⓘ

Sep 12, 2024 – 04:04 PM EDT

PDB ID : 9CIH  
Title : Crystal structure of human polymerase eta with incoming dCMPnPP nucleotide across O4-methyl threofuranosyl thymidine in DNA template at extension stage  
Authors : Tomar, R.; Stone, M.P.; Egli, M.  
Deposited on : 2024-07-03  
Resolution : 2.15 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 2022.3.0, CSD as543be (2022)  
Xtriage (Phenix) : 1.20.1  
EDS : 3.0  
buster-report : 1.1.7 (2018)  
Percentile statistics : 20231227.v01 (using entries in the PDB archive December 27th 2023)  
CCP4 : 9.0.002 (Gargrove)  
Density-Fitness : 1.0.11  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.38.3


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.15 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	164625	1881 (2.16-2.16)
Clashscore	180529	2047 (2.16-2.16)
Ramachandran outliers	177936	2027 (2.16-2.16)
Sidechain outliers	177891	2026 (2.16-2.16)
RSRZ outliers	164620	1882 (2.16-2.16)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	435	 2% 82% 15%
2	T	12	 58% 33% 8%
3	P	8	 12% 75% 25%

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard

residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
4	GOL	A	501	-	X	X	-

## 2 Entry composition [i](#)

There are 7 unique types of molecules in this entry. The entry contains 4096 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called DNA polymerase eta.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	424	3344	2097	601	622	24	0	4	0

There are 3 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-2	GLY	-	expression tag	UNP Q9Y253
A	-1	PRO	-	expression tag	UNP Q9Y253
A	0	HIS	-	expression tag	UNP Q9Y253

- Molecule 2 is a DNA chain called DNA (5'-D(\*CP\*AP\*TP\*GP\*(A1A0L)P\*TP\*GP\*AP\*C P\*GP\*CP\*T)-3').

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	P			
2	T	12	227	108	39	69	11	0	0	1

- Molecule 3 is a DNA chain called DNA (5'-D(\*AP\*GP\*CP\*GP\*TP\*CP\*AP\*A)-3').


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	P			
3	P	8	162	78	33	44	7	0	0	0

- Molecule 4 is GLYCEROL (three-letter code: GOL) (formula: C<sub>3</sub>H<sub>8</sub>O<sub>3</sub>) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	1	Total C O 6 3 3	0	0
4	A	1	Total C O 6 3 3	0	0
4	T	1	Total C O 6 3 3	0	0

- Molecule 5 is 2'-deoxy-5'-O-[(R)-hydroxy{[(R)-hydroxy(phosphonoxy)phosphoryl]amino}phosphoryl]cytidine (three-letter code: 0KX) (formula: C<sub>9</sub>H<sub>17</sub>N<sub>4</sub>O<sub>12</sub>P<sub>3</sub>) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
			Total	C	N	O	P		
5	A	1	28	9	4	12	3	0	0

- Molecule 6 is MAGNESIUM ION (three-letter code: MG) (formula: Mg) (labeled as "Ligand of Interest" by depositor).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	Mg		
6	A	2	2	2	0	0


- Molecule 7 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	O		
7	A	276	276	276	0	0
7	T	25	25	25	0	0
7	P	14	14	14	0	0

### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: DNA polymerase eta



- Molecule 2: DNA (5'-D(\*CP\*AP\*TP\*GP\*(A1A0L)P\*TP\*GP\*AP\*CP\*GP\*CP\*T)-3')



- Molecule 3: DNA (5'-D(\*AP\*GP\*CP\*GP\*TP\*CP\*AP\*A)-3')



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 61	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	98.47Å 98.47Å 81.90Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	42.20 – 2.15 42.20 – 2.15	Depositor EDS
% Data completeness (in resolution range)	99.9 (42.20-2.15) 99.9 (42.20-2.15)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	0.10	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.20 (at 2.16Å)	Xtrriage
Refinement program	PHENIX 1.20.1_4487	Depositor
R, $R_{free}$	0.170 , 0.232 0.170 , 0.232	Depositor DCC
$R_{free}$ test set	1309 reflections (5.25%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	21.5	Xtrriage
Anisotropy	0.003	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.35 , 49.6	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.51$ , $\langle L^2 \rangle = 0.34$	Xtrriage
Estimated twinning fraction	0.043 for h,-h-k,-l	Xtrriage
$F_o, F_c$ correlation	0.95	EDS
Total number of atoms	4096	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	27.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.76% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: MG, GOL, A1A0L, 0KX

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.40	0/3412	0.60	0/4602
2	T	0.93	0/230	1.18	1/352 (0.3%)
3	P	0.95	0/182	0.93	0/279
All	All	0.49	0/3824	0.67	1/5233 (0.0%)

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	T	7	DG	C1'-O4'-C4'	-5.36	104.74	110.10

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3344	0	3393	36	1
2	T	227	0	114	6	0
3	P	162	0	91	7	0
4	A	12	0	14	7	0
4	T	6	0	7	1	0
5	A	28	0	17	2	0
6	A	2	0	0	0	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
7	A	276	0	0	9	1
7	P	14	0	0	0	0
7	T	25	0	0	1	0
All	All	4096	0	3636	49	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.

All (49) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:126:GLN:NE2	7:A:602:HOH:O	2.05	0.88
1:A:79:GLN:OE1	7:A:601:HOH:O	1.99	0.81
3:P:7:DA:H2''	3:P:8:DA:H3'	1.70	0.72
1:A:161:VAL:HG12	1:A:162:GLN:HG3	1.73	0.70
2:T:7:DG:H21	3:P:8:DA:N6	1.93	0.67
1:A:356:ARG:NH2	7:A:605:HOH:O	2.29	0.65
1:A:42:TRP:CD1	1:A:43:LYS:HG3	2.34	0.61
1:A:383:ARG:NH2	7:A:604:HOH:O	2.28	0.60
1:A:10:ALA:HB3	1:A:119:VAL:HG13	1.84	0.60
4:T:101:GOL:O3	4:T:101:GOL:O1	2.17	0.60
4:A:501:GOL:H11	3:P:8:DA:C8	2.40	0.57
1:A:55:ARG:CZ	4:A:502:GOL:H32	2.36	0.55
1:A:111:ARG:NH1	4:A:501:GOL:O1	2.41	0.53
1:A:161:VAL:N	7:A:614:HOH:O	2.41	0.53
4:A:501:GOL:H12	2:T:7:DG:H4'	1.90	0.52
1:A:322:SER:HB3	1:A:423:PHE:CD1	2.45	0.52
1:A:369:SER:HB3	1:A:423:PHE:HB3	1.91	0.52
1:A:109:ILE:HD11	1:A:314:GLN:HG3	1.91	0.52
1:A:108:VAL:HG22	7:A:812:HOH:O	2.11	0.51
1:A:317:LYS:NZ	2:T:9:DC:OP2	2.42	0.51
1:A:81:ARG:NH2	7:A:617:HOH:O	2.43	0.51
1:A:138:SER:OG	1:A:140[B]:ASP:OD2	2.23	0.50
1:A:290:PHE:O	1:A:294:ASN:HB2	2.11	0.50
1:A:40:LYS:HD2	1:A:44:GLY:O	2.12	0.49
2:T:9:DC:H2''	2:T:10:DG:C8	2.48	0.49
1:A:372:VAL:CG1	1:A:417:PRO:HB2	2.44	0.47
1:A:189:GLN:OE1	7:A:603:HOH:O	2.21	0.47
1:A:39:TYR:O	1:A:45:GLY:HA2	2.15	0.46
1:A:119:VAL:HG22	1:A:121:LEU:HG	1.97	0.46
4:A:501:GOL:C1	2:T:7:DG:H4'	2.46	0.46

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:375:ASP:OD1	1:A:377:ARG:HG3	2.16	0.45
1:A:113:SER:HB2	3:P:8:DA:H1'	1.97	0.45
4:A:501:GOL:H31	3:P:8:DA:C8	2.52	0.45
1:A:324:ASN:OD1	1:A:421:MET:HG3	2.16	0.45
4:A:501:GOL:H31	3:P:8:DA:H8	1.82	0.44
1:A:255:ILE:HD13	1:A:275:MET:HG3	1.98	0.44
1:A:178:LEU:HD12	1:A:178:LEU:HA	1.85	0.44
1:A:323:LYS:HG2	1:A:325:PHE:CZ	2.53	0.44
2:T:8:DA:OP2	7:T:201:HOH:O	2.21	0.44
1:A:130:GLN:HA	1:A:130:GLN:OE1	2.19	0.43
5:A:503:0KX:O5'	5:A:503:0KX:H15	2.19	0.42
7:A:689:HOH:O	3:P:8:DA:H4'	2.19	0.42
5:A:503:0KX:H7	5:A:503:0KX:O2B	2.19	0.42
1:A:365:GLN:HB3	1:A:427:THR:HG23	2.01	0.41
1:A:180:ILE:HA	1:A:188:LEU:HD22	2.02	0.41
1:A:411:ILE:HD12	1:A:412:GLN:H	1.86	0.41
1:A:376:LYS:HA	1:A:376:LYS:HD2	1.94	0.41
1:A:193:GLY:O	1:A:197:VAL:HG23	2.21	0.40
1:A:352:LEU:HB3	1:A:390:TYR:CE2	2.56	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:293:LYS:NZ	7:A:601:HOH:O[6_565]	2.00	0.20

## 5.3 Torsion angles [\(i\)](#)

### 5.3.1 Protein backbone [\(i\)](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	424/435 (98%)	415 (98%)	9 (2%)	0	<b>100</b> <b>100</b>

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	367/372 (99%)	354 (96%)	13 (4%)	31 30

All (13) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	41	SER
1	A	62	SER
1	A	108	VAL
1	A	115	ASP
1	A	119	VAL
1	A	176	ASP
1	A	179	GLN
1	A	311	LYS
1	A	328	LYS
1	A	333	THR
1	A	334	ARG
1	A	409	SER
1	A	430	SER

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (5) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	79	GLN
1	A	126	GLN
1	A	135	GLN
1	A	280	GLN
1	A	405	ASN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

1 non-standard protein/DNA/RNA residue is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
2	A1A0L	T	5	2	18,21,22	3.97	10 (55%)	22,30,33	1.80	5 (22%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	A1A0L	T	5	2	-	1/7/23/24	0/2/2/2

All (10) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	T	5	A1A0L	C2'-C3'	-9.70	1.28	1.53
2	T	5	A1A0L	O4'-C1'	-6.54	1.32	1.42
2	T	5	A1A0L	O4'-C4'	5.23	1.55	1.43
2	T	5	A1A0L	C2'-C1'	4.70	1.68	1.53
2	T	5	A1A0L	O2-C2	-4.64	1.15	1.23
2	T	5	A1A0L	C4'-C3'	4.45	1.60	1.52
2	T	5	A1A0L	O4-C4	-4.43	1.23	1.34
2	T	5	A1A0L	C6-C5	3.36	1.40	1.34
2	T	5	A1A0L	C2-N1	-3.17	1.33	1.40
2	T	5	A1A0L	C6-N1	-3.07	1.32	1.38

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	T	5	A1A0L	C5-C6-N1	-4.32	118.62	123.31
2	T	5	A1A0L	C2-N3-C4	3.59	122.64	116.63
2	T	5	A1A0L	C3'-C2'-C1'	3.37	107.33	99.91
2	T	5	A1A0L	C3-C5-C6	-2.68	119.23	122.85

*Continued on next page...*

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	T	5	A1A0L	C1-O4-C4	2.02	123.29	116.84

There are no chirality outliers.

All (1) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	T	5	A1A0L	O4'-C1'-N1-C6

There are no ring outliers.

No monomer is involved in short contacts.

## 5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

Of 6 ligands modelled in this entry, 2 are monoatomic - leaving 4 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
5	OKX	A	503	6	28,29,29	3.86	19 (67%)	39,45,45	1.01	1 (2%)
4	GOL	A	502	-	5,5,5	1.17	1 (20%)	5,5,5	1.16	0
4	GOL	T	101	-	5,5,5	1.39	1 (20%)	5,5,5	1.02	0
4	GOL	A	501	-	5,5,5	1.65	2 (40%)	5,5,5	1.48	1 (20%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	OKX	A	503	6	-	4/19/34/34	0/2/2/2
4	GOL	A	502	-	-	0/4/4/4	-
4	GOL	T	101	-	-	0/4/4/4	-
4	GOL	A	501	-	-	4/4/4/4	-

All (23) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
5	A	503	OKX	C3'-C4'	-7.12	1.34	1.53
5	A	503	OKX	C6-C5	6.47	1.50	1.35
5	A	503	OKX	C2-N3	6.28	1.48	1.36
5	A	503	OKX	O4'-C4'	6.06	1.58	1.45
5	A	503	OKX	C2'-C1'	-5.35	1.37	1.52
5	A	503	OKX	PB-O3B	4.83	1.65	1.59
5	A	503	OKX	C4-N3	4.80	1.44	1.34
5	A	503	OKX	O4'-C1'	4.77	1.53	1.42
5	A	503	OKX	PB-O2B	4.68	1.53	1.46
5	A	503	OKX	C4-N4	4.39	1.44	1.33
5	A	503	OKX	C2'-C3'	4.20	1.63	1.52
5	A	503	OKX	C2-N1	3.86	1.48	1.40
5	A	503	OKX	PB-N3A	3.68	1.73	1.63
5	A	503	OKX	O2-C2	-3.58	1.17	1.23
5	A	503	OKX	C6-N1	3.56	1.46	1.38
5	A	503	OKX	PA-O1A	2.94	1.50	1.46
5	A	503	OKX	C5-C4	2.93	1.49	1.42
4	T	101	GOL	O2-C2	-2.90	1.35	1.43
5	A	503	OKX	PA-O5'	2.89	1.68	1.57
4	A	501	GOL	O2-C2	-2.77	1.35	1.43
5	A	503	OKX	PA-N3A	2.33	1.69	1.63
4	A	501	GOL	O3-C3	-2.09	1.33	1.42
4	A	502	GOL	O2-C2	-2.05	1.37	1.43

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
5	A	503	OKX	O1A-PA-N3A	-3.07	107.25	111.77
4	A	501	GOL	C3-C2-C1	-2.62	102.19	111.80

There are no chirality outliers.

All (8) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	A	501	GOL	C1-C2-C3-O3
5	A	503	0KX	PB-O3B-PG-O1G
5	A	503	0KX	PB-N3A-PA-O1A
4	A	501	GOL	O2-C2-C3-O3
4	A	501	GOL	O1-C1-C2-C3
4	A	501	GOL	O1-C1-C2-O2
5	A	503	0KX	C5'-O5'-PA-O2A
5	A	503	0KX	PB-O3B-PG-O2G



There are no ring outliers.



4 monomers are involved in 10 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
5	A	503	0KX	2	0
4	A	502	GOL	1	0
4	T	101	GOL	1	0
4	A	501	GOL	6	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.







## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	424/435 (97%)	-0.25	8 (1%) 66 70	10, 22, 54, 79	4 (0%)
2	T	11/12 (91%)	-0.45	0 100 100	24, 29, 46, 47	0
3	P	8/8 (100%)	-0.12	1 (12%) 9 11	26, 32, 35, 51	0
All	All	443/455 (97%)	-0.25	9 (2%) 64 69	10, 23, 54, 79	4 (0%)

All (9) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
3	P	8	DA	3.1
1	A	406	CYS	2.5
1	A	413	THR	2.3
1	A	181	ASP	2.2
1	A	180	ILE	2.1
1	A	431	ALA	2.1
1	A	408	THR	2.1
1	A	161	VAL	2.1
1	A	378	LEU	2.0

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q < 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
2	A1A0L	T	5	20/21	0.96	0.06	18,23,26,27	0

### 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

### 6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
4	GOL	T	101	6/6	0.85	0.13	23,27,38,38	0
4	GOL	A	501	6/6	0.90	0.18	15,25,38,42	0
4	GOL	A	502	6/6	0.92	0.11	17,26,36,44	0
5	OKX	A	503	28/28	0.98	0.05	9,15,20,21	0
6	MG	A	504	1/1	0.98	0.02	13,13,13,13	0
6	MG	A	505	1/1	0.98	0.03	13,13,13,13	0


The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

**Electron density around GOL T 101:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)















**Electron density around 0KX A 503:**


$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)



**Electron density around MG A 504:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)





## 6.5 Other polymers [i](#)

There are no such residues in this entry.