



Full wwPDB X-ray Structure Validation Report ⓘ

Jan 20, 2026 – 10:06 AM EST

PDB ID : 9DAY / pdb_00009day
Title : Crystal Structure of the Transcriptional Regulator HcpR from Porphyromonas Gingivalis
Authors : Musayev, F.N.; Escalante, C.R.; Belvin, B.R.; Lewis, J.P.
Deposited on : 2024-08-22
Resolution : 2.30 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
buster-report	:	1.1.7 (2018)
Percentile statistics	:	20231227.v01 (using entries in the PDB archive December 27th 2023)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.47


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.30 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	164625	5963 (2.30-2.30)
Clashscore	180529	6698 (2.30-2.30)
Ramachandran outliers	177936	6640 (2.30-2.30)
Sidechain outliers	177891	6640 (2.30-2.30)
RSRZ outliers	164620	5963 (2.30-2.30)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	242	19% 88% 10% .
1	B	242	29% 81% 12% 6%

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 3673 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Transcriptional regulator, putative.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	237	Total	C	N	O	S	0	0	0
			1841	1177	312	344	8			
1	B	228	Total	C	N	O	S	0	0	0
			1626	1034	277	307	8			

There are 32 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	-13	MET	-	initiating methionine	UNP Q7MVK4
A	-12	ASP	-	expression tag	UNP Q7MVK4
A	-11	HIS	-	expression tag	UNP Q7MVK4
A	-10	HIS	-	expression tag	UNP Q7MVK4
A	-9	HIS	-	expression tag	UNP Q7MVK4
A	-8	HIS	-	expression tag	UNP Q7MVK4
A	-7	HIS	-	expression tag	UNP Q7MVK4
A	-6	HIS	-	expression tag	UNP Q7MVK4
A	-5	GLU	-	expression tag	UNP Q7MVK4
A	-4	ASN	-	expression tag	UNP Q7MVK4
A	-3	LEU	-	expression tag	UNP Q7MVK4
A	-2	TYR	-	expression tag	UNP Q7MVK4
A	-1	PHE	-	expression tag	UNP Q7MVK4
A	0	GLN	-	expression tag	UNP Q7MVK4
A	1	GLY	-	expression tag	UNP Q7MVK4
A	2	SER	-	expression tag	UNP Q7MVK4
B	-13	MET	-	initiating methionine	UNP Q7MVK4
B	-12	ASP	-	expression tag	UNP Q7MVK4
B	-11	HIS	-	expression tag	UNP Q7MVK4
B	-10	HIS	-	expression tag	UNP Q7MVK4
B	-9	HIS	-	expression tag	UNP Q7MVK4
B	-8	HIS	-	expression tag	UNP Q7MVK4
B	-7	HIS	-	expression tag	UNP Q7MVK4
B	-6	HIS	-	expression tag	UNP Q7MVK4
B	-5	GLU	-	expression tag	UNP Q7MVK4

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
B	-4	ASN	-	expression tag	UNP Q7MVK4
B	-3	LEU	-	expression tag	UNP Q7MVK4
B	-2	TYR	-	expression tag	UNP Q7MVK4
B	-1	PHE	-	expression tag	UNP Q7MVK4
B	0	GLN	-	expression tag	UNP Q7MVK4
B	1	GLY	-	expression tag	UNP Q7MVK4
B	2	SER	-	expression tag	UNP Q7MVK4

- Molecule 2 is GLYCEROL (CCD ID: GOL) (formula: $C_3H_8O_3$) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	C	O	0	0
			6	3	3		
2	A	1	Total	C	O	0	0
			6	3	3		
2	B	1	Total	C	O	0	0
			6	3	3		

- Molecule 3 is FORMIC ACID (CCD ID: FMT) (formula: CH_2O_2) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O	0	0
			3	1	2		
3	A	1	Total	C	O	0	0
			3	1	2		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	104	Total	O	0	0
			104	104		
4	B	78	Total	O	0	0
			78	78		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Transcriptional regulator, putative



- Molecule 1: Transcriptional regulator, putative



4 Data and refinement statistics

Property	Value	Source
Space group	P 41 21 2	Depositor
Cell constants a, b, c, α , β , γ	106.28Å 106.28Å 116.77Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	26.57 – 2.30 26.57 – 2.30	Depositor EDS
% Data completeness (in resolution range)	98.6 (26.57-2.30) 98.5 (26.57-2.30)	Depositor EDS
R_{merge}	0.05	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.39 (at 2.31Å)	Xtriage
Refinement program	PHENIX (1.20.1_4487: ???)	Depositor
R, R_{free}	0.215 , 0.256 0.215 , 0.255	Depositor DCC
R_{free} test set	1554 reflections (5.12%)	wwPDB-VP
Wilson B-factor (Å ²)	39.0	Xtriage
Anisotropy	0.427	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 59.4	EDS
L-test for twinning ²	$\langle L \rangle = 0.51$, $\langle L^2 \rangle = 0.35$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	3673	wwPDB-VP
Average B, all atoms (Å ²)	62.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.02% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality ⓘ

5.1 Standard geometry ⓘ

Bond lengths and bond angles in the following residue types are not validated in this section: GOL, FMT

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.47	2/1871 (0.1%)	0.75	2/2519 (0.1%)
1	B	0.42	2/1645 (0.1%)	0.67	3/2218 (0.1%)
All	All	0.45	4/3516 (0.1%)	0.71	5/4737 (0.1%)

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	183	LYS	C-N	-7.71	1.23	1.33
1	A	183	LYS	C-N	-7.70	1.23	1.33
1	A	186	SER	C-N	-5.56	1.26	1.33
1	B	186	SER	C-N	-5.54	1.26	1.33

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	175	ARG	O-C-N	-6.85	113.16	122.48
1	B	187	ASP	O-C-N	5.41	127.85	122.12
1	A	187	ASP	O-C-N	5.38	127.82	122.12
1	B	184	GLU	O-C-N	5.26	127.70	122.12
1	A	184	GLU	O-C-N	5.20	127.63	122.12

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within

the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1841	0	1850	20	0
1	B	1626	0	1538	18	0
2	A	12	0	16	3	0
2	B	6	0	8	0	0
3	A	6	0	2	0	0
4	A	104	0	0	3	0
4	B	78	0	0	2	0
All	All	3673	0	3414	35	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (35) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:85:ARG:HE	2:A:301:GOL:H11	1.43	0.83
1:A:67:GLU:HB3	1:A:78:ILE:HD13	1.64	0.78
1:B:116:LYS:NZ	4:B:401:HOH:O	2.26	0.67
1:B:52:LYS:NZ	4:B:403:HOH:O	2.32	0.62
1:B:0:GLN:HG3	1:B:31:GLU:HG2	1.83	0.61
1:B:144:LEU:O	1:B:148:ILE:HG12	2.02	0.59
1:A:167:LEU:HD13	1:A:176:ILE:HG21	1.84	0.58
1:A:65:ARG:HG2	1:A:65:ARG:HH11	1.70	0.56
1:A:167:LEU:HB2	1:A:176:ILE:HD13	1.87	0.56
1:B:21:MET:HE1	1:B:129:LEU:HB2	1.88	0.55
1:A:97:ASN:HB3	1:B:146:LYS:HG3	1.90	0.54
1:A:27:LEU:HG	1:A:31:GLU:HG3	1.90	0.53
1:A:40:LEU:HD12	1:A:64:ILE:HD12	1.91	0.53
1:A:65:ARG:HG2	1:A:65:ARG:NH1	2.25	0.51
1:A:9:LEU:HD13	1:A:30:LEU:HD12	1.94	0.50
1:A:85:ARG:NE	2:A:301:GOL:H11	2.21	0.48
1:B:9:LEU:HG	1:B:13:LYS:HE3	1.94	0.48
1:A:70:GLY:HA3	1:B:149:HIS:CE1	2.51	0.46
1:B:41:LYS:O	1:B:44:GLU:HB3	2.16	0.46
1:A:147:LYS:HG3	1:B:148:ILE:HD12	1.98	0.45
1:B:186:SER:CB	1:B:193:ARG:HA	2.46	0.45
1:B:208:ILE:HG22	1:B:209:ILE:HG23	1.97	0.45
1:A:116:LYS:NZ	4:A:403:HOH:O	2.39	0.44
1:B:9:LEU:HD12	1:B:30:LEU:HD12	2.00	0.44
1:A:68:MET:CE	1:A:101:VAL:HG22	2.48	0.43

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:167:LEU:HD13	1:A:176:ILE:CG2	2.48	0.43
1:B:27:LEU:HD22	1:B:31:GLU:OE2	2.18	0.43
1:B:52:LYS:HE2	1:B:98:ILE:HG22	2.00	0.43
1:B:41:LYS:NZ	1:B:44:GLU:HB2	2.33	0.43
1:A:86:ILE:O	2:A:301:GOL:H12	2.19	0.43
1:A:31:GLU:HG2	4:A:454:HOH:O	2.19	0.42
1:B:12:TRP:CD1	1:B:26:LEU:HD11	2.55	0.42
1:B:38:GLU:OE1	1:B:113:ARG:HD2	2.21	0.41
1:A:41:LYS:HD3	1:A:41:LYS:HA	1.76	0.40
1:A:34:SER:O	4:A:402:HOH:O	2.22	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	235/242 (97%)	233 (99%)	2 (1%)	0	100	100
1	B	222/242 (92%)	216 (97%)	6 (3%)	0	100	100
All	All	457/484 (94%)	449 (98%)	8 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	196/211 (93%)	196 (100%)	0	100	100
1	B	152/211 (72%)	150 (99%)	2 (1%)	65	79
All	All	348/422 (82%)	346 (99%)	2 (1%)	84	92

All (2) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	33	CYS
1	B	151	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	150	GLN
1	B	149	HIS

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

5 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	FMT	A	303	-	2,2,2	0.78	0	1,1,1	0.18	0
2	GOL	A	301	-	5,5,5	0.88	0	5,5,5	1.10	1 (20%)
2	GOL	A	302	-	5,5,5	0.95	0	5,5,5	1.01	0
3	FMT	A	304	-	2,2,2	0.71	0	1,1,1	0.23	0
2	GOL	B	301	-	5,5,5	0.94	0	5,5,5	1.04	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	GOL	A	301	-	-	2/4/4/4	-
2	GOL	A	302	-	-	2/4/4/4	-
2	GOL	B	301	-	-	4/4/4/4	-

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	301	GOL	C3-C2-C1	-2.04	104.33	111.80

There are no chirality outliers.

All (8) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
2	B	301	GOL	C1-C2-C3-O3
2	A	301	GOL	C1-C2-C3-O3
2	A	302	GOL	O1-C1-C2-C3
2	B	301	GOL	O2-C2-C3-O3
2	A	302	GOL	O1-C1-C2-O2
2	A	301	GOL	O2-C2-C3-O3
2	B	301	GOL	O1-C1-C2-C3
2	B	301	GOL	O1-C1-C2-O2



There are no ring outliers.


1 monomer is involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	301	GOL	3	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.







5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ > 2		OWAB(Å ²)	Q < 0.9
1	A	237/242 (97%)	0.68	45 (18%)	4 5	27, 49, 120, 130	0
1	B	228/242 (94%)	1.16	71 (31%)	1 1	26, 50, 139, 158	0
All	All	465/484 (96%)	0.91	116 (24%)	2 2	26, 50, 134, 158	0

All (116) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	176	ILE	6.8
1	B	182	TRP	6.7
1	B	227	LEU	6.3
1	B	167	LEU	6.2
1	B	174	ASN	5.8
1	B	169	THR	5.7
1	B	218	ILE	5.7
1	B	190	GLY	5.6
1	B	185	LEU	5.5
1	B	164	LEU	5.4
1	B	2	SER	5.3
1	B	211	VAL	5.3
1	B	192	ASN	5.2
1	B	189	PHE	5.1
1	B	180	SER	5.1
1	B	-1	PHE	5.0
1	A	182	TRP	5.0
1	B	166	GLN	5.0
1	B	203	LEU	4.8
1	B	173	SER	4.8
1	B	162	ASP	4.7
1	B	212	ASP	4.7
1	A	216	ILE	4.6
1	B	219	LEU	4.5

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	176	ILE	4.5
1	A	177	VAL	4.4
1	B	194	GLN	4.4
1	B	163	TYR	4.4
1	B	228	GLU	4.4
1	B	220	GLN	4.3
1	B	216	ILE	4.3
1	B	168	TYR	4.3
1	A	214	LYS	4.1
1	B	195	SER	4.0
1	A	173	SER	4.0
1	B	226	ARG	4.0
1	A	189	PHE	4.0
1	B	186	SER	3.9
1	A	98	ILE	3.8
1	B	165	PHE	3.8
1	B	160	ILE	3.8
1	B	171	ASP	3.7
1	B	184	GLU	3.7
1	B	200	LEU	3.6
1	A	97	ASN	3.6
1	A	213	GLY	3.6
1	A	194	GLN	3.6
1	A	211	VAL	3.6
1	B	225	SER	3.5
1	B	224	LEU	3.4
1	B	209	ILE	3.3
1	A	196	LEU	3.3
1	B	210	ARG	3.3
1	B	0	GLN	3.2
1	B	222	ASN	3.1
1	A	212	ASP	3.1
1	B	181	SER	3.1
1	A	185	LEU	3.1
1	B	223	ARG	3.1
1	A	200	LEU	3.1
1	A	210	ARG	3.1
1	B	208	ILE	3.0
1	B	215	SER	3.0
1	A	215	SER	3.0
1	A	46	TYR	2.9
1	B	188	ARG	2.9

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	193	ARG	2.9
1	A	69	VAL	2.9
1	B	183	LYS	2.9
1	B	193	ARG	2.9
1	B	156	LEU	2.9
1	B	3	PRO	2.8
1	A	174	ASN	2.8
1	B	155	SER	2.8
1	B	214	LYS	2.8
1	B	172	GLY	2.8
1	B	159	LYS	2.7
1	A	219	LEU	2.7
1	A	70	GLY	2.6
1	A	186	SER	2.6
1	A	195	SER	2.6
1	B	177	VAL	2.6
1	A	218	ILE	2.6
1	B	179	GLU	2.5
1	B	137	ILE	2.5
1	A	-8	HIS	2.5
1	A	160	ILE	2.5
1	A	183	LYS	2.4
1	B	221	PRO	2.4
1	B	213	GLY	2.4
1	A	187	ASP	2.4
1	A	175	ARG	2.4
1	B	161	GLY	2.3
1	A	178	VAL	2.3
1	B	1	GLY	2.3
1	B	205	GLU	2.3
1	A	190	GLY	2.2
1	A	228	GLU	2.2
1	B	196	LEU	2.2
1	A	163	TYR	2.2
1	A	197	ALA	2.2
1	A	191	VAL	2.2
1	A	180	SER	2.2
1	A	156	LEU	2.2
1	B	53	LEU	2.2
1	B	178	VAL	2.2
1	B	217	GLU	2.1
1	A	192	ASN	2.1

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	188	ARG	2.1
1	B	158	GLY	2.1
1	A	184	GLU	2.1
1	A	32	SER	2.0
1	B	199	SER	2.0
1	A	170	LYS	2.0
1	B	197	ALA	2.0
1	B	202	GLN	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
3	FMT	A	303	3/3	0.79	0.18	68,68,69,69	0
2	GOL	A	302	6/6	0.85	0.17	50,57,59,70	0
2	GOL	A	301	6/6	0.85	0.14	41,48,56,62	0
2	GOL	B	301	6/6	0.91	0.12	44,46,49,51	0
3	FMT	A	304	3/3	0.94	0.09	42,42,43,45	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.




Electron density around FMT A 303:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around GOL A 302:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)




**Electron density around GOL A 301:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around GOL B 301:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around FMT A 304:

$2mF_o - DF_c$ (at 0.7 rmsd) in gray
 $mF_o - DF_c$ (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.