



Full wwPDB X-ray Structure Validation Report ⓘ

Oct 28, 2025 – 08:03 AM EDT

PDB ID : 9DCT / pdb_00009dct
Title : Structure of the TelB-associated type VII secretion system chaperone SIR_0178
Authors : Gkrakopoulou, P.; Kim, Y.; Whitney, J.C.
Deposited on : 2024-08-27
Resolution : 3.24 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20231227.v01 (using entries in the PDB archive December 27th 2023)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.46


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 3.24 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	164625	1999 (3.28-3.20)
Clashscore	180529	2147 (3.28-3.20)
Ramachandran outliers	177936	2118 (3.28-3.20)
Sidechain outliers	177891	2117 (3.28-3.20)
RSRZ outliers	164620	2001 (3.28-3.20)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	251	
1	B	251	
1	C	251	
1	D	251	

2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 7319 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called DUF4176 domain-containing protein.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	228	Total	C	N	O	S	0	0	0
			1848	1210	282	350	6			
1	B	225	Total	C	N	O	S	0	0	0
			1826	1196	279	345	6			
1	C	221	Total	C	N	O	S	0	0	0
			1799	1181	274	338	6			
1	D	228	Total	C	N	O	S	0	0	0
			1845	1208	282	348	7			

There are 56 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-13	MET	-	initiating methionine	UNP T1ZCG6
A	-12	GLY	-	expression tag	UNP T1ZCG6
A	-11	SER	-	expression tag	UNP T1ZCG6
A	-10	SER	-	expression tag	UNP T1ZCG6
A	-9	HIS	-	expression tag	UNP T1ZCG6
A	-8	HIS	-	expression tag	UNP T1ZCG6
A	-7	HIS	-	expression tag	UNP T1ZCG6
A	-6	HIS	-	expression tag	UNP T1ZCG6
A	-5	HIS	-	expression tag	UNP T1ZCG6
A	-4	HIS	-	expression tag	UNP T1ZCG6
A	-3	SER	-	expression tag	UNP T1ZCG6
A	-2	GLN	-	expression tag	UNP T1ZCG6
A	-1	ASP	-	expression tag	UNP T1ZCG6
A	0	PRO	-	expression tag	UNP T1ZCG6
B	-13	MET	-	initiating methionine	UNP T1ZCG6
B	-12	GLY	-	expression tag	UNP T1ZCG6
B	-11	SER	-	expression tag	UNP T1ZCG6
B	-10	SER	-	expression tag	UNP T1ZCG6
B	-9	HIS	-	expression tag	UNP T1ZCG6
B	-8	HIS	-	expression tag	UNP T1ZCG6
B	-7	HIS	-	expression tag	UNP T1ZCG6

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
B	-6	HIS	-	expression tag	UNP T1ZCG6
B	-5	HIS	-	expression tag	UNP T1ZCG6
B	-4	HIS	-	expression tag	UNP T1ZCG6
B	-3	SER	-	expression tag	UNP T1ZCG6
B	-2	GLN	-	expression tag	UNP T1ZCG6
B	-1	ASP	-	expression tag	UNP T1ZCG6
B	0	PRO	-	expression tag	UNP T1ZCG6
C	-13	MET	-	initiating methionine	UNP T1ZCG6
C	-12	GLY	-	expression tag	UNP T1ZCG6
C	-11	SER	-	expression tag	UNP T1ZCG6
C	-10	SER	-	expression tag	UNP T1ZCG6
C	-9	HIS	-	expression tag	UNP T1ZCG6
C	-8	HIS	-	expression tag	UNP T1ZCG6
C	-7	HIS	-	expression tag	UNP T1ZCG6
C	-6	HIS	-	expression tag	UNP T1ZCG6
C	-5	HIS	-	expression tag	UNP T1ZCG6
C	-4	HIS	-	expression tag	UNP T1ZCG6
C	-3	SER	-	expression tag	UNP T1ZCG6
C	-2	GLN	-	expression tag	UNP T1ZCG6
C	-1	ASP	-	expression tag	UNP T1ZCG6
C	0	PRO	-	expression tag	UNP T1ZCG6
D	-13	MET	-	initiating methionine	UNP T1ZCG6
D	-12	GLY	-	expression tag	UNP T1ZCG6
D	-11	SER	-	expression tag	UNP T1ZCG6
D	-10	SER	-	expression tag	UNP T1ZCG6
D	-9	HIS	-	expression tag	UNP T1ZCG6
D	-8	HIS	-	expression tag	UNP T1ZCG6
D	-7	HIS	-	expression tag	UNP T1ZCG6
D	-6	HIS	-	expression tag	UNP T1ZCG6
D	-5	HIS	-	expression tag	UNP T1ZCG6
D	-4	HIS	-	expression tag	UNP T1ZCG6
D	-3	SER	-	expression tag	UNP T1ZCG6
D	-2	GLN	-	expression tag	UNP T1ZCG6
D	-1	ASP	-	expression tag	UNP T1ZCG6
D	0	PRO	-	expression tag	UNP T1ZCG6


- Molecule 2 is CHLORIDE ION (CCD ID: CL) (formula: Cl).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	1	Total Cl 1 1	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


• Molecule 1: DUF4176 domain-containing protein



• Molecule 1: DUF4176 domain-containing protein




• Molecule 1: DUF4176 domain-containing protein



• Molecule 1: DUF4176 domain-containing protein





LEU
ARG
LYS
GLU
LEU
LYS

4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	67.81Å 69.85Å 253.94Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	46.34 – 3.24 46.34 – 3.24	Depositor EDS
% Data completeness (in resolution range)	92.0 (46.34-3.24) 92.6 (46.34-3.24)	Depositor EDS
R_{merge}	0.17	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.83 (at 3.25Å)	Xtriage
Refinement program	PHENIX 1.20.1_4487	Depositor
R, R_{free}	0.259 , 0.308 0.259 , 0.310	Depositor DCC
R_{free} test set	941 reflections (4.70%)	wwPDB-VP
Wilson B-factor (Å ²)	123.5	Xtriage
Anisotropy	0.571	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.33 , 104.0	EDS
L-test for twinning ²	$\langle L \rangle = 0.46$, $\langle L^2 \rangle = 0.29$	Xtriage
Estimated twinning fraction	0.043 for k,h,-l	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	7319	wwPDB-VP
Average B, all atoms (Å ²)	148.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.37% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: CL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.08	0/1890	0.24	0/2566
1	B	0.08	0/1869	0.24	0/2538
1	C	0.07	0/1841	0.22	0/2499
1	D	0.08	0/1888	0.23	0/2562
All	All	0.08	0/7488	0.23	0/10165

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1848	0	1831	30	0
1	B	1826	0	1807	20	0
1	C	1799	0	1784	21	0
1	D	1845	0	1832	33	0
2	A	1	0	0	0	0
All	All	7319	0	7254	96	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.

All (96) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:120:LEU:HD22	1:B:178:LEU:HD21	1.79	0.63
1:C:59:GLN:HG2	1:C:65:SER:HB2	1.83	0.61
1:A:43:LYS:NZ	1:A:179:ILE:O	2.29	0.60
1:D:114:GLU:O	1:D:180:LYS:N	2.31	0.59
1:B:155:ASP:HB2	1:B:175:SER:HA	1.84	0.59
1:D:50:LYS:NZ	1:D:212:ALA:O	2.35	0.59
1:C:227:THR:HG22	1:C:231:LEU:HD13	1.86	0.57
1:D:155:ASP:HB2	1:D:175:SER:HA	1.87	0.56
1:D:140:GLY:N	1:D:157:PHE:O	2.38	0.56
1:C:203:GLN:NE2	1:C:208:GLN:O	2.40	0.55
1:A:48:LEU:HD12	1:A:53:GLN:HB2	1.89	0.55
1:D:10:PHE:HZ	1:D:99:ILE:HG12	1.72	0.55
1:B:108:PRO:HG3	1:B:188:GLU:HG3	1.89	0.54
1:C:115:LEU:HD21	1:C:174:VAL:HG22	1.90	0.54
1:B:195:MET:HG3	1:D:162:PRO:HG2	1.90	0.53
1:B:106:LEU:O	1:B:196:LYS:NZ	2.29	0.52
1:D:145:LEU:HD22	1:D:152:TYR:HB2	1.91	0.52
1:B:48:LEU:HD12	1:B:53:GLN:HB2	1.92	0.52
1:A:155:ASP:HB2	1:A:175:SER:HA	1.90	0.52
1:D:203:GLN:HG3	1:D:208:GLN:HB2	1.91	0.52
1:D:120:LEU:HD21	1:D:178:LEU:HD21	1.92	0.52
1:A:187:LEU:HD13	1:C:184:HIS:HB2	1.92	0.51
1:A:127:MET:HB3	1:D:120:LEU:HD22	1.94	0.50
1:D:115:LEU:HD21	1:D:174:VAL:HG22	1.94	0.50
1:D:145:LEU:HD23	1:D:149:PHE:HB3	1.93	0.50
1:B:111:THR:HG23	1:B:185:MET:HA	1.93	0.50
1:A:154:ILE:HG21	1:A:173:PHE:HD2	1.77	0.49
1:C:147:LYS:HD3	1:C:148:ASP:N	2.28	0.49
1:D:52:LYS:HA	1:D:70:ARG:HH12	1.76	0.49
1:A:173:PHE:HE1	1:D:170:PRO:HB3	1.78	0.49
1:D:17:TRP:HH2	1:D:204:LEU:HA	1.78	0.48
1:B:143:LEU:HB2	1:B:154:ILE:HB	1.95	0.48
1:D:17:TRP:HE1	1:D:209:LEU:HD21	1.78	0.48
1:A:125:GLN:HG3	1:A:129:GLY:C	2.39	0.48
1:D:77:LEU:HD13	1:D:86:ILE:HG12	1.95	0.48
1:B:106:LEU:HG	1:B:142:ARG:HH22	1.79	0.47
1:A:225:TYR:HD1	1:A:231:LEU:HD11	1.79	0.47
1:C:17:TRP:HB3	1:C:20:VAL:HG23	1.96	0.47
1:D:48:LEU:O	1:D:70:ARG:NH1	2.47	0.47
1:B:121:ASP:HB3	1:B:124:LEU:HB3	1.96	0.47

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:70:ARG:HB2	1:D:76:ILE:HD13	1.96	0.47
1:D:48:LEU:HD12	1:D:53:GLN:HB2	1.97	0.46
1:C:104:GLY:O	1:C:200:ARG:NH2	2.36	0.46
1:A:137:MET:HE3	1:A:161:TRP:HD1	1.81	0.46
1:A:166:VAL:HG12	1:A:168:ASN:H	1.79	0.46
1:A:191:TRP:CG	1:B:229:PRO:HB3	2.50	0.46
1:C:192:GLU:HG3	1:C:196:LYS:HE3	1.96	0.46
1:A:229:PRO:HG3	1:B:191:TRP:CD2	2.51	0.46
1:D:69:GLU:N	1:D:69:GLU:OE1	2.49	0.46
1:C:231:LEU:HD12	1:C:231:LEU:HA	1.83	0.46
1:A:153:GLU:N	1:A:209:LEU:O	2.46	0.46
1:C:161:TRP:CE2	1:C:162:PRO:HB3	2.50	0.46
1:D:142:ARG:HG2	1:D:199:LEU:HD11	1.98	0.46
1:D:4:TYR:HE1	1:D:9:ILE:HD12	1.81	0.46
1:A:128:LEU:HD21	1:D:170:PRO:HB2	1.98	0.45
1:C:49:ILE:HD12	1:C:89:GLU:HG3	1.98	0.45
1:B:47:LYS:HG2	1:B:119:LEU:HD21	1.99	0.45
1:D:57:GLU:HB3	1:D:67:LEU:HD13	1.97	0.45
1:D:115:LEU:HB2	1:D:134:ALA:HB1	1.98	0.45
1:B:142:ARG:HA	1:B:142:ARG:HD3	1.71	0.45
1:A:70:ARG:HB2	1:A:76:ILE:HD12	1.99	0.44
1:A:9:ILE:HG22	1:A:13:VAL:HG23	1.99	0.44
1:B:87:TRP:HB3	1:B:92:PHE:HB2	2.00	0.44
1:A:117:LEU:HA	1:A:120:LEU:HD12	1.99	0.44
1:B:4:TYR:CZ	1:B:200:ARG:HD3	2.52	0.44
1:A:165:GLU:OE1	1:C:159:ARG:NH2	2.51	0.43
1:C:145:LEU:HG	1:C:146:ALA:H	1.83	0.43
1:A:145:LEU:HD23	1:A:149:PHE:HB3	1.99	0.43
1:B:112:VAL:HB	1:B:184:HIS:HB3	1.99	0.43
1:A:159:ARG:HG3	1:A:165:GLU:HA	1.99	0.42
1:C:57:GLU:HB3	1:C:65:SER:OG	2.19	0.42
1:D:49:ILE:HG23	1:D:89:GLU:HG3	2.00	0.42
1:A:7:VAL:HG13	1:A:33:ILE:HD13	2.01	0.42
1:A:59:GLN:H	1:A:59:GLN:HG2	1.50	0.42
1:C:210:SER:HB3	1:C:213:PHE:CD2	2.55	0.42
1:A:153:GLU:O	1:A:211:THR:N	2.52	0.42
1:B:46:TYR:CZ	1:B:50:LYS:HG3	2.54	0.42
1:D:169:ILE:HD13	1:D:169:ILE:HA	1.80	0.41
1:C:115:LEU:HB3	1:C:120:LEU:HD21	2.02	0.41
1:C:228:THR:HA	1:C:231:LEU:HB2	2.01	0.41
1:D:81:LYS:HE2	1:D:81:LYS:HB3	1.95	0.41

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:48:LEU:HD12	1:C:53:GLN:HB2	2.03	0.41
1:A:154:ILE:HG21	1:A:173:PHE:CD2	2.56	0.41
1:A:47:LYS:HE3	1:A:51:LEU:HD12	2.02	0.41
1:D:12:GLU:H	1:D:12:GLU:HG3	1.69	0.41
1:C:1:MET:HE3	1:C:1:MET:HB2	1.86	0.41
1:D:64:LEU:H	1:D:64:LEU:HG	1.50	0.41
1:D:117:LEU:HB2	1:D:129:GLY:HA2	2.03	0.40
1:A:58:TYR:O	1:A:65:SER:HA	2.21	0.40
1:B:44:GLY:HA3	1:B:56:TYR:CZ	2.56	0.40
1:D:17:TRP:HA	1:D:18:PRO:HD3	1.96	0.40
1:A:19:THR:HB	1:A:94:LEU:HD21	2.03	0.40
1:A:83:LYS:HE2	1:A:83:LYS:HB3	1.83	0.40
1:B:156:TYR:HB2	1:B:174:VAL:HB	2.04	0.40
1:A:118:GLU:H	1:A:118:GLU:HG3	1.57	0.40
1:C:219:GLN:O	1:C:223:LEU:HG	2.22	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	224/251 (89%)	210 (94%)	14 (6%)	0	100	100
1	B	221/251 (88%)	213 (96%)	8 (4%)	0	100	100
1	C	217/251 (86%)	209 (96%)	8 (4%)	0	100	100
1	D	224/251 (89%)	213 (95%)	11 (5%)	0	100	100
All	All	886/1004 (88%)	845 (95%)	41 (5%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	202/223 (91%)	184 (91%)	18 (9%)	8	29
1	B	199/223 (89%)	185 (93%)	14 (7%)	12	39
1	C	196/223 (88%)	186 (95%)	10 (5%)	20	50
1	D	202/223 (91%)	186 (92%)	16 (8%)	10	35
All	All	799/892 (90%)	741 (93%)	58 (7%)	11	38

All (58) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	19	THR
1	A	24	VAL
1	A	27	ASP
1	A	59	GLN
1	A	60	SER
1	A	62	LEU
1	A	64	LEU
1	A	65	SER
1	A	83	LYS
1	A	118	GLU
1	A	127	MET
1	A	130	GLU
1	A	150	GLU
1	A	157	PHE
1	A	177	MET
1	A	190	GLU
1	A	223	LEU
1	A	231	LEU
1	B	8	GLU
1	B	9	ILE
1	B	24	VAL
1	B	25	THR
1	B	30	GLU
1	B	73	ASP

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	106	LEU
1	B	124	LEU
1	B	142	ARG
1	B	166	VAL
1	B	169	ILE
1	B	174	VAL
1	B	192	GLU
1	B	225	TYR
1	C	2	THR
1	C	3	GLU
1	C	27	ASP
1	C	73	ASP
1	C	157	PHE
1	C	166	VAL
1	C	168	ASN
1	C	188	GLU
1	C	214	MET
1	C	231	LEU
1	D	12	GLU
1	D	13	VAL
1	D	19	THR
1	D	24	VAL
1	D	61	PHE
1	D	64	LEU
1	D	118	GLU
1	D	124	LEU
1	D	127	MET
1	D	128	LEU
1	D	135	LEU
1	D	141	ARG
1	D	142	ARG
1	D	147	LYS
1	D	166	VAL
1	D	169	ILE

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (6) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	125	GLN
1	A	184	HIS
1	A	219	GLN
1	B	75	GLN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	D	53	GLN
1	D	168	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 1 ligands modelled in this entry, 1 is monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	228/251 (90%)	-0.56	2 (0%) 81 69	106, 148, 184, 197	0
1	B	225/251 (89%)	-0.58	0 100 100	94, 130, 166, 206	0
1	C	221/251 (88%)	-0.67	0 100 100	110, 140, 178, 209	0
1	D	228/251 (90%)	-0.54	0 100 100	102, 164, 207, 223	0
All	All	902/1004 (89%)	-0.59	2 (0%) 92 88	94, 145, 195, 223	0

All (2) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	138	LEU	2.1
1	A	124	LEU	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
2	CL	A	301	1/1	0.84	0.16	138,138,138,138	0

6.5 Other polymers [i](#)

There are no such residues in this entry.