



# Full wwPDB X-ray Structure Validation Report ⓘ

Feb 3, 2025 – 08:26 pm GMT

PDB ID : 9GC5  
Title : Highly optimized CNS penetrant inhibitors of EGFR Exon20 Insertion Mutations  
Authors : Hargreaves, D.  
Deposited on : 2024-08-01  
Resolution : 1.91 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.4, CSD as541be (2020)  
Xtrriage (Phenix) : 1.13  
EDS : 3.0  
buster-report : 1.1.7 (2018)  
Percentile statistics : 20231227.v01 (using entries in the PDB archive December 27th 2023)  
CCP4 : 9.0.003 (Gargrove)  
Density-Fitness : 1.0.11  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.40


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 1.91 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	164625	7293 (1.90-1.90)
Clashscore	180529	8090 (1.90-1.90)
Ramachandran outliers	177936	8022 (1.90-1.90)
Sidechain outliers	177891	8022 (1.90-1.90)
RSRZ outliers	164620	7292 (1.90-1.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	330	 6% 81% 10% 9%
1	B	330	 9% 85% 5% 9%

## 2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 5088 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Epidermal growth factor receptor.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	300	2416	1550	410	438	18	0	0	0
1	B	300	2416	1550	410	438	18	0	0	0

There are 6 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	693	GLY	-	expression tag	UNP P00533
A	694	SER	-	expression tag	UNP P00533
A	948	ARG	VAL	engineered mutation	UNP P00533
B	693	GLY	-	expression tag	UNP P00533
B	694	SER	-	expression tag	UNP P00533
B	948	ARG	VAL	engineered mutation	UNP P00533

- Molecule 2 is 1-[2-[5-(1,3-benzoxazol-4-yl)-2,4-bis(fluoranyl)phenyl]-3-pyrimidin-4-yl-4,6-dihydropyrrolo[3,4-d]imidazol-5-yl]propan-1-one (three-letter code: A1IZ8) (formula: C<sub>25</sub>H<sub>18</sub>F<sub>2</sub>N<sub>6</sub>O<sub>2</sub>) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
			Total	C	F	N	O		
2	A	1	35	25	2	6	2	0	0
2	B	1	35	25	2	6	2	0	0

- Molecule 3 is SULFATE ION (three-letter code: SO<sub>4</sub>) (formula: O<sub>4</sub>S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
			Total	O	S		
3	A	1	5	4	1	0	0


- Molecule 4 is water.

<b>Mol</b>	<b>Chain</b>	<b>Residues</b>	<b>Atoms</b>		<b>ZeroOcc</b>	<b>AltConf</b>
4	A	81	Total 81	O 81	0	0
4	B	100	Total 100	O 100	0	0


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Epidermal growth factor receptor



- Molecule 1: Epidermal growth factor receptor



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	35.87Å 76.15Å 77.97Å 66.88° 83.58° 87.00°	Depositor
Resolution (Å)	16.38 – 1.91 16.38 – 1.91	Depositor EDS
% Data completeness (in resolution range)	63.5 (16.38-1.91) 63.4 (16.38-1.91)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.05 (at 1.91Å)	Xtrriage
Refinement program	BUSTER 2.11.8 (26-JUL-2023)	Depositor
R, $R_{free}$	0.220 , 0.261 0.215 , 0.259	Depositor DCC
$R_{free}$ test set	23166 reflections (4.84%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	31.5	Xtrriage
Anisotropy	0.052	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.36 , 39.7	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.51$ , $\langle L^2 \rangle = 0.34$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
$F_o, F_c$ correlation	0.93	EDS
Total number of atoms	5088	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	37.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 10.76% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality

### 5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: SO4, A1IZ8

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.41	0/2469	0.58	0/3341
1	B	0.44	0/2469	0.60	0/3341
All	All	0.43	0/4938	0.59	0/6682

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2416	0	2456	17	0
1	B	2416	0	2456	13	0
2	A	35	0	0	0	0
2	B	35	0	0	0	0
3	A	5	0	0	0	0
4	A	81	0	0	0	0
4	B	100	0	0	0	0
All	All	5088	0	4912	26	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.



All (26) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:988:HIS:CE1	1:B:990:PRO:HA	2.05	0.91
1:B:949:LYS:HB3	1:B:959:PRO:HD3	1.82	0.62
1:A:737:LYS:NZ	1:B:804:GLU:HG2	2.15	0.61
1:A:737:LYS:HZ2	1:B:804:GLU:HG2	1.67	0.60
1:A:811:SER:OG	1:A:975:PRO:HB2	2.10	0.52
1:A:724:GLY:HA2	1:A:748:ARG:HG3	1.92	0.50
1:A:705:ARG:HG2	1:A:707:LEU:HD13	1.95	0.49
1:A:860:LYS:O	1:A:861:LEU:HB2	2.12	0.48
1:A:842:ASN:OD1	1:A:855:ASP:OD2	2.31	0.47
1:B:962:ARG:CZ	1:B:962:ARG:HB2	2.42	0.47
1:B:811:SER:OG	1:B:975:PRO:HB2	2.15	0.47
1:B:829:GLU:HA	1:B:893:HIS:CE1	2.50	0.47
1:A:1001:LEU:HD13	1:B:778:LEU:HD23	1.97	0.46
1:B:731:TRP:CZ2	1:B:733:PRO:HB3	2.50	0.46
1:B:989:LEU:HD12	1:B:989:LEU:H	1.81	0.46
1:B:882:ALA:HA	1:B:898:TRP:CD2	2.51	0.46
1:A:737:LYS:NZ	1:B:804:GLU:CG	2.79	0.45
1:B:960:LYS:HB2	1:B:963:GLU:HG3	1.99	0.43
1:A:882:ALA:HA	1:A:898:TRP:CD2	2.54	0.43
1:A:926:ILE:HA	1:A:929:LYS:HE3	2.01	0.43
1:A:926:ILE:HG23	1:A:931:GLU:HB2	2.02	0.42
1:A:769:VAL:O	1:A:776:ARG:HB3	2.20	0.41
1:A:941:ILE:HD12	1:A:941:ILE:HA	1.98	0.41
1:A:967:GLU:O	1:A:971:MET:HG3	2.20	0.41
1:A:718:LEU:HD12	1:A:726:VAL:HG12	2.02	0.40
1:A:835:HIS:O	1:A:836:ARG:HB2	2.21	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	296/330 (90%)	285 (96%)	10 (3%)	1 (0%)	37	29
1	B	296/330 (90%)	288 (97%)	8 (3%)	0	100	100
All	All	592/660 (90%)	573 (97%)	18 (3%)	1 (0%)	44	36

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	990	PRO

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	269/289 (93%)	262 (97%)	7 (3%)	41	36
1	B	269/289 (93%)	263 (98%)	6 (2%)	47	43
All	All	538/578 (93%)	525 (98%)	13 (2%)	44	39

All (13) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	705	ARG
1	A	720	SER
1	A	757	LYS
1	A	778	LEU
1	A	793	MET
1	A	913	LYS
1	A	987	MET
1	B	757	LYS
1	B	778	LEU
1	B	804	GLU
1	B	808	ASN
1	B	970	LYS
1	B	989	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (8) such

sidechains are listed below:

Mol	Chain	Res	Type
1	A	773	HIS
1	A	982	GLN
1	B	773	HIS
1	B	805	HIS
1	B	808	ASN
1	B	849	GLN
1	B	888	HIS
1	B	988	HIS

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

3 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
3	SO4	A	1102	-	4,4,4	0.18	0	6,6,6	0.63	0
2	A1IZ8	B	1101	1	31,40,40	0.60	0	34,59,59	1.08	2 (5%)
2	A1IZ8	A	1101	1	31,40,40	0.61	0	34,59,59	0.98	2 (5%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral

centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	A1IZ8	B	1101	1	-	4/15/26/26	0/6/6/6
2	A1IZ8	A	1101	1	-	4/15/26/26	0/6/6/6

There are no bond length outliers.

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed( $^{\circ}$ )	Ideal( $^{\circ}$ )
2	B	1101	A1IZ8	C18-N3-C13	3.84	129.74	124.57
2	A	1101	A1IZ8	C18-N3-C13	3.45	129.21	124.57
2	B	1101	A1IZ8	C9-C10-C12	3.29	123.65	118.41
2	A	1101	A1IZ8	C9-C10-C12	2.79	122.85	118.41

There are no chirality outliers.

All (8) torsion outliers are listed below:



Mol	Chain	Res	Type	Atoms
2	B	1101	A1IZ8	N1-C13-C3-C4
2	B	1101	A1IZ8	N2-C22-C23-C24
2	A	1101	A1IZ8	N1-C13-C3-C4
2	B	1101	A1IZ8	N1-C13-C3-C2
2	A	1101	A1IZ8	N2-C22-C23-C24
2	A	1101	A1IZ8	C-C5-C6-C7
2	A	1101	A1IZ8	N1-C13-C3-C2
2	B	1101	A1IZ8	O1-C22-C23-C24

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier.

The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	300/330 (90%)	0.56	21 (7%) 24 25	20, 36, 64, 80	0
1	B	300/330 (90%)	0.53	29 (9%) 15 15	19, 34, 61, 81	0
All	All	600/660 (90%)	0.54	50 (8%) 19 20	19, 35, 64, 81	0

All (50) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	750	ALA	7.3
1	A	750	ALA	7.2
1	A	989	LEU	6.6
1	B	989	LEU	6.3
1	B	753	PRO	5.6
1	B	988	HIS	5.2
1	A	751	THR	5.1
1	B	699	PRO	4.5
1	A	755	ALA	4.4
1	B	861	LEU	4.3
1	B	751	THR	4.1
1	B	1011	VAL	4.1
1	A	988	HIS	4.0
1	B	754	LYS	4.0
1	A	990	PRO	3.6
1	A	753	PRO	3.5
1	B	807	ASP	3.5
1	B	755	ALA	3.4
1	A	1011	VAL	3.3
1	A	754	LYS	3.2
1	B	749	GLU	3.1
1	B	917	GLY	3.1
1	A	699	PRO	2.9
1	A	861	LEU	2.7

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	B	990	PRO	2.7
1	B	747	LEU	2.6
1	B	808	ASN	2.6
1	B	727	TYR	2.5
1	A	1010	VAL	2.5
1	B	719	GLY	2.4
1	A	858	LEU	2.4
1	A	737	LYS	2.4
1	A	856	PHE	2.3
1	B	809	ILE	2.3
1	B	918	ILE	2.3
1	B	981	ILE	2.3
1	B	987	MET	2.3
1	A	717	VAL	2.3
1	B	1010	VAL	2.2
1	B	962	ARG	2.2
1	B	985	GLU	2.2
1	A	757	LYS	2.2
1	A	1012	ASP	2.1
1	B	717	VAL	2.1
1	B	876	VAL	2.1
1	A	1007	MET	2.0
1	A	918	ILE	2.0
1	A	749	GLU	2.0
1	B	762	GLU	2.0
1	B	748	ARG	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [\(i\)](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [\(i\)](#)

There are no monosaccharides in this entry.


## 6.4 Ligands [\(i\)](#)


In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.



Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
2	A1IZ8	A	1101	35/35	0.90	0.09	32,33,35,36	0
2	A1IZ8	B	1101	35/35	0.92	0.07	27,30,32,34	0
3	SO4	A	1102	5/5	0.93	0.10	62,63,63,63	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.





## 6.5 Other polymers [i](#)

There are no such residues in this entry.