



Full wwPDB X-ray Structure Validation Report ⓘ

Sep 8, 2025 – 02:16 pm BST

PDB ID : 9GPD / pdb_00009gpd
Title : ManDH5 E303Q mutant in complex with mannotriose a beta-D-Mannanase of GH5 family from Dictyoglomus thermophilum
Authors : Sivron, Y.; Romano, A.; Shoham, Y.; Shoham, G.
Deposited on : 2024-09-07
Resolution : 1.35 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	FAILED
Mogul	:	1.8.4, CSD as541be (2020)
Xtriage (Phenix)	:	2.0rc1
EDS	:	3.0
Percentile statistics	:	20231227.v01 (using entries in the PDB archive December 27th 2023)
CCP4	:	9.0.006 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.45.1


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.35 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	164625	1089 (1.36-1.36)
RSRZ outliers	164620	1088 (1.36-1.36)

MolProbity failed to run properly - the sequence quality summary graphics cannot be shown.

2 Entry composition [i](#)

There are 3 unique types of molecules in this entry. The entry contains 10115 atoms, of which 4623 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called DUF5060 domain-containing protein.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	563	Total	C	H	N	O	S	0	9	0
			9356	3131	4591	770	855	9			

There is a discrepancy between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	303	GLN	GLU	engineered mutation	UNP A0A7C3MIF0

- Molecule 2 is an oligosaccharide called beta-D-mannopyranose-(1-4)-beta-D-mannopyranose-(1-4)-beta-D-mannopyranose.



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
2	B	3	Total	C	H	O	0	0	0
			66	18	32	16			

- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	684	Total	O	0	9
			693	693		

MolProbity failed to run properly - this section is therefore empty.

3 Data and refinement statistics

Property	Value	Source
Space group	I 2 2 2	Depositor
Cell constants a, b, c, α , β , γ	95.78Å 99.11Å 153.84Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	83.30 – 1.35 83.30 – 1.35	Depositor EDS
% Data completeness (in resolution range)	99.7 (83.30-1.35) 99.6 (83.30-1.35)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.37 (at 1.35Å)	Xtriage
Refinement program	PHENIX 1.14_3228	Depositor
R, R_{free}	0.156 , 0.167 0.159 , 0.171	Depositor DCC
R_{free} test set	7938 reflections (4.83%)	wwPDB-VP
Wilson B-factor (Å ²)	13.1	Xtriage
Anisotropy	0.337	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.39 , 43.9	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	0.009 for k,h,-l	Xtriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	10115	wwPDB-VP
Average B, all atoms (Å ²)	21.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.72% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

4 Model quality [i](#)

4.1 Standard geometry [i](#)

MolProbity failed to run properly - this section is therefore empty.

4.2 Too-close contacts [i](#)

MolProbity failed to run properly - this section is therefore empty.

4.3 Torsion angles [i](#)

4.3.1 Protein backbone [i](#)

MolProbity failed to run properly - this section is therefore empty.

4.3.2 Protein sidechains [i](#)

MolProbity failed to run properly - this section is therefore empty.

4.3.3 RNA [i](#)

MolProbity failed to run properly - this section is therefore empty.

4.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

4.5 Carbohydrates [i](#)

3 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	BMA	B	1	2	12,12,12	2.36	5 (41%)	17,17,17	1.71	5 (29%)
2	BMA	B	2	2	11,11,12	1.52	2 (18%)	15,15,17	2.15	7 (46%)
2	BMA	B	3	2	11,11,12	1.56	3 (27%)	15,15,17	2.18	3 (20%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	BMA	B	1	2	-	0/2/22/22	0/1/1/1
2	BMA	B	2	2	-	0/2/19/22	0/1/1/1
2	BMA	B	3	2	-	0/2/19/22	0/1/1/1

All (10) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	1	BMA	C1-C2	4.70	1.63	1.52
2	B	1	BMA	O2-C2	-3.13	1.35	1.43
2	B	2	BMA	C2-C3	2.94	1.56	1.52
2	B	1	BMA	C3-C2	2.87	1.59	1.52
2	B	3	BMA	C2-C3	2.79	1.56	1.52
2	B	1	BMA	O5-C5	-2.64	1.37	1.44
2	B	3	BMA	C1-C2	2.50	1.57	1.52
2	B	2	BMA	O2-C2	-2.44	1.38	1.43
2	B	3	BMA	C4-C5	2.42	1.58	1.53
2	B	1	BMA	C4-C3	2.01	1.57	1.52

All (15) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	3	BMA	O5-C5-C6	5.39	115.65	107.20
2	B	3	BMA	C1-O5-C5	-4.81	105.67	112.19
2	B	2	BMA	C1-O5-C5	4.62	118.46	112.19
2	B	1	BMA	O5-C1-C2	3.88	117.21	110.28
2	B	2	BMA	C2-C3-C4	2.96	116.02	110.89
2	B	2	BMA	C1-C2-C3	-2.95	106.05	109.67
2	B	1	BMA	O2-C2-C3	2.85	116.93	110.35
2	B	2	BMA	O4-C4-C3	-2.66	104.19	110.35
2	B	1	BMA	O1-C1-O5	-2.53	102.79	110.38
2	B	2	BMA	O2-C2-C1	-2.40	104.23	109.15

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	1	BMA	C1-C2-C3	-2.26	105.63	110.31
2	B	2	BMA	C6-C5-C4	2.22	118.20	113.00
2	B	3	BMA	O2-C2-C1	-2.19	104.67	109.15
2	B	1	BMA	O4-C4-C5	-2.18	103.88	109.30
2	B	2	BMA	O5-C5-C4	-2.04	105.87	110.83


There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.



4.6 Ligand geometry [i](#)

There are no ligands in this entry.

4.7 Other polymers [i](#)

There are no such residues in this entry.

4.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.

5 Fit of model and data

5.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	563/568 (99%)	0.22	60 (10%) 12 15	8, 17, 38, 53	7 (1%)

All (60) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	403	TRP	6.7
1	A	160	CYS	5.6
1	A	419	ILE	5.4
1	A	124[A]	TYR	5.3
1	A	521	ILE	5.0
1	A	474	ASN	4.8
1	A	507	ASN	4.7
1	A	18	SER	4.4
1	A	510	SER	4.3
1	A	466	THR	4.2
1	A	520	LEU	4.2
1	A	509	ASP	3.9
1	A	511	SER	3.7
1	A	20	MET	3.6
1	A	508	MET	3.6
1	A	519	LEU	3.5
1	A	563	LEU	3.4
1	A	17	ILE	3.3
1	A	538	ILE	3.3
1	A	505	ILE	3.3
1	A	544	ASP	3.3
1	A	465	LEU	3.2
1	A	564	LEU	3.2
1	A	467	ASN	3.1
1	A	463	GLU	3.1
1	A	107	LYS	3.1
1	A	543	LEU	3.0

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	200	ARG	2.9
1	A	468	TYR	2.8
1	A	398	GLY	2.8
1	A	21	GLU	2.8
1	A	502	LYS	2.7
1	A	57	TYR	2.7
1	A	496	LYS	2.7
1	A	113	GLY	2.7
1	A	523	PRO	2.7
1	A	19	LYS	2.7
1	A	482	LEU	2.6
1	A	454	ASP	2.6
1	A	475	ARG	2.5
1	A	522	LYS	2.5
1	A	119	ASP	2.4
1	A	422	GLY	2.4
1	A	545	LYS	2.4
1	A	470	PHE	2.4
1	A	481	GLY	2.4
1	A	3	ARG	2.3
1	A	473	TYR	2.3
1	A	547	VAL	2.3
1	A	4	TYR	2.3
1	A	22	LEU	2.2
1	A	7	ASN	2.2
1	A	462	PHE	2.2
1	A	524	PRO	2.1
1	A	117	SER	2.1
1	A	62	ARG	2.1
1	A	455	ILE	2.1
1	A	400	ASN	2.0
1	A	480	ILE	2.0
1	A	2	SER	2.0

5.2 Non-standard residues in protein, DNA, RNA chains [i](#)


There are no non-standard protein/DNA/RNA residues in this entry.

5.3 Carbohydrates [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
2	BMA	B	3	11/12	0.86	0.13	28,34,40,46	0
2	BMA	B	2	11/12	0.93	0.10	18,25,30,36	0
2	BMA	B	1	12/12	0.94	0.10	12,27,35,41	0

The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.



5.4 Ligands [i](#)

There are no ligands in this entry.

5.5 Other polymers [i](#)

There are no such residues in this entry.