

Full wwPDB X-ray Structure Validation Report (i)

Sep 22, 2025 – 02:11 PM JST

PDB ID : 9JNG / pdb 00009jng

Title : Crystal Structure of SME-1 E166A mutant with Tebipenem

Authors: Dhankhar, K.; Hazra, S.

Deposited on : 2024-09-23

Resolution : 2.26 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
https://www.wwpdb.org/validation/2017/XrayValidationReportHelp
with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (1)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0

Mogul : 1.8.5 (274361), CSD as541be (2020)

Xtriage (Phenix) : 2.0 EDS : 3.0

buster-report : 1.1.7 (2018)

Percentile statistics : 20231227.v01 (using entries in the PDB archive December 27th 2023)

CCP4 : 9.0.010 (Gargrove)

Density-Fitness : 1.0.12

Ideal geometry (proteins) : Engh & Huber (2001) Ideal geometry (DNA, RNA) : Parkinson et al. (1996)


Validation Pipeline (wwPDB-VP) : 2.46

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: $X\text{-}RAY\ DIFFRACTION$

The reported resolution of this entry is 2.26 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive $(\# \mathrm{Entries})$	$\begin{array}{c} {\rm Similar\ resolution} \\ (\#{\rm Entries},{\rm resolution\ range}({\rm \AA})) \end{array}$
R_{free}	164625	1763 (2.26-2.26)
Clashscore	180529	1919 (2.26-2.26)
Ramachandran outliers	177936	1884 (2.26-2.26)
Sidechain outliers	177891	1885 (2.26-2.26)
RSRZ outliers	164620	1763 (2.26-2.26)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain		
1	A	275	81%	14%	• • •
1	В	275	81%	13%	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:



Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	DGE	A	302	X	-	-	-
3	DGE	В	303	X	-	-	-



2 Entry composition (i)

There are 6 unique types of molecules in this entry. The entry contains 4339 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

• Molecule 1 is a protein called beta-lactamase.


\mathbf{Mol}	Chain	Residues	Atoms			ZeroOcc	AltConf	Trace		
1	A	267	Total 2073	C 1296	N 369	O 401	S 7	0	2	0
1	В	267	Total 2062	C 1290	N 365	O 400	S 7	0	1	0

There are 18 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-1	MET	-	initiating methionine	UNP Q54488
A	0	GLY	-	expression tag	UNP Q54488
A	143	ALA	GLU	engineered mutation	UNP Q54488
A	268	HIS	-	expression tag	UNP Q54488
A	269	HIS	-	expression tag	UNP Q54488
A	270	HIS	-	expression tag	UNP Q54488
A	271	HIS	-	expression tag	UNP Q54488
A	272	HIS	-	expression tag	UNP Q54488
A	273	HIS	-	expression tag	UNP Q54488
В	-1	MET	-	initiating methionine	UNP Q54488
В	0	GLY	-	expression tag	UNP Q54488
В	143	ALA	GLU	engineered mutation	UNP Q54488
В	268	HIS	-	expression tag	UNP Q54488
В	269	HIS	-	expression tag	UNP Q54488
В	270	HIS	_	expression tag	UNP Q54488
В	271	HIS	-	expression tag	UNP Q54488
В	272	HIS	-	expression tag	UNP Q54488
В	273	HIS	_	expression tag	UNP Q54488


• Molecule 2 is DI(HYDROXYETHYL)ETHER (CCD ID: PEG) (formula: $C_4H_{10}O_3$) (labeled as "Ligand of Interest" by depositor).





Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	1	Total C O 7 4 3	0	0
2	В	1	Total C O 7 4 3	0	0

• Molecule 3 is $(2S,3R,4S)-4-\{[1-(4,5-dihydro-1,3-thiazol-2-yl)azetidin-3-yl]sulfanyl\}-2-[(2S,3R)-3-hydroxy-1-oxobutan-2-yl]-3-methyl-3,4-dihydro-2H-pyrrole-5-carboxylic acid (CCD ID: DGE) (formula: <math>C_{16}H_{23}N_3O_4S_2$) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf		
9	Λ	1	Total	С	N	О	S	0	0
3	A	1	25	16	3	4	2	U	U

Continued on next page...




Continued from previous page...

Mo	Chain	Residues		Ato	$\mathbf{m}\mathbf{s}$			ZeroOcc	AltConf
9	D	1	Total	С	N	О	S	0	0
)	D	1	25	16	3	4	2	U	0

• Molecule 4 is CHLORIDE ION (CCD ID: CL) (formula: Cl) (labeled as "Ligand of Interest" by depositor).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	2	Total Cl 2 2	0	0

• Molecule 5 is 1,2-ETHANEDIOL (CCD ID: EDO) (formula: $C_2H_6O_2$) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	В	1	Total 4	C 2	O 2	0	0

• Molecule 6 is water.


Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	A	65	Total O 65 65	0	0
6	В	69	Total O 69 69	0	0



3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: beta-lactamase





4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants	71.77Å 51.69Å 76.40Å	Donositor
a, b, c, α , β , γ	90.00° 114.22° 90.00°	Depositor
Resolution (Å)	23.29 - 2.26	Depositor
rtesolution (A)	23.29 - 2.26	EDS
% Data completeness	99.8 (23.29-2.26)	Depositor
(in resolution range)	99.8 (23.29-2.26)	EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$< I/\sigma(I) > 1$	3.11 (at 2.26Å)	Xtriage
Refinement program	REFMAC 5.8.0425	Depositor
D D.	0.179 , 0.270	Depositor
R, R_{free}	0.181 , 0.272	DCC
R_{free} test set	1169 reflections (4.83%)	wwPDB-VP
Wilson B-factor (Å ²)	22.3	Xtriage
Anisotropy	0.522	Xtriage
Bulk solvent $k_{sol}(e/Å^3)$, $B_{sol}(Å^2)$	0.36, 38.3	EDS
L-test for twinning ²	$ < L >=0.48, < L^2>=0.31$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	4339	wwPDB-VP
Average B, all atoms (Å ²)	26.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: The analyses of the Patterson function reveals a significant off-origin peak that is 30.51 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 1.2919e-03. The detected translational NCS is most likely also responsible for the elevated intensity ratio.

²Theoretical values of <|L|>, $<L^2>$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



¹Intensities estimated from amplitudes.

5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: DGE, PEG, EDO, CL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond	lengths	Bo	ond angles
MIOI	Chain	RMSZ	# Z > 5	RMSZ	# Z > 5
1	A	0.60	0/2108	1.28	$11/2842 \ (0.4\%)$
1	В	0.61	0/2097	1.30	$11/2828 \ (0.4\%)$
All	All	0.61	0/4205	1.29	$22/5670 \ (0.4\%)$

There are no bond length outliers.

All (22) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	\mathbf{Z}	$Observed(^o)$	$\mathrm{Ideal}(^{o})$
1	В	230	LYS	CB-CA-C	11.00	127.49	109.89
1	В	174	ASN	CB-CA-C	-9.96	93.33	110.17
1	A	77	ASP	CA-CB-CG	7.08	119.68	112.60
1	В	87	LYS	CB-CA-C	6.76	123.13	110.63
1	В	20	ARG	CB-CA-C	6.05	120.08	109.80
1	В	202	VAL	N-CA-CB	6.04	118.51	111.87
1	A	72	LYS	CG-CD-CE	5.75	124.53	111.30
1	A	20	ARG	CB-CA-C	5.75	118.63	110.24
1	A	249	LYS	CB-CA-C	5.75	119.21	109.50
1	A	202	VAL	N-CA-CB	5.72	116.53	111.67
1	В	27	ASP	CA-CB-CG	5.63	118.23	112.60
1	В	244	LYS	N-CA-CB	5.55	118.06	110.01
1	В	87	LYS	CB-CG-CD	5.49	123.92	111.30
1	В	243	ARG	CD-NE-CZ	5.29	131.81	124.40
1	A	15	GLU	CB-CG-CD	5.22	121.47	112.60
1	В	87	LYS	CG-CD-CE	5.18	123.22	111.30
1	A	174	ASN	CA-CB-CG	-5.14	107.46	112.60
1	В	249	LYS	CB-CA-C	5.10	118.23	109.51
1	A	153	ASP	CA-CB-CG	5.06	117.66	112.60
1	A	27	ASP	CA-CB-CG	5.02	117.62	112.60
1	A	145[A]	GLU	CB-CG-CD	5.01	121.11	112.60
1	A	145[B]	GLU	CB-CG-CD	5.01	121.11	112.60



There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2073	0	2060	21	1
1	В	2062	0	2048	22	0
2	A	7	0	10	2	0
2	В	7	0	10	1	0
3	A	25	0	0	0	0
3	В	25	0	0	1	0
4	A	2	0	0	0	0
5	В	4	0	6	1	0
6	A	65	0	0	5	1
6	В	69	0	0	4	0
All	All	4339	0	4134	46	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

All (46) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	$\begin{array}{c} {\rm Interatomic} \\ {\rm distance} \ ({\rm \AA}) \end{array}$	Clash overlap (Å)
1:B:141:ARG:HB3	1:B:145[B]:GLU:HG3	1.48	0.95
1:A:66:ASP:H	1:A:69:GLN:HE21	1.28	0.80
1:B:66:ASP:H	1:B:69:GLN:HE21	1.26	0.80
1:B:141:ARG:HB3	1:B:145[B]:GLU:CG	2.16	0.73
1:A:2:LYS:CB	6:A:465:HOH:O	2.37	0.73
1:A:112:ASN:O	1:A:116:GLU:HG2	1.90	0.71
1:B:112:ASN:O	1:B:116:GLU:HG2	1.92	0.69
1:A:141:ARG:HB3	1:A:145[A]:GLU:HG3	1.74	0.68
1:B:66:ASP:H	1:B:69:GLN:NE2	1.90	0.68
1:A:66:ASP:H	1:A:69:GLN:NE2	1.92	0.67
1:B:76:ARG:HD3	1:B:78:LEU:HD23	1.77	0.65
1:A:76:ARG:HD3	1:A:78:LEU:HD23	1.78	0.64
2:B:301:PEG:H21	5:B:302:EDO:H11	1.81	0.62

Continued on next page...



Continued from previous page...

A		Interatomic	Clash
Atom-1	Atom-2	distance (Å)	overlap (Å)
1:B:67:ILE:HG22	6:B:452:HOH:O	2.01	0.61
1:B:17:PHE:CE2	1:B:243:ARG:HD2	2.40	0.57
1:B:82:SER:HB3	1:B:85:THR:OG1	2.07	0.55
1:A:17:PHE:CE2	1:A:243:ARG:HD2	2.41	0.55
1:A:82:SER:HB3	1:A:85:THR:OG1	2.07	0.54
1:A:88:TYR:OH	1:B:153:ASP:HB2	2.09	0.52
1:B:169:LYS:NZ	6:B:403:HOH:O	2.42	0.52
1:B:136:GLU:HG3	1:B:162:ALA:HB2	1.91	0.51
1:A:61:GLN:HB2	1:A:180:VAL:HG21	1.95	0.49
1:B:17:PHE:CD2	1:B:243:ARG:HD2	2.48	0.48
1:B:36:TYR:CE1	1:B:37:ARG:HG3	2.49	0.47
1:B:129:MET:HG3	1:B:137:PHE:CD1	2.51	0.46
1:A:129:MET:HG3	1:A:137:PHE:CD1	2.50	0.45
1:B:139:LEU:HD11	1:B:146:LEU:HD22	1.99	0.44
1:B:49:LYS:HB2	1:B:211:LYS:HE3	2.00	0.44
1:A:17:PHE:CD2	1:A:243:ARG:HD2	2.53	0.44
1:A:136:GLU:HG3	1:A:162:ALA:HB2	2.00	0.44
2:A:301:PEG:H22	6:A:410:HOH:O	2.18	0.43
1:B:183:ILE:HG13	6:B:407:HOH:O	2.18	0.43
1:B:138:ARG:O	1:B:156:ASP:HA	2.18	0.43
2:A:301:PEG:C2	6:A:410:HOH:O	2.66	0.43
1:A:49:LYS:HB2	1:A:211:LYS:HE3	2.00	0.43
1:B:264:GLN:O	1:B:267:ASP:HB2	2.18	0.42
1:A:17:PHE:HB3	6:A:443:HOH:O	2.19	0.42
1:A:132:ILE:HD13	1:A:170:LEU:HD21	2.01	0.42
1:B:11:LYS:HE2	6:B:468:HOH:O	2.20	0.42
1:A:138:ARG:O	1:A:156:ASP:HA	2.20	0.42
3:B:303:DGE:CAD	3:B:303:DGE:CAC	2.97	0.42
1:B:36:TYR:CD1	1:B:37:ARG:HG3	2.55	0.41
1:A:158:SER:N	6:A:407:HOH:O	2.45	0.41
1:A:103:LEU:C	1:A:103:LEU:HD23	2.46	0.41
1:A:264:GLN:O	1:A:267:ASP:HB2	2.21	0.41
1:A:17:PHE:O	1:A:243:ARG:NH1	2.52	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1 Atom-2		$\begin{array}{c} {\rm Interatomic} \\ {\rm distance} \ ({\rm \AA}) \end{array}$	Clash overlap (Å)
1:A:260[A]:ARG:NH2	6:A:461:HOH:O[2_445]	2.13	0.07



5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Perce	\mathbf{ntiles}
1	A	$267/275 \ (97\%)$	263 (98%)	4 (2%)	0	100	100
1	В	$266/275 \ (97\%)$	262 (98%)	4 (2%)	0	100	100
All	All	533/550 (97%)	525 (98%)	8 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles		
1	A	$216/223 \ (97\%)$	210 (97%)	6 (3%)	38 47		
1	В	215/223 (96%)	204 (95%)	11 (5%)	20 21		
All	All	431/446 (97%)	414 (96%)	17 (4%)	29 33		

All (17) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	17	PHE
1	A	70	LYS
1	A	74	GLU
1	A	119	LEU
1	A	136	GLU
1	A	243	ARG
1	В	8	LYS
1	В	17	PHE

Continued on next page...



Continued from previous page...

Mol	Chain	Res	Type
1	В	70	LYS
1	В	119	LEU
1	В	123	GLU
1	В	136	GLU
1	В	145[A]	GLU
1	В	145[B]	GLU
1	В	230	LYS
1	В	243	ARG
1	В	264	GLN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (7) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	69	GLN
1	A	168	ASN
1	A	186	ASN
1	A	231	ASN
1	В	69	GLN
1	В	168	ASN
1	В	231	ASN

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

There are no oligosaccharides in this entry.

5.6 Ligand geometry (i)

Of 7 ligands modelled in this entry, 2 are monoatomic - leaving 5 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and



the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type Chain Res			Link	Bond lengths		Bond angles			
MIOI	Type	Chain	Res	LILIK	Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
5	EDO	В	302	-	3,3,3	0.22	0	2,2,2	0.20	0
3	DGE	A	302	1	23,27,27	1.65	5 (21%)	14,39,39	2.37	6 (42%)
3	DGE	В	303	1	23,27,27	1.23	3 (13%)	14,39,39	2.41	5 (35%)
2	PEG	В	301	-	6,6,6	0.30	0	5,5,5	0.31	0
2	PEG	A	301	-	6,6,6	0.21	0	5,5,5	0.25	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	EDO	В	302	-	-	1/1/1/1	-
3	DGE	A	302	1	1/1/10/11	8/12/53/53	0/3/3/3
3	DGE	В	303	1	1/1/10/11	7/12/53/53	0/3/3/3
2	PEG	В	301	-	-	2/4/4/4	-
2	PEG	A	301	-	-	3/4/4/4	-

All (8) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	$\operatorname{Observed}(\operatorname{\mathring{A}})$	Ideal(A)
3	A	302	DGE	CAY-NAP	3.82	1.31	1.28
3	A	302	DGE	CAN-CBA	3.51	1.58	1.54
3	A	302	DGE	CAM-CBA	3.13	1.58	1.54
3	В	303	DGE	CAY-NAP	2.83	1.30	1.28
3	В	303	DGE	CAM-CBA	2.64	1.57	1.54
3	A	302	DGE	CAX-NBG	2.60	1.40	1.34
3	В	303	DGE	CAN-CBA	2.18	1.57	1.54
3	A	302	DGE	CAX-NAO	2.06	1.29	1.27

All (11) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	\mathbf{Z}	$Observed(^o)$	$Ideal(^{o})$
3	В	303	DGE	CBA-CAM-NBG	-5.56	84.22	87.96
3	В	303	DGE	CBA-CAN-NBG	-5.26	84.42	87.96

Continued on next page...



Continued from previous page...

Mol	Chain	Res	Type	Atoms	\mathbf{Z}	$\mathbf{Observed}(^o)$	$\mathbf{Ideal}(^o)$
3	A	302	DGE	CBA-CAN-NBG	-4.17	85.15	87.96
3	A	302	DGE	CBA-CAM-NBG	-3.75	85.43	87.96
3	A	302	DGE	OAG-CAW-CBF	-3.69	115.90	125.23
3	A	302	DGE	CBF-CBD-NAP	-3.16	105.80	111.94
3	A	302	DGE	OAH-CAV-OAF	-2.94	116.88	123.61
3	A	302	DGE	OAI-CAZ-CAC	-2.81	101.43	109.74
3	В	303	DGE	OAH-CAV-OAF	-2.76	117.29	123.61
3	В	303	DGE	CAJ-CAK-SAS	-2.16	103.19	106.47
3	В	303	DGE	CBF-CBD-NAP	-2.14	107.78	111.94

All (2) chirality outliers are listed below:

Mol	Chain	Res	Type	Atom
3	A	302	DGE	CBE
3	В	303	DGE	CAZ

All (21) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	A	302	DGE	CAC-CAZ-CBF-CAW
3	A	302	DGE	OAI-CAZ-CBF-CAW
3	A	302	DGE	OAI-CAZ-CBF-CBD
3	A	302	DGE	OAH-CAV-CAY-NAP
3	A	302	DGE	OAF-CAV-CAY-NAP
3	A	302	DGE	OAH-CAV-CAY-CBE
3	A	302	DGE	OAF-CAV-CAY-CBE
3	В	303	DGE	CAC-CAZ-CBF-CAW
3	В	303	DGE	CBB-CBD-CBF-CAZ
2	В	301	PEG	O1-C1-C2-O2
2	В	301	PEG	O2-C3-C4-O4
5	В	302	EDO	O1-C1-C2-O2
2	A	301	PEG	O1-C1-C2-O2
2	A	301	PEG	C4-C3-O2-C2
2	A	301	PEG	C1-C2-O2-C3
3	В	303	DGE	OAH-CAV-CAY-CBE
3	В	303	DGE	CAC-CAZ-CBF-CBD
3	В	303	DGE	OAH-CAV-CAY-NAP
3	В	303	DGE	OAF-CAV-CAY-NAP
3	В	303	DGE	OAI-CAZ-CBF-CAW
3	A	302	DGE	CBB-CBE-SAT-CBA


There are no ring outliers.




4				1 1	•	4	1 .	1 1
-4	monomers	are	1000	lved	ın	4	short	contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
5	В	302	EDO	1	0
3	В	303	DGE	1	0
2	В	301	PEG	1	0
2	A	301	PEG	2	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less then 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.

















5.7 Other polymers (i)

There are no such residues in this entry.



5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ>2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95^{th} percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ $>$	# RSRZ > 2		Z>2	$OWAB(A^2)$	Q<0.9
1	A	$267/275 \ (97\%)$	-0.76	0	100	100	12, 24, 36, 47	2 (0%)
1	В	267/275~(97%)	-0.74	0	100	100	14, 25, 39, 48	1 (0%)
All	All	534/550~(97%)	-0.75	0	100	100	12, 25, 38, 48	3 (0%)

There are no RSRZ outliers to report.

6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates (i)

There are no oligosaccharides in this entry.


6.4 Ligands (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95^{th} percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	$\mathbf{B} ext{-}\mathbf{factors}(\mathring{\mathbf{A}}^2)$	Q < 0.9
3	DGE	A	302	25/25	0.91	0.10	23,32,55,56	0
2	PEG	В	301	7/7	0.92	0.09	30,32,39,40	0
2	PEG	A	301	7/7	0.93	0.09	29,30,31,32	0
3	DGE	В	303	25/25	0.94	0.08	22,30,52,56	0
5	EDO	В	302	4/4	0.96	0.06	29,32,32,34	0
4	CL	A	303	1/1	0.98	0.16	51,51,51,51	0
4	CL	A	304	1/1	0.99	0.03	37,37,37,37	0




The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.







Electron density around PEG B 301:

 $2 {\rm mF}_o\text{-}{\rm DF}_c$ (at 0.7 rmsd) in gray ${\rm mF}_o\text{-}{\rm DF}_c$ (at 3 rmsd) in purple (negative) and green (positive)










Electron density around PEG A 301:


 $2 {\rm mF}_o\text{-}{\rm DF}_c$ (at 0.7 rmsd) in gray ${\rm mF}_o\text{-}{\rm DF}_c$ (at 3 rmsd) in purple (negative) and green (positive)






















Electron density around CL A 303:

 $2 {
m mF}_o {
m -DF}_c$ (at 0.7 rmsd) in gray ${
m mF}_o {
m -DF}_c$ (at 3 rmsd) in purple (negative) and green (positive)









6.5 Other polymers (i)

There are no such residues in this entry.

