

Full wwPDB X-ray Structure Validation Report (i)

Oct 29, 2025 – 02:09 PM JST

PDB ID : 9KM1 / pdb 00009km1

Title: Crystal Structure of the bromodomain of human BRD9 in complex with the

inhibitor Y22077

Authors: Chen, Z.; Zhang, C.; Xu, H.; Wu, X.; Zhang, Y.; Xu, Y.

Deposited on : 2024-11-15

Resolution : 2.68 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org
A user guide is available at

https://www.wwpdb.org/validation/2017/XrayValidationReportHelp with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (1)) were used in the production of this report:

 $Mol Probity \quad : \quad \text{4-5-2 with Phenix} 2.0$

Mogul : 1.8.5 (274361), CSD as541be (2020)

Xtriage (Phenix) : 2.0 EDS : 3.0

buster-report : 1.1.7 (2018)

Percentile statistics : 20231227.v01 (using entries in the PDB archive December 27th 2023)

CCP4 : 9.0.010 (Gargrove)

Density-Fitness : 1.0.12

Ideal geometry (proteins) : Engh & Huber (2001) Ideal geometry (DNA, RNA) : Parkinson et al. (1996)


Validation Pipeline (wwPDB-VP) : 2.46

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: $X\text{-}RAY\ DIFFRACTION$

The reported resolution of this entry is 2.68 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive $(\# \mathrm{Entries})$	$\begin{array}{c} {\rm Similar\ resolution} \\ (\#{\rm Entries},{\rm resolution\ range}(\mathring{\rm A})) \end{array}$
R_{free}	164625	4708 (2.70-2.66)
Clashscore	180529	5138 (2.70-2.66)
Ramachandran outliers	177936	5071 (2.70-2.66)
Sidechain outliers	177891	5071 (2.70-2.66)
RSRZ outliers	164620	4708 (2.70-2.66)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain					
1	A	149	9%	66%		30%	_	
1	В	149	6%	64%	5%	30%	-	



2 Entry composition (i)

There are 4 unique types of molecules in this entry. The entry contains 1779 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

• Molecule 1 is a protein called Bromodomain-containing protein 9.

	\mathbf{Mol}	Chain	Residues	${f Atoms}$			ZeroOcc	AltConf	Trace		
	1	ι Λ	105	Total	С	N	О	S	0	0	0
	1 A	Α		858	557	144	149	8		U	
Ī	1	D	104	Total	С	N	О	S	0	0	0
	1	Б	104	850	551	143	148	8			

There are 86 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	111	MET	-	initiating methionine	UNP Q9H8M2
A	112	LYS	-	expression tag	UNP Q9H8M2
A	113	LYS	-	expression tag	UNP Q9H8M2
A	114	GLY	-	expression tag	UNP Q9H8M2
A	115	HIS	-	expression tag	UNP Q9H8M2
A	116	HIS	-	expression tag	UNP Q9H8M2
A	117	HIS	-	expression tag	UNP Q9H8M2
A	118	HIS	-	expression tag	UNP Q9H8M2
A	119	HIS	-	expression tag	UNP Q9H8M2
A	120	HIS	-	expression tag	UNP Q9H8M2
A	121	GLU	-	expression tag	UNP Q9H8M2
A	122	ASN	-	expression tag	UNP Q9H8M2
A	123	LEU	-	expression tag	UNP Q9H8M2
A	124	TYR	-	expression tag	UNP Q9H8M2
A	125	PHE	-	expression tag	UNP Q9H8M2
A	126	GLN	-	expression tag	UNP Q9H8M2
A	127	GLY	-	expression tag	UNP Q9H8M2
A	128	GLY	-	expression tag	UNP Q9H8M2
A	129	SER	-	expression tag	UNP Q9H8M2
A	130	LEU	-	expression tag	UNP Q9H8M2
A	131	LYS	-	expression tag	UNP Q9H8M2
A	132	LEU	-	expression tag	UNP Q9H8M2
A	133	SER	=	expression tag	UNP Q9H8M2
A	240	GLU	-	expression tag	UNP Q9H8M2
A	241	ARG	-	expression tag	UNP Q9H8M2



 $Continued\ from\ previous\ page...$


Chain	Residue	Modelled Modelled	Actual	Comment	Reference
A	242	LEU	-	expression tag	UNP Q9H8M2
A	243	LEU	-	expression tag	UNP Q9H8M2
A	244	ALA	-	expression tag	UNP Q9H8M2
A	245	LEU	_	expression tag	UNP Q9H8M2
A	246	LYS	-	expression tag	UNP Q9H8M2
A	247	ARG	-	expression tag	UNP Q9H8M2
A	248	SER	-	expression tag	UNP Q9H8M2
A	249	MET	-	expression tag	UNP Q9H8M2
A	250	SER	-	expression tag	UNP Q9H8M2
A	251	PHE	-	expression tag	UNP Q9H8M2
A	252	MET	-	expression tag	UNP Q9H8M2
A	253	GLN	-	expression tag	UNP Q9H8M2
A	254	ASP	-	expression tag	UNP Q9H8M2
A	255	MET	-	expression tag	UNP Q9H8M2
A	256	ASP	-	expression tag	UNP Q9H8M2
A	257	PHE	-	expression tag	UNP Q9H8M2
A	258	SER	-	expression tag	UNP Q9H8M2
A	259	GLN	_	expression tag	UNP Q9H8M2
В	111	MET	-	initiating methionine	UNP Q9H8M2
В	112	LYS	-	expression tag	UNP Q9H8M2
В	113	LYS	_	expression tag	UNP Q9H8M2
В	114	GLY	-	expression tag	UNP Q9H8M2
В	115	HIS	-	expression tag	UNP Q9H8M2
В	116	HIS	-	expression tag	UNP Q9H8M2
В	117	HIS	-	expression tag	UNP Q9H8M2
В	118	HIS	-	expression tag	UNP Q9H8M2
В	119	HIS	-	expression tag	UNP Q9H8M2
В	120	HIS	-	expression tag	UNP Q9H8M2
В	121	GLU	-	expression tag	UNP Q9H8M2
В	122	ASN	-	expression tag	UNP Q9H8M2
В	123	LEU	-	expression tag	UNP Q9H8M2
В	124	TYR	-	expression tag	UNP Q9H8M2
В	125	PHE	-	expression tag	UNP Q9H8M2
В	126	GLN	-	expression tag	UNP Q9H8M2
В	127	GLY	-	expression tag	UNP Q9H8M2
В	128	GLY		expression tag	UNP Q9H8M2
В	129	SER	-	expression tag	UNP Q9H8M2
В	130	LEU	-	expression tag	UNP Q9H8M2
В	131	LYS	-	expression tag	UNP Q9H8M2
В	132	LEU	-	expression tag	UNP Q9H8M2
В	133	SER	-	expression tag	UNP Q9H8M2
В	240	GLU	-	expression tag	UNP Q9H8M2



Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
В	241	ARG	-	expression tag	UNP Q9H8M2
В	242	LEU	-	expression tag	UNP Q9H8M2
В	243	LEU	-	expression tag	UNP Q9H8M2
В	244	ALA	-	expression tag	UNP Q9H8M2
В	245	LEU	-	expression tag	UNP Q9H8M2
В	246	LYS	-	expression tag	UNP Q9H8M2
В	247	ARG	-	expression tag	UNP Q9H8M2
В	248	SER	-	expression tag	UNP Q9H8M2
В	249	MET	-	expression tag	UNP Q9H8M2
В	250	SER	-	expression tag	UNP Q9H8M2
В	251	PHE	-	expression tag	UNP Q9H8M2
В	252	MET	-	expression tag	UNP Q9H8M2
В	253	GLN	-	expression tag	UNP Q9H8M2
В	254	ASP	-	expression tag	UNP Q9H8M2
В	255	MET	-	expression tag	UNP Q9H8M2
В	256	ASP	-	expression tag	UNP Q9H8M2
В	257	PHE	-	expression tag	UNP Q9H8M2
В	258	SER	-	expression tag	UNP Q9H8M2
В	259	GLN	-	expression tag	UNP Q9H8M2

• Molecule 2 is 1-[1-[4-(imidazol-1-ylmethyl)-3,5-dimethoxy-phenyl]indolizin-3-yl]propan-1-on e (CCD ID: A1EFZ) (formula: $C_{23}H_{23}N_3O_3$) (labeled as "Ligand of Interest" by depositor).




Mol	Chain	Residues	A	Atoms		ZeroOcc	AltConf	
2	A	1	Total 29	C 23	N 3	O 3	0	0



 $Continued\ from\ previous\ page...$

Mol	Chain	Residues	A	ton	ns		ZeroOcc	AltConf
2	В	1	Total 29	C 23	N 3	O 3	0	0

• Molecule 3 is 1,2-ETHANEDIOL (CCD ID: EDO) (formula: $C_2H_6O_2$).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	1	Total C O 4 2 2	0	0
3	В	1	Total C O 4 2 2	0	0

• Molecule 4 is water.


\mathbf{Mol}	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	1	Total O 1 1	0	0
4	В	4	Total O 4 4	0	0



3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Bromodomain-containing protein 9





4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 2 21 21	Depositor
Cell constants	29.82Å 68.29Å 145.65Å	Donositor
a, b, c, α , β , γ	90.00° 90.00° 90.00°	Depositor
Resolution (Å)	72.82 - 2.68	Depositor
rtesolution (A)	72.82 - 2.68	EDS
% Data completeness	99.8 (72.82-2.68)	Depositor
(in resolution range)	99.8 (72.82-2.68)	EDS
R_{merge}	0.08	Depositor
R_{sym}	(Not available)	Depositor
$< I/\sigma(I) > 1$	2.71 (at 2.69Å)	Xtriage
Refinement program	REFMAC 5.8.0189	Depositor
D D.	0.250 , 0.300	Depositor
R, R_{free}	0.252 , 0.294	DCC
R_{free} test set	460 reflections (5.13%)	wwPDB-VP
Wilson B-factor (Å ²)	65.4	Xtriage
Anisotropy	0.337	Xtriage
Bulk solvent $k_{sol}(e/Å^3)$, $B_{sol}(Å^2)$	0.36, 48.2	EDS
L-test for twinning ²	$ < L >=0.48, < L^2>=0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.92	EDS
Total number of atoms	1779	wwPDB-VP
Average B, all atoms (Å ²)	72.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: The largest off-origin peak in the Patterson function is 6.57% of the height of the origin peak. No significant pseudotranslation is detected.

²Theoretical values of <|L|>, $<L^2>$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



¹Intensities estimated from amplitudes.

5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: EDO, A1EFZ

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond	lengths	Bond angles		
	Chain	RMSZ	# Z > 5	RMSZ	# Z > 5	
1	A	0.70	0/880	0.92	0/1181	
1	В	0.66	0/872	0.89	0/1170	
All	All	0.68	0/1752	0.90	0/2351	

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	858	0	868	2	0
1	В	850	0	857	2	0
2	A	29	0	0	0	0
2	В	29	0	0	0	0
3	A	4	0	6	0	0
3	В	4	0	6	0	0
4	A	1	0	0	0	0
4	В	4	0	0	0	0
All	All	1779	0	1737	4	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 1.



All (4) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	$\begin{array}{c} \text{Interatomic} \\ \text{distance (Å)} \end{array}$	$\begin{array}{c} \text{Clash} \\ \text{overlap } (\text{\AA}) \end{array}$
1:A:145:LEU:HB2	1:A:190:ILE:HD11	1.77	0.66
1:B:146:GLU:HG3	1:B:190:ILE:HG21	2.00	0.43
1:A:205:PHE:CZ	1:A:233:GLY:HA3	2.53	0.43
1:B:233:GLY:O	1:B:237:MET:HG2	2.22	0.40

There are no symmetry-related clashes.

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured Allowed		Outliers Percenti		ntiles
1	A	103/149 (69%)	102 (99%)	1 (1%)	0	100	100
1	В	102/149 (68%)	101 (99%)	1 (1%)	0	100	100
All	All	205/298~(69%)	203 (99%)	2 (1%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles		
1	A	93/132 (70%)	90 (97%)	3 (3%)	34 60		
1	В	92/132 (70%)	88 (96%)	4 (4%)	25 49		



Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
All	All	185/264 (70%)	178 (96%)	7 (4%)	28 53	

All (7) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	190	ILE
1	A	239	LYS
1	A	241	ARG
1	В	175	MET
1	В	187	LYS
1	В	230	LEU
1	В	240	GLU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	151	GLN
1	В	231	HIS

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

There are no oligosaccharides in this entry.

5.6 Ligand geometry (i)

4 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond



length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths		Bond angles			
WIOI	Type				Counts	RMSZ	# Z >2	Counts	RMSZ	# Z > 2
2	A1EFZ	В	301	-	28,32,32	2.07	6 (21%)	32,45,45	2.15	10 (31%)
3	EDO	В	302	_	3,3,3	0.54	0	2,2,2	0.30	0
3	EDO	A	302	_	3,3,3	0.81	0	2,2,2	0.53	0
2	A1EFZ	A	301	_	28,32,32	2.42	7 (25%)	32,45,45	1.98	5 (15%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	A1EFZ	В	301	-	-	4/14/18/18	0/4/4/4
3	EDO	В	302	-	-	1/1/1/1	-
3	EDO	A	302	-	-	1/1/1/1	-
2	A1EFZ	A	301	-	-	8/14/18/18	0/4/4/4

All (13) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	$\operatorname{Observed}(\textup{\AA})$	$Ideal(\AA)$
2	A	301	A1EFZ	CAI-CAM	-8.61	1.34	1.49
2	В	301	A1EFZ	CAI-CAM	-6.77	1.37	1.49
2	В	301	A1EFZ	CAS-CAP	-4.58	1.43	1.51
2	A	301	A1EFZ	CAS-CAP	-4.58	1.43	1.51
2	A	301	A1EFZ	CAF-CAE	-3.39	1.34	1.41
2	A	301	A1EFZ	CAH-CAG	-3.29	1.34	1.39
2	A	301	A1EFZ	CAC-NAD	3.23	1.43	1.38
2	A	301	A1EFZ	CAH-CAI	-3.02	1.33	1.40
2	В	301	A1EFZ	CAF-CAE	-2.94	1.35	1.41
2	В	301	A1EFZ	CAH-CAG	-2.46	1.35	1.39
2	В	301	A1EFZ	CAH-CAI	-2.21	1.35	1.40
2	A	301	A1EFZ	CAI-CAE	-2.18	1.35	1.40
2	В	301	A1EFZ	CAC-NAD	2.17	1.41	1.38

All (15) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	\mathbf{Z}	$Observed(^o)$	$\operatorname{Ideal}({}^{o})$
2	A	301	A1EFZ	CBC-CAK-CAJ	-8.28	101.87	114.03

 $\overline{C}ontinued\ on\ next\ page...$



Continued from previous page...

Mol	Chain	Res	Type	Atoms	\mathbf{Z}	$Observed(^o)$	$Ideal(^{o})$
2	В	301	A1EFZ	CBC-CAK-CAJ	-7.77	102.62	114.03
2	В	301	A1EFZ	OAL-CAJ-CAG	4.23	124.81	120.14
2	A	301	A1EFZ	OAL-CAJ-CAG	3.87	124.41	120.14
2	В	301	A1EFZ	CAS-NAT-CAU	3.25	130.65	126.25
2	В	301	A1EFZ	CAS-NAT-CAX	-2.84	122.34	125.60
2	A	301	A1EFZ	CBB-OBA-CAO	2.51	121.32	117.53
2	В	301	A1EFZ	CAZ-OAY-CAQ	2.49	121.29	117.53
2	A	301	A1EFZ	CAS-NAT-CAU	2.47	129.59	126.25
2	В	301	A1EFZ	CAR-CAM-CAI	2.32	124.50	120.70
2	В	301	A1EFZ	OAL-CAJ-CAK	-2.14	117.08	120.63
2	В	301	A1EFZ	CAM-CAN-CAO	2.09	122.95	120.13
2	В	301	A1EFZ	CBB-OBA-CAO	2.08	120.67	117.53
2	В	301	A1EFZ	CAK-CAJ-CAG	-2.05	115.92	118.00
2	A	301	A1EFZ	CAU-NAT-CAX	-2.04	106.20	108.21

There are no chirality outliers.

All (14) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	301	A1EFZ	OAL-CAJ-CAK-CBC
2	A	301	A1EFZ	CAG-CAJ-CAK-CBC
2	В	301	A1EFZ	CAG-CAJ-CAK-CBC
2	A	301	A1EFZ	CAR-CAQ-OAY-CAZ
2	A	301	A1EFZ	CAP-CAQ-OAY-CAZ
3	A	302	EDO	O1-C1-C2-O2
3	В	302	EDO	O1-C1-C2-O2
2	В	301	A1EFZ	OAL-CAJ-CAK-CBC
2	В	301	A1EFZ	CAP-CAS-NAT-CAX
2	A	301	A1EFZ	CAN-CAO-OBA-CBB
2	A	301	A1EFZ	CAP-CAS-NAT-CAU
2	A	301	A1EFZ	CAP-CAO-OBA-CBB
2	A	301	A1EFZ	CAP-CAS-NAT-CAX
2	В	301	A1EFZ	CAP-CAS-NAT-CAU


There are no ring outliers.

No monomer is involved in short contacts.


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less then 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier.



Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.







5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ>2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95^{th} percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ $>$	$\#\mathrm{RSRZ}{>}2$	$\mathbf{OWAB}(\mathrm{\AA}^2)$	Q < 0.9
1	A	105/149 (70%)	0.91	14 (13%) 8 7	43, 69, 108, 130	0
1	В	104/149~(69%)	0.76	9 (8%) 17 16	45, 65, 109, 117	0
All	All	209/298 (70%)	0.84	23 (11%) 12 11	43, 68, 109, 130	0

All (23) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	242	LEU	4.8
1	A	241	ARG	3.5
1	В	239	LYS	3.5
1	A	190	ILE	3.3
1	A	141	ILE	3.0
1	A	175	MET	2.7
1	A	195	TYR	2.7
1	В	191	VAL	2.7
1	A	138	SER	2.7
1	В	140	PRO	2.6
1	В	241	ARG	2.6
1	A	239	LYS	2.6
1	A	191	VAL	2.5
1	В	141	ILE	2.5
1	В	148	PHE	2.4
1	A	150	ARG	2.4
1	A	163	PHE	2.3
1	A	231	HIS	2.2
1	В	145	LEU	2.2
1	В	238	SER	2.1
1	В	234	PHE	2.1
1	A	193	ASN	2.0
1	A	205	PHE	2.0



6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates (i)

There are no oligosaccharides in this entry.

6.4 Ligands (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95^{th} percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	$\mathbf{B} ext{-}\mathbf{factors}(\mathring{\mathbf{A}}^2)$	Q < 0.9
3	EDO	A	302	4/4	0.76	0.21	47,48,49,50	0
3	EDO	В	302	4/4	0.84	0.24	61,61,66,68	0
2	A1EFZ	A	301	29/29	0.94	0.09	40,48,58,60	0
2	A1EFZ	В	301	29/29	0.94	0.09	41,47,55,60	0


The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.



Electron density around A1EFZ A 301: $2 \mathrm{mF}_o\text{-}\mathrm{DF}_c$ (at 0.7 rmsd) in gray ${ m mF}_o{ m -DF}_c$ (at 3 rmsd) in purple (negative) and green (positive) Electron density around A1EFZ B 301: $2 \mathrm{mF}_o\text{-}\mathrm{DF}_c$ (at 0.7 rmsd) in gray

 $2 {\rm mF}_o\text{-}{\rm DF}_c$ (at 0.7 rmsd) in gray ${\rm mF}_o\text{-}{\rm DF}_c$ (at 3 rmsd) in purple (negative) and green (positive)







6.5 Other polymers (i)

There are no such residues in this entry.

