

Full wwPDB X-ray Structure Validation Report (i)

Oct 5, 2025 – 02:47 PM JST

PDB ID : 9LOI / pdb 00009loi

Title: Crystal Structure of the Heterodimeric HIF-2 Complex with Agonist SD-16

Authors: Wu, D.; Zhuang, J.; Li, F.; Chen, X.; Diao, X.

Deposited on : 2025-01-23

Resolution : 2.78 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
https://www.wwpdb.org/validation/2017/XrayValidationReportHelp
with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (1)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0

Mogul : 1.8.5 (274361), CSD as541be (2020)

Xtriage (Phenix) : 2.0

EDS : 3.0

buster-report : 1.1.7 (2018)

Percentile statistics : 20231227.v01 (using entries in the PDB archive December 27th 2023)

CCP4 : 9.0.010 (Gargrove)

Density-Fitness : 1.0.12

Ideal geometry (proteins) : Engh & Huber (2001) Ideal geometry (DNA, RNA) : Parkinson et al. (1996)


Validation Pipeline (wwPDB-VP) : 2.46

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: X-RAY DIFFRACTION

The reported resolution of this entry is 2.78 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive $(\# \mathrm{Entries})$	$\begin{array}{c} {\rm Similar\ resolution} \\ (\#{\rm Entries},{\rm resolution\ range}(\mathring{\rm A})) \end{array}$
R_{free}	164625	4924 (2.80-2.76)
Clashscore	180529	5458 (2.80-2.76)
Ramachandran outliers	177936	5386 (2.80-2.76)
Sidechain outliers	177891	5388 (2.80-2.76)
RSRZ outliers	164620	4926 (2.80-2.76)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain							
1	A	384	3%	42%		22%	·	33%		
2	В	368	2%	58%			21%	·	20%	



2 Entry composition (i)

There are 4 unique types of molecules in this entry. The entry contains 4502 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

• Molecule 1 is a protein called Aryl hydrocarbon receptor nuclear translocator.

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf	Trace		
1	Λ	257	Total	С	N	О	S	0	0	0
1	Α	201	2089	1325	364	384	16	0	U	U

There is a discrepancy between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
Α	81	MET	-	initiating methionine	UNP P53762

• Molecule 2 is a protein called Endothelial PAS domain-containing protein 1.


Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace	
2	В	296	Total 2379	C 1503	N 404	O 448	S 24	0	0	0

There are 8 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
В	2	MET	-	initiating methionine	UNP P97481
В	363	GLU	-	expression tag	UNP P97481
В	364	HIS	-	expression tag	UNP P97481
В	365	HIS	-	expression tag	UNP P97481
В	366	HIS	-	expression tag	UNP P97481
В	367	HIS	-	expression tag	UNP P97481
В	368	HIS	-	expression tag	UNP P97481
В	369	HIS	-	expression tag	UNP P97481

• Molecule 3 is 6-chloranyl-1,1-bis(oxidanylidene)- {N}-(2-pyrrolidin-1-ylphenyl)-1,2-be nzothiazol-3-amine (CCD ID: A1EKA) (formula: C₁₇H₁₆ClN₃O₂S) (labeled as "Ligand of Interest" by depositor).





Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	
2	D	1	Total	С	Cl	N	О	S	0	0
3	Б	1	24	17	1	3	2	1	0	U

• Molecule 4 is water.


\mathbf{Mol}	Chain	Residues	${f Atoms}$	ZeroOcc	AltConf
4	A	3	Total O 3 3	0	0
4	В	7	Total O 7 7	0	0



3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Aryl hydrocarbon receptor nuclear translocator



4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants	49.64Å 99.61Å 77.66Å	Donositon
a, b, c, α , β , γ	90.00° 107.08° 90.00°	Depositor
Resolution (Å)	37.12 - 2.78	Depositor
Resolution (A)	37.12 - 2.78	EDS
% Data completeness	92.9 (37.12-2.78)	Depositor
(in resolution range)	92.9 (37.12-2.78)	EDS
R_{merge}	0.12	Depositor
R_{sym}	(Not available)	Depositor
$< I/\sigma(I) > 1$	$1.60 \; (at \; 2.77\text{Å})$	Xtriage
Refinement program	PHENIX (1.20.1_4487: ???)	Depositor
R, R_{free}	0.220 , 0.275	Depositor
it, it free	0.226 , 0.274	DCC
R_{free} test set	803 reflections (4.39%)	wwPDB-VP
Wilson B-factor (Å ²)	77.5	Xtriage
Anisotropy	0.241	Xtriage
Bulk solvent $k_{sol}(e/Å^3)$, $B_{sol}(Å^2)$	$0.35 \; , 70.6$	EDS
L-test for twinning ²	$< L >=0.51, < L^2>=0.34$	Xtriage
Estimated twinning fraction	0.028 for h,-k,-h-l	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	4502	wwPDB-VP
Average B, all atoms (Å ²)	88.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: The largest off-origin peak in the Patterson function is 5.94% of the height of the origin peak. No significant pseudotranslation is detected.

²Theoretical values of <|L|>, $<L^2>$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



¹Intensities estimated from amplitudes.

5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: A1EKA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain		nd lengths	Bond angles		
IVIOI		RMSZ	# Z > 5	RMSZ	# Z > 5	
1	A	0.33	1/2130 (0.0%)	0.59	3/2872 (0.1%)	
2	В	0.31	0/2427	0.58	3/3275 (0.1%)	
All	All	0.32	1/4557 (0.0%)	0.58	6/6147 (0.1%)	

All (1) bond length outliers are listed below:

\mathbf{Mol}	Chain	Res	Type	Atoms	Z	$\operatorname{Observed}(\text{\AA})$	$\operatorname{Ideal}(ext{\AA})$
1	A	116	VAL	N-CA	5.37	1.50	1.46

All (6) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	\mathbf{Z}	$Observed(^o)$	$Ideal(^{o})$
1	A	125	LYS	CA-C-N	5.82	127.11	119.84
1	A	125	LYS	C-N-CA	5.82	127.11	119.84
2	В	134	PHE	CB-CA-C	5.47	117.66	109.13
2	В	313	HIS	N-CA-C	-5.30	98.18	107.73
2	В	268	HIS	N-CA-C	-5.10	103.35	109.72
1	A	192	SER	CA-C-O	-5.01	114.59	120.20

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.



Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2089	0	2064	72	0
2	В	2379	0	2337	59	0
3	В	24	0	0	4	0
4	A	3	0	0	0	0
4	В	7	0	0	0	0
All	All	4502	0	4401	124	0

The all-atom clash score is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clash score for this structure is 14.

All (124) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic	Clash
Atom-1	Atom-2	$\operatorname{distance}\left(\operatorname{\mathring{A}} ight)$	overlap (Å)
1:A:192:SER:O	1:A:195:PRO:HD2	1.44	1.17
1:A:435:GLU:HG3	1:A:437:LEU:HD21	1.38	1.03
1:A:448:ASN:HD21	2:B:251:ASP:HB2	1.34	0.92
1:A:435:GLU:CG	1:A:437:LEU:HD21	2.02	0.90
1:A:448:ASN:ND2	2:B:251:ASP:HB2	1.89	0.87
1:A:435:GLU:HG3	1:A:437:LEU:CD2	2.05	0.85
1:A:431:SER:OG	1:A:435:GLU:HG2	1.76	0.84
1:A:137:SER:HA	1:A:140:LYS:HE2	1.63	0.81
1:A:192:SER:O	1:A:195:PRO:CD	2.29	0.80
1:A:435:GLU:CD	1:A:437:LEU:HD21	2.11	0.76
1:A:448:ASN:HD21	2:B:251:ASP:CB	1.99	0.75
2:B:240:ASP:OD1	2:B:240:ASP:N	2.16	0.73
1:A:260:ARG:HD3	1:A:311:TYR:HB3	1.72	0.71
1:A:125:LYS:H	1:A:125:LYS:CD	2.05	0.70
2:B:111:LEU:O	2:B:123:GLN:NE2	2.25	0.69
1:A:366:ARG:HB2	1:A:375:PHE:HB3	1.76	0.68
1:A:119:CYS:HB3	1:A:134:MET:HG2	1.75	0.67
2:B:177:THR:HG23	2:B:179:ARG:H	1.60	0.67
1:A:178:ILE:HG12	1:A:338:VAL:HG22	1.77	0.66
2:B:148:THR:O	2:B:166:ARG:NH2	2.30	0.65
1:A:415:VAL:HG12	1:A:423:LEU:HB2	1.79	0.64
2:B:265:ILE:HG12	2:B:317:VAL:HG11	1.78	0.64
1:A:196:VAL:HG23	1:A:197:LEU:HG	1.77	0.64
1:A:112:LEU:HD21	1:A:135:ALA:HB2	1.81	0.63
2:B:101:VAL:HG22	2:B:110:PHE:HB3	1.83	0.60
1:A:111:GLU:HA	1:A:114:ASP:HB2	1.83	0.60
2:B:247:ARG:HG3	2:B:338:MET:SD	2.42	0.60
1:A:184:GLY:O	1:A:208:THR:HA	2.01	0.60
2:B:134:PHE:O	2:B:173:LYS:HG3	2.02	0.59

Continued on next page...



 $Continued\ from\ previous\ page...$

Continued from previ		Interatomic	Clash
Atom-1	Atom-2	${\rm distance}(\mathring{\rm A})$	overlap (Å)
1:A:160:THR:HG22	1:A:161:ASP:H	1.69	0.58
1:A:175:PHE:CD1	1:A:196:VAL:HG21	2.38	0.58
1:A:104:LYS:HE2	1:A:108:TYR:CE2	2.40	0.57
1:A:104:LYS:CE	1:A:108:TYR:OH	2.52	0.57
1:A:304:VAL:HB	1:A:343:LEU:HD11	1.86	0.57
2:B:111:LEU:HD22	2:B:115:ILE:HD12	1.86	0.56
2:B:101:VAL:CG2	2:B:110:PHE:HB3	2.36	0.55
1:A:125:LYS:H	1:A:125:LYS:HD3	1.73	0.54
2:B:38:ALA:HB2	2:B:56:ILE:HD11	1.91	0.53
2:B:104:GLN:OE1	2:B:104:GLN:N	2.29	0.53
2:B:326:ILE:HD13	2:B:338:MET:HG2	1.90	0.53
1:A:104:LYS:HE3	1:A:108:TYR:OH	2.08	0.53
2:B:175:THR:HB	2:B:183:VAL:O	2.08	0.53
1:A:119:CYS:CB	1:A:134:MET:HG2	2.38	0.53
2:B:286:SER:O	2:B:290:THR:HG22	2.10	0.52
2:B:70:LEU:HA	2:B:73:VAL:HG12	1.91	0.52
1:A:447:GLN:N	1:A:447:GLN:OE1	2.44	0.51
2:B:359:THR:C	2:B:361:SER:H	2.19	0.51
1:A:445:THR:HG22	1:A:457:ILE:HG12	1.92	0.51
2:B:359:THR:C	2:B:361:SER:N	2.69	0.51
2:B:326:ILE:HD11	2:B:336:CYS:SG	2.51	0.50
1:A:413:GLN:O	1:A:417:LYS:HG2	2.11	0.50
1:A:125:LYS:CD	1:A:125:LYS:N	2.72	0.50
1:A:127:ASP:HB3	1:A:130:THR:HG23	1.92	0.50
1:A:169:LEU:HD12	1:A:173:ASP:HA	1.92	0.50
1:A:104:LYS:HE2	1:A:108:TYR:CZ	2.47	0.50
1:A:179:VAL:HA	1:A:186:VAL:HA	1.95	0.49
1:A:129:LEU:HA	1:A:132:LEU:HD12	1.94	0.49
1:A:126:PRO:HB2	1:A:130:THR:OG1	2.13	0.49
1:A:456:TYR:CE1	1:A:458:ILE:HD11	2.47	0.49
2:B:246:SER:OG	2:B:248:HIS:NE2	2.43	0.48
3:B:401:A1EKA:N1	3:B:401:A1EKA:C17	2.76	0.48
1:A:186:VAL:HG21	1:A:189:VAL:HG22	1.93	0.48
1:A:448:ASN:ND2	2:B:251:ASP:CB	2.66	0.48
2:B:63:PHE:CD1	2:B:64:LEU:HD23	2.49	0.48
1:A:431:SER:OG	1:A:435:GLU:CG	2.56	0.47
3:B:401:A1EKA:N3	3:B:401:A1EKA:C9	2.77	0.47
1:A:204:TRP:HZ2	1:A:267:MET:HE3	1.79	0.47
1:A:169:LEU:HD21	1:A:195:PRO:HG2	1.97	0.47
1:A:126:PRO:O	1:A:131:ILE:HD11	2.16	0.46
1:A:364:ILE:HG13	1:A:377:ASP:OD2	2.16	0.46

Continued on next page...



 $Continued\ from\ previous\ page...$

Continuea from preva		Interatomic	Clash
Atom-1	Atom-2	${\rm distance}(\mathring{\rm A})$	overlap (Å)
2:B:38:ALA:HB1	2:B:48:SER:HB2	1.96	0.46
1:A:164:LEU:O	1:A:168:ILE:HG12	2.15	0.46
2:B:219:LEU:HD23	2:B:219:LEU:HA	1.83	0.46
2:B:328:ASN:HB3	2:B:331:ASN:OD1	2.15	0.46
1:A:104:LYS:HE2	1:A:108:TYR:OH	2.16	0.46
2:B:245:LEU:HD11	2:B:340:VAL:HG22	1.98	0.46
2:B:261:ILE:HG13	2:B:265:ILE:HD12	1.98	0.46
2:B:326:ILE:HG12	2:B:336:CYS:O	2.14	0.46
2:B:108:MET:HE2	2:B:131:ILE:HA	1.97	0.45
2:B:248:HIS:O	2:B:336:CYS:HB2	2.16	0.45
1:A:311:TYR:CD2	2:B:93:LYS:HD3	2.52	0.45
2:B:69:LEU:HD23	2:B:202:TYR:CD2	2.51	0.45
1:A:387:GLN:N	1:A:390:GLU:OE2	2.46	0.45
1:A:456:TYR:HE1	1:A:458:ILE:HD11	1.80	0.45
1:A:127:ASP:OD1	1:A:128:LYS:N	2.50	0.45
1:A:223:GLU:H	1:A:223:GLU:HG2	1.51	0.45
1:A:139:MET:HE1	2:B:37:LEU:HD12	1.98	0.44
1:A:363:PHE:CE1	1:A:376:VAL:HG23	2.52	0.44
2:B:246:SER:HB3	2:B:341:ASN:HD21	1.82	0.44
1:A:310:GLY:HA3	1:A:338:VAL:O	2.17	0.44
1:A:412:PHE:O	1:A:415:VAL:HG22	2.18	0.44
1:A:311:TYR:O	1:A:337:LEU:HD12	2.17	0.44
1:A:176:LEU:HB2	2:B:91:TYR:CE1	2.53	0.44
1:A:401:HIS:ND1	1:A:403:GLU:HB3	2.32	0.44
1:A:419:LYS:HE2	1:A:454:ILE:HD11	2.00	0.43
1:A:418:LEU:HD12	1:A:423:LEU:HD21	2.00	0.43
1:A:439:MET:HG3	1:A:462:THR:O	2.18	0.43
2:B:267:TYR:OH	2:B:310:LEU:O	2.35	0.43
2:B:29:LYS:O	2:B:33:VAL:HG23	2.19	0.43
2:B:87:MET:HE3	2:B:87:MET:HB2	1.90	0.43
2:B:34:PHE:CZ	2:B:53:LYS:HE3	2.53	0.43
1:A:260:ARG:NH2	2:B:95:LEU:O	2.53	0.42
1:A:417:LYS:HA	1:A:417:LYS:HD3	1.87	0.42
2:B:119:MET:HE1	2:B:191:LYS:HB3	2.01	0.42
1:A:395:ASN:O	1:A:398:GLU:HB2	2.20	0.42
2:B:166:ARG:HG3	2:B:222:LEU:HD11	2.00	0.42
1:A:378:HIS:O	1:A:381:VAL:HG12	2.19	0.42
2:B:250:MET:HE3	2:B:250:MET:HB3	1.75	0.42
2:B:69:LEU:HD23	2:B:202:TYR:HD2	1.84	0.41
2:B:101:VAL:HG12	2:B:223:ILE:CG1	2.50	0.41
2:B:101:VAL:O	2:B:101:VAL:HG23	2.20	0.41

Continued on next page...



Continued from previous page...

Atom-1	Atom-2	Interatomic	Clash
Atom-1	Atom-2	${f distance}({ m \AA})$	overlap (Å)
1:A:441:THR:HG23	1:A:459:CYS:HB3	2.02	0.41
2:B:51:LEU:HD23	2:B:51:LEU:HA	1.93	0.41
2:B:96:GLU:O	2:B:228:PRO:HD2	2.20	0.41
2:B:284:LEU:HD23	2:B:284:LEU:HA	1.84	0.41
2:B:309:MET:HE3	3:B:401:A1EKA:C2	2.51	0.41
1:A:171:ALA:O	2:B:197:GLY:HA2	2.20	0.41
2:B:269:PRO:O	2:B:273:LEU:HD13	2.21	0.41
2:B:202:TYR:CD1	2:B:202:TYR:C	2.99	0.41
2:B:244:PHE:CZ	3:B:401:A1EKA:CL1	3.11	0.41
1:A:261:ARG:HA	1:A:261:ARG:HD2	1.85	0.41
1:A:258:GLY:N	1:A:312:ILE:HD11	2.36	0.40
2:B:232:PRO:HG2	2:B:338:MET:HB3	2.02	0.40
2:B:43:LEU:HB2	2:B:44:PRO:HD2	2.04	0.40

There are no symmetry-related clashes.

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Perce	ntiles
1	A	243/384 (63%)	238 (98%)	5 (2%)	0	100	100
2	В	288/368 (78%)	282 (98%)	6 (2%)	0	100	100
All	All	531/752 (71%)	520 (98%)	11 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.



The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric Outliers		Percentiles		
1	A	239/347 (69%)	225 (94%)	14 (6%)	16 41		
2	В	274/337 (81%)	264 (96%)	10 (4%)	30 61		
All	All	513/684 (75%)	489 (95%)	24 (5%)	22 51		

All (24) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	115	MET
1	A	118	THR
1	A	119	CYS
1	A	129	LEU
1	A	160	THR
1	A	183	THR
1	A	187	VAL
1	A	189	VAL
1	A	223	GLU
1	A	269	CYS
1	A A	377	ASP
1	A	407	LEU
1	A	422	VAL
1	A	458	ILE
2	В	115	ILE
2	В	196	THR
2	В	221	CYS
2	В	226	CYS
2 2	В	240	ASP
	В	258	ASP
2	В	287	GLU
2	В	319	LEU
2	В	321	THR
2	В	326	ILE

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	302	HIS
2	В	313	HIS
2	В	335	GLN



5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains i

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

There are no oligosaccharides in this entry.

5.6 Ligand geometry (i)

1 ligand is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Во	ond leng	$\overline{ ext{ths}}$	В	ond ang	eles
WIOI	Type	e Chain	nes	LIIIK	Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	A1EKA	В	401	-	26,27,27	0.57	0	36,40,40	1.07	1 (2%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	A1EKA	В	401	_	-	2/8/30/30	0/4/4/4

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	\mathbf{Z}	$Observed(^o)$	$Ideal(^{o})$
3	В	401	A1EKA	C2-C3-C6	-5.26	121.40	123.79



There are no chirality outliers.

All (2) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	В	401	A1EKA	C12-C13-N2-C14
3	В	401	A1EKA	C8-C13-N2-C14


There are no ring outliers.

1 monomer is involved in 4 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	В	401	A1EKA	4	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less then 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ>2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95^{th} percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ $>$	$\# \mathrm{RSRZ}{>}2$	$OWAB(Å^2)$	Q < 0.9
1	A	257/384~(66%)	0.52	11 (4%) 40 35	60, 89, 140, 170	0
2	В	296/368 (80%)	0.34	8 (2%) 56 50	51, 75, 141, 187	0
All	All	553/752 (73%)	0.42	19 (3%) 48 42	51, 81, 141, 187	0

All (19) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	В	149	LEU	3.8
1	A	116	VAL	3.6
2	В	362	LEU	2.9
1	A	134	MET	2.8
2	В	70	LEU	2.8
1	A	191	ASP	2.7
2	В	33	VAL	2.5
1	A	434	ARG	2.4
1	A	312	ILE	2.3
1	A	426	MET	2.2
1	A	137	SER	2.1
1	A	132	LEU	2.1
1	A	119	CYS	2.1
1	A	118	THR	2.1
2	В	327	TYR	2.1
1	A	435	GLU	2.1
2	В	237	ILE	2.1
2	В	73	VAL	2.0
2	В	202	TYR	2.0

6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.



6.3 Carbohydrates (i)


There are no oligosaccharides in this entry.

6.4 Ligands (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95^{th} percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	$\mathbf{B} ext{-}\mathbf{factors}(\mathbf{\mathring{A}}^2)$	Q<0.9
3	A1EKA	В	401	24/24	0.95	0.10	67,71,74,75	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.





6.5 Other polymers (i)

There are no such residues in this entry.

