



# Full wwPDB X-ray Structure Validation Report ⓘ

Mar 30, 2026 – 04:52 pm BST

PDB ID : 9SBB / pdb\_00009sbb  
Title : Monoclonal Antibodies from COVID-19 Convalescent Patients Target Cryptic Epitopes for Universal SARS-CoV-2 Neutralization  
Authors : Dessau, M.; Harit, A.  
Deposited on : 2025-08-08  
Resolution : 2.12 Å (reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0  
Mogul : 1.8.4, CSD as541be (2020)  
Xtriage (Phenix) : 2.0  
EDS : 3.0  
buster-report : 1.1.7 (2018)  
Percentile statistics : 20231227.v01 (using entries in the PDB archive December 27th 2023)  
CCP4 : 9.0.010 (Gargrove)  
Density-Fitness : 1.0.12  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.48.1


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.12 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	164625	7689 (2.14-2.10)
Clashscore	180529	8431 (2.14-2.10)
Ramachandran outliers	177936	8366 (2.14-2.10)
Sidechain outliers	177891	8367 (2.14-2.10)
RSRZ outliers	164620	7689 (2.14-2.10)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	H	229	 4% 77% 17% • 5%
2	L	217	 2% 92% 6% • 6%
3	R	219	 6% 80% 8% 11%

## 2 Entry composition i

There are 8 unique types of molecules in this entry. The entry contains 4997 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called TAU-2310 Fab Heavy Chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	H	217	1621	1030	265	320	6	0	0	0

- Molecule 2 is a protein called TAU-2310 Fab Light Chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	L	213	1580	985	259	331	5	0	0	0

- Molecule 3 is a protein called Spike protein S1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
3	R	194	1539	987	256	288	8	0	1	0

There are 10 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
R	536	PRO	-	expression tag	UNP P0DTC2
R	537	GLY	-	expression tag	UNP P0DTC2
R	538	THR	-	expression tag	UNP P0DTC2
R	539	LYS	-	expression tag	UNP P0DTC2
R	540	HIS	-	expression tag	UNP P0DTC2
R	541	HIS	-	expression tag	UNP P0DTC2
R	542	HIS	-	expression tag	UNP P0DTC2
R	543	HIS	-	expression tag	UNP P0DTC2
R	544	HIS	-	expression tag	UNP P0DTC2
R	545	HIS	-	expression tag	UNP P0DTC2

- Molecule 4 is GLYCEROL (CCD ID: GOL) (formula: C<sub>3</sub>H<sub>8</sub>O<sub>3</sub>) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	L	1	Total C O 6 3 3	0	0
4	R	1	Total C O 6 3 3	0	0
4	R	1	Total C O 6 3 3	0	0

- Molecule 5 is DI(HYDROXYETHYL)ETHER (CCD ID: PEG) (formula:  $C_4H_{10}O_3$ ) (labeled as "Ligand of Interest" by depositor).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	L	1	Total	C	O	0	0
			7	4	3		

- Molecule 6 is LITHIUM ION (CCD ID: LI) (formula: Li) (labeled as "Ligand of Interest" by depositor).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	L	1	Total	Li	0	0
			1	1		

- Molecule 7 is 2-acetamido-2-deoxy-beta-D-glucopyranose (CCD ID: NAG) (formula: C<sub>8</sub>H<sub>15</sub>NO<sub>6</sub>) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
7	R	1	Total	C	N	O	0	0
			14	8	1	5		

- Molecule 8 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
8	H	70	Total	O	0	0
			70	70		
8	L	74	Total	O	0	0
			74	74		
8	R	73	Total	O	0	0
			73	73		

### 3 Residue-property plots [i](#)


These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: TAU-2310 Fab Heavy Chain

Chain H: 





- Molecule 2: TAU-2310 Fab Light Chain

Chain L: 



- Molecule 3: Spike protein S1

Chain R: 



## 4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	111.80Å 87.97Å 95.76Å 90.00° 108.25° 90.00°	Depositor
Resolution (Å)	60.42 – 2.12 60.42 – 2.12	Depositor EDS
% Data completeness (in resolution range)	98.6 (60.42-2.12) 98.7 (60.42-2.12)	Depositor EDS
$R_{merge}$	0.13	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.38 (at 2.12Å)	Xtrriage
Refinement program	PHENIX v1.21.2_5419	Depositor
R, $R_{free}$	0.197 , 0.230 0.197 , 0.231	Depositor DCC
$R_{free}$ test set	2449 reflections (4.88%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	35.5	Xtrriage
Anisotropy	0.464	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.35 , 39.7	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.32$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
$F_o, F_c$ correlation	0.96	EDS
Total number of atoms	4997	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	48.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.36% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality i

### 5.1 Standard geometry i

Bond lengths and bond angles in the following residue types are not validated in this section: PEG, NAG, GOL, LI

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	H	1.01	0/1662	1.31	5/2267 (0.2%)
2	L	0.60	0/1618	0.85	0/2209
3	R	0.58	0/1586	0.90	1/2159 (0.0%)
All	All	0.76	0/4866	1.04	6/6635 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
3	R	0	3

There are no bond length outliers.

All (6) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	H	160	PRO	N-CA-CB	-7.08	94.81	102.60
1	H	130	PRO	N-CA-C	5.55	120.01	111.57
3	R	405	ASP	CA-CB-CG	5.26	117.86	112.60
1	H	194	THR	CB-CA-C	5.26	118.43	109.75
1	H	116	GLY	CA-C-O	-5.11	118.70	122.23
1	H	106	TYR	CB-CA-C	-5.04	100.94	110.01

There are no chirality outliers.

All (3) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
3	R	403	ARG	Sidechain

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Group
3	R	408	ARG	Sidechain
3	R	454	ARG	Sidechain

## 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	H	1621	0	1572	31	0
2	L	1580	0	1520	11	0
3	R	1539	0	1457	10	0
4	L	6	0	8	2	0
4	R	12	0	16	0	0
5	L	7	0	10	0	0
6	L	1	0	0	0	0
7	R	14	0	13	2	0
8	H	70	0	0	0	0
8	L	74	0	0	3	0
8	R	73	0	0	0	0
All	All	4997	0	4596	51	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (51) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:H:154:LYS:HE2	1:H:182:GLN:HE22	1.48	0.79
1:H:134:PRO:HD3	1:H:220:LYS:HE2	1.68	0.74
1:H:137:PRO:O	1:H:138:SER:HB3	1.86	0.72
1:H:158:PRO:O	1:H:211:HIS:HE1	1.77	0.68
1:H:5:LEU:HD12	1:H:115:GLN:HE22	1.59	0.68
3:R:368:LEU:HA	3:R:371:SER:OG	1.94	0.67
1:H:67:ARG:HG2	1:H:84:ASN:O	1.96	0.66
1:H:170:LEU:HD21	1:H:193:VAL:HG21	1.78	0.64
2:L:1:GLN:N	2:L:1:GLN:OE1	2.29	0.62
1:H:195:VAL:HG21	1:H:205:TYR:CZ	2.34	0.62

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:R:520:ALA:HB1	3:R:521:PRO:HD2	1.82	0.60
2:L:109:THR:HG23	8:L:431:HOH:O	2.01	0.60
1:H:211:HIS:HD2	1:H:214:SER:OG	1.87	0.57
1:H:1:GLU:N	3:R:519:HIS:ND1	2.52	0.56
1:H:137:PRO:O	1:H:138:SER:CB	2.54	0.55
1:H:211:HIS:HD2	1:H:214:SER:CB	2.20	0.55
2:L:137:LEU:HD12	2:L:183:LEU:HD23	1.90	0.54
1:H:87:ARG:HG3	1:H:89:GLU:HG2	1.91	0.53
1:H:189:LEU:C	1:H:189:LEU:HD12	2.34	0.52
1:H:154:LYS:HE2	1:H:182:GLN:NE2	2.21	0.52
1:H:195:VAL:HG22	1:H:196:PRO:HD2	1.92	0.51
1:H:211:HIS:HD2	1:H:214:SER:HB3	1.77	0.49
3:R:367:VAL:O	3:R:371:SER:HB3	2.13	0.49
3:R:359:SER:HA	3:R:524:VAL:CG2	2.43	0.48
3:R:339:GLY:HA2	7:R:601:NAG:HN2	1.79	0.48
1:H:211:HIS:CD2	1:H:214:SER:HB3	2.49	0.48
3:R:368:LEU:HD11	7:R:601:NAG:H81	1.96	0.47
1:H:195:VAL:HG21	1:H:205:TYR:CE1	2.50	0.46
1:H:5:LEU:HD12	1:H:115:GLN:NE2	2.27	0.45
3:R:365:TYR:CD2	3:R:387:LEU:HB3	2.52	0.45
1:H:199:SER:OG	1:H:203:GLN:HB3	2.17	0.45
4:L:301:GOL:H2	8:L:401:HOH:O	2.17	0.45
2:L:1:GLN:N	2:L:1:GLN:CD	2.76	0.43
1:H:161:VAL:HG23	1:H:189:LEU:HD21	2.00	0.42
1:H:39:GLN:C	1:H:92:ALA:HB1	2.44	0.42
1:H:150:GLY:HA2	1:H:165:TRP:CZ2	2.54	0.42
1:H:150:GLY:HA2	1:H:165:TRP:CH2	2.54	0.42
2:L:42:PRO:CG	4:L:301:GOL:H11	2.50	0.41
2:L:94:THR:HG23	2:L:98:ALA:O	2.21	0.41
2:L:125:PRO:HD3	2:L:137:LEU:HD23	2.02	0.41
1:H:154:LYS:CE	1:H:182:GLN:HE22	2.25	0.41
3:R:395:VAL:HG22	3:R:515:PHE:HD1	1.86	0.41
1:H:161:VAL:CG2	1:H:189:LEU:HD21	2.50	0.41
2:L:194:ARG:O	2:L:212:ALA:HB1	2.21	0.41
1:H:51:ILE:HG12	1:H:72:ARG:HD2	2.02	0.41
1:H:68:PHE:CE1	1:H:83:ILE:HG23	2.56	0.41
2:L:145:TYR:HA	2:L:146:PRO:C	2.46	0.41
2:L:109:THR:HG21	8:L:434:HOH:O	2.21	0.40
1:H:57:SER:OG	3:R:462:LYS:HE2	2.21	0.40
2:L:194:ARG:O	2:L:213:PRO:HD2	2.21	0.40
1:H:91:THR:HG23	1:H:120:THR:HA	2.03	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	H	213/229 (93%)	204 (96%)	9 (4%)	0	100	100
2	L	211/217 (97%)	208 (99%)	3 (1%)	0	100	100
3	R	193/219 (88%)	187 (97%)	6 (3%)	0	100	100
All	All	617/665 (93%)	599 (97%)	18 (3%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	H	178/187 (95%)	175 (98%)	3 (2%)	56	62
2	L	180/184 (98%)	179 (99%)	1 (1%)	84	89
3	R	168/191 (88%)	168 (100%)	0	100	100
All	All	526/562 (94%)	522 (99%)	4 (1%)	79	84

All (4) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	H	160	PRO
1	H	195	VAL

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	H	196	PRO
2	L	120	VAL

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (9) such sidechains are listed below:

Mol	Chain	Res	Type
1	H	82	GLN
1	H	84	ASN
1	H	182	GLN
1	H	211	HIS
2	L	113	GLN
3	R	334	ASN
3	R	394	ASN
3	R	487	ASN
3	R	501	ASN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

Of 6 ligands modelled in this entry, 1 is monoatomic - leaving 5 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
4	GOL	L	301	-	5,5,5	0.08	0	5,5,5	0.33	0
7	NAG	R	601	3	14,14,15	0.40	0	17,19,21	1.08	2 (11%)
4	GOL	R	602	-	5,5,5	0.09	0	5,5,5	0.29	0
5	PEG	L	302	-	6,6,6	0.11	0	5,5,5	0.10	0
4	GOL	R	603	-	5,5,5	0.10	0	5,5,5	0.29	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	GOL	L	301	-	-	1/4/4/4	-
7	NAG	R	601	3	-	3/6/23/26	0/1/1/1
4	GOL	R	602	-	-	2/4/4/4	-
5	PEG	L	302	-	-	1/4/4/4	-
4	GOL	R	603	-	-	2/4/4/4	-

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
7	R	601	NAG	O5-C1-C2	2.67	115.50	111.29
7	R	601	NAG	C1-C2-N2	-2.23	106.69	110.49

There are no chirality outliers.

All (9) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
4	R	602	GOL	O1-C1-C2-C3
7	R	601	NAG	C3-C2-N2-C7
7	R	601	NAG	C8-C7-N2-C2
7	R	601	NAG	O7-C7-N2-C2
4	L	301	GOL	C1-C2-C3-O3
4	R	603	GOL	C1-C2-C3-O3
4	R	602	GOL	O1-C1-C2-O2
4	R	603	GOL	O2-C2-C3-O3
5	L	302	PEG	O1-C1-C2-O2



There are no ring outliers.


2 monomers are involved in 4 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	L	301	GOL	2	0
7	R	601	NAG	2	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.







## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	H	217/229 (94%)	0.22	9 (4%) 42 44	28, 42, 88, 104	0
2	L	213/217 (98%)	-0.02	4 (1%) 66 68	24, 42, 80, 97	0
3	R	194/219 (88%)	0.18	14 (7%) 23 25	26, 44, 88, 107	1 (0%)
All	All	624/665 (93%)	0.13	27 (4%) 40 43	24, 43, 87, 107	1 (0%)

All (27) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	H	224	PRO	5.0
3	R	369	TYR	4.2
3	R	385	THR	4.0
2	L	1	GLN	3.8
1	H	138	SER	3.8
3	R	368	LEU	3.6
1	H	146	THR	3.5
1	H	200	LEU	3.2
2	L	2	SER	3.2
2	L	213	PRO	3.0
1	H	194	THR	2.9
1	H	115	GLN	2.8
3	R	521	PRO	2.7
3	R	334	ASN	2.7
3	R	371	SER	2.7
3	R	527	PRO	2.7
3	R	367	VAL	2.5
3	R	373	SER	2.4
3	R	386	LYS	2.3
2	L	159	PRO	2.3
3	R	522	ALA	2.3
3	R	387	LEU	2.2
1	H	195	VAL	2.2

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
3	R	365	TYR	2.1
1	H	149	LEU	2.1
1	H	202	THR	2.0
3	R	335	LEU	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

## 6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
7	NAG	R	601	14/15	0.59	0.24	91,121,130,130	0
4	GOL	L	301	6/6	0.80	0.14	57,67,68,72	0
4	GOL	R	603	6/6	0.83	0.15	71,79,85,85	0
5	PEG	L	302	7/7	0.91	0.14	51,60,72,74	0
4	GOL	R	602	6/6	0.93	0.11	41,44,46,47	0
6	LI	L	303	1/1	0.94	0.51	20,20,20,20	0




The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

**Electron density around NAG R 601:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)




**Electron density around GOL L 301:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)






**Electron density around GOL R 603:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)







**Electron density around PEG L 302:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)

**Electron density around GOL R 602:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)





## 6.5 Other polymers [i](#)

There are no such residues in this entry.