



Full wwPDB X-ray Structure Validation Report ⓘ

Dec 1, 2025 – 09:18 pm GMT

PDB ID : 9SXC / pdb_00009sxc
Title : Crystal structure of the Legionella pneumophila effector SidL (Lpg0437) in complex with its metaeffector LegA11 (Lpg0436)
Authors : Machtens, D.A.; Willerding, J.M.; Eschenburg, S.; Reubold, T.F.
Deposited on : 2025-10-09
Resolution : 2.40 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Xtriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20231227.v01 (using entries in the PDB archive December 27th 2023)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.46


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.40 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	164625	4642 (2.40-2.40)
Clashscore	180529	5218 (2.40-2.40)
Ramachandran outliers	177936	5158 (2.40-2.40)
Sidechain outliers	177891	5159 (2.40-2.40)
RSRZ outliers	164620	4642 (2.40-2.40)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	624	 2% 58% 11% 30%
2	B	275	 77% 12% 11%

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 5658 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Dot/Icm secretion system substrate.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	436	Total	C	N	O	S	0	0	0
			3499	2245	574	665	15			

There are 6 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	17	GLY	-	expression tag	UNP Q5ZYD5
A	18	ALA	-	expression tag	UNP Q5ZYD5
A	19	MET	-	expression tag	UNP Q5ZYD5
A	20	GLY	-	expression tag	UNP Q5ZYD5
A	21	GLY	-	expression tag	UNP Q5ZYD5
A	22	SER	-	expression tag	UNP Q5ZYD5

- Molecule 2 is a protein called Ankyrin repeat-containing protein.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	246	Total	C	N	O	S	0	0	0
			1973	1252	331	381	9			

There are 6 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	-5	GLY	-	expression tag	UNP Q5ZYD6
B	-4	ALA	-	expression tag	UNP Q5ZYD6
B	-3	MET	-	expression tag	UNP Q5ZYD6
B	-2	GLY	-	expression tag	UNP Q5ZYD6
B	-1	GLY	-	expression tag	UNP Q5ZYD6
B	0	SER	-	expression tag	UNP Q5ZYD6

- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	105	Total 105	O 105	0	0
3	B	81	Total 81	O 81	0	0

4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	68.53Å 93.38Å 133.67Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	47.85 – 2.40 47.85 – 2.40	Depositor EDS
% Data completeness (in resolution range)	99.9 (47.85-2.40) 100.0 (47.85-2.40)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	0.99 (at 2.32Å)	Xtriage
Refinement program	PHENIX 1.20.1-4487	Depositor
R, R_{free}	0.224 , 0.268 0.224 , 0.267	Depositor DCC
R_{free} test set	1847 reflections (5.00%)	wwPDB-VP
Wilson B-factor (Å ²)	52.5	Xtriage
Anisotropy	0.328	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.31 , 38.6	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	5658	wwPDB-VP
Average B, all atoms (Å ²)	66.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.45% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.14	0/3556	0.30	0/4788
2	B	0.13	0/2011	0.31	0/2714
All	All	0.13	0/5567	0.30	0/7502

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3499	0	3564	42	0
2	B	1973	0	1966	21	0
3	A	105	0	0	0	0
3	B	81	0	0	2	0
All	All	5658	0	5530	58	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (58) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:10:LYS:NZ	3:B:301:HOH:O	2.21	0.74

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:177:ILE:HG21	1:A:189:LEU:HD12	1.72	0.72
2:B:95:LYS:HG3	2:B:122:VAL:HG21	1.76	0.67
1:A:254:ASN:ND2	1:A:323:ALA:O	2.31	0.63
1:A:321:LEU:O	1:A:324:SER:OG	2.17	0.61
1:A:309:ILE:O	1:A:314:GLU:N	2.33	0.61
1:A:256:THR:HB	1:A:260:GLU:HB2	1.84	0.60
1:A:381:LYS:HG3	2:B:49:THR:HG21	1.84	0.59
1:A:309:ILE:HA	1:A:317:VAL:HG21	1.86	0.58
1:A:314:GLU:HG2	1:A:315:LYS:HG3	1.86	0.57
2:B:142:ILE:HG21	2:B:148:LEU:HD23	1.86	0.55
1:A:310:GLU:HG2	1:A:311:GLN:HG2	1.86	0.55
2:B:261:GLY:HA2	2:B:264:THR:HG22	1.88	0.55
1:A:314:GLU:N	1:A:314:GLU:OE1	2.40	0.54
2:B:117:PRO:HB3	2:B:145:MET:HE1	1.93	0.51
1:A:316:GLU:N	1:A:316:GLU:OE1	2.46	0.48
2:B:81:PRO:HB3	2:B:86:LYS:HA	1.96	0.48
1:A:110:THR:HB	1:A:187:GLY:HA3	1.96	0.48
2:B:118:HIS:CE1	2:B:156:ASP:HB3	2.49	0.48
1:A:539:LYS:HD3	2:B:262:ILE:HD11	1.94	0.48
2:B:185:GLN:NE2	3:B:306:HOH:O	2.46	0.48
2:B:171:LEU:O	2:B:178:ILE:HA	2.14	0.47
1:A:291:ARG:HB3	1:A:294:ASP:HB2	1.97	0.47
1:A:328:LYS:HB3	1:A:329:ARG:H	1.56	0.47
1:A:80:SER:HA	1:A:83:LYS:HE2	1.98	0.46
1:A:89:ILE:HG21	1:A:101:LYS:HG3	1.96	0.46
1:A:159:ALA:HA	1:A:205:VAL:HG22	1.97	0.46
1:A:263:ILE:HD13	1:A:619:ALA:HB1	1.98	0.46
1:A:226:LEU:O	1:A:567:LYS:NZ	2.43	0.46
1:A:145:PRO:HG3	1:A:189:LEU:HD21	1.97	0.46
1:A:358:THR:HG22	2:B:245:GLU:HA	1.98	0.46
1:A:534:VAL:HG11	1:A:585:VAL:HG13	1.97	0.46
1:A:309:ILE:H	1:A:309:ILE:HG13	1.55	0.45
2:B:102:PHE:O	2:B:106:VAL:HG23	2.16	0.45
1:A:108:LEU:HD21	1:A:135:LEU:HD22	1.99	0.45
1:A:179:GLU:HG2	1:A:191:MET:HG2	1.99	0.45
1:A:254:ASN:HD22	1:A:323:ALA:C	2.25	0.45
1:A:340:PHE:CE2	1:A:344:ARG:HG3	2.52	0.44
2:B:168:LYS:O	2:B:250:ARG:NH2	2.50	0.44
2:B:120:LEU:HD21	2:B:141:ILE:HG21	1.99	0.44
1:A:260:GLU:HG2	1:A:337:PRO:HG2	1.99	0.44
1:A:88:ALA:HB3	1:A:123:ILE:HD12	2.00	0.44

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:141:ILE:O	2:B:145:MET:HG3	2.17	0.44
1:A:272:LEU:HB3	1:A:274:ILE:HD11	2.00	0.43
1:A:265:THR:HG21	1:A:299:ILE:HG21	2.00	0.43
1:A:368:LYS:HE3	1:A:368:LYS:HB2	1.88	0.43
1:A:328:LYS:HE3	1:A:329:ARG:N	2.33	0.42
1:A:564:VAL:O	1:A:568:ARG:HG3	2.20	0.42
1:A:253:GLY:O	1:A:255:LYS:N	2.47	0.42
2:B:16:LEU:HG	2:B:20:MET:HE2	2.02	0.41
1:A:251:LEU:HD22	1:A:627:VAL:HB	2.01	0.41
1:A:396:ARG:HH22	2:B:254:GLU:CD	2.28	0.41
2:B:99:LEU:HG	2:B:103:LYS:HE3	2.02	0.41
1:A:198:LEU:HB3	1:A:199:PRO:HD3	2.02	0.41
2:B:122:VAL:HA	2:B:250:ARG:HB3	2.03	0.41
1:A:122:GLN:HG3	2:B:178:ILE:HD13	2.03	0.41
1:A:545:LYS:HA	1:A:550:LEU:HD22	2.02	0.40
1:A:277:LYS:HA	1:A:277:LYS:HD2	1.88	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	426/624 (68%)	405 (95%)	17 (4%)	4 (1%)	14	22
2	B	242/275 (88%)	235 (97%)	6 (2%)	1 (0%)	30	44
All	All	668/899 (74%)	640 (96%)	23 (3%)	5 (1%)	19	29

All (5) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	253	GLY
1	A	328	LYS

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	254	ASN
2	B	190	VAL
1	A	274	ILE

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	396/561 (71%)	384 (97%)	12 (3%)	36	57
2	B	222/244 (91%)	216 (97%)	6 (3%)	40	60
All	All	618/805 (77%)	600 (97%)	18 (3%)	37	58

All (18) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	77	ILE
1	A	132	LEU
1	A	182	LEU
1	A	217	LEU
1	A	252	VAL
1	A	264	LEU
1	A	307	GLU
1	A	309	ILE
1	A	327	LEU
1	A	328	LYS
1	A	329	ARG
1	A	579	VAL
2	B	49	THR
2	B	57	VAL
2	B	72	VAL
2	B	90	ILE
2	B	190	VAL
2	B	206	SER

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (4) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	555	ASN
2	B	123	ASN
2	B	215	ASN
2	B	231	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	436/624 (69%)	0.24	12 (2%) 55 51	36, 66, 105, 116	0
2	B	246/275 (89%)	0.14	3 (1%) 76 73	39, 61, 86, 106	0
All	All	682/899 (75%)	0.20	15 (2%) 62 59	36, 64, 103, 116	0

All (15) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	364	ILE	5.0
2	B	192	PRO	3.6
1	A	77	ILE	2.7
2	B	264	THR	2.5
1	A	629	LYS	2.4
1	A	252	VAL	2.3
1	A	261	PHE	2.3
1	A	256	THR	2.2
1	A	245	LEU	2.2
1	A	362	ASP	2.2
1	A	300	ILE	2.1
2	B	206	SER	2.1
1	A	321	LEU	2.0
1	A	308	VAL	2.0
1	A	143	ALA	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands

There are no ligands in this entry.

6.5 Other polymers

There are no such residues in this entry.