



Full wwPDB X-ray Structure Validation Report ⓘ

Nov 25, 2025 – 06:13 PM EST

PDB ID : 9YK4 / pdb_00009yk4
Title : Crystal structure of CYP3A4 bound to imidazole and tetraethylene glycol
Authors : Sevrioukova, I.F.
Deposited on : 2025-10-06
Resolution : 1.78 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtrriage (Phenix) : 2.0
EDS : 3.0
buster-report : 1.1.7 (2018)
Percentile statistics : 20231227.v01 (using entries in the PDB archive December 27th 2023)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.46


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.78 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	164625	1191 (1.78-1.78)
Clashscore	180529	1282 (1.78-1.78)
Ramachandran outliers	177936	1270 (1.78-1.78)
Sidechain outliers	177891	1270 (1.78-1.78)
RSRZ outliers	164620	1191 (1.78-1.78)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	487	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	IMD	A	602	-	-	X	-

2 Entry composition [i](#)

There are 7 unique types of molecules in this entry. The entry contains 4184 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Cytochrome P450 3A4.


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	466	3803	2483	620	676	24	0	12	0

There are 25 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	?	-	LEU	deletion	UNP P08684
A	?	-	ILE	deletion	UNP P08684
A	?	-	PRO	deletion	UNP P08684
A	?	-	ASP	deletion	UNP P08684
A	?	-	LEU	deletion	UNP P08684
A	?	-	ALA	deletion	UNP P08684
A	?	-	MET	deletion	UNP P08684
A	?	-	GLU	deletion	UNP P08684
A	?	-	THR	deletion	UNP P08684
A	?	-	TRP	deletion	UNP P08684
A	?	-	LEU	deletion	UNP P08684
A	?	-	LEU	deletion	UNP P08684
A	?	-	LEU	deletion	UNP P08684
A	?	-	ALA	deletion	UNP P08684
A	?	-	VAL	deletion	UNP P08684
A	?	-	SER	deletion	UNP P08684
A	?	-	LEU	deletion	UNP P08684
A	?	-	VAL	deletion	UNP P08684
A	?	-	LEU	deletion	UNP P08684
A	?	-	LEU	deletion	UNP P08684
A	468	SER	CYS	engineered mutation	UNP P08684
A	504	HIS	-	expression tag	UNP P08684
A	505	HIS	-	expression tag	UNP P08684
A	506	HIS	-	expression tag	UNP P08684
A	507	HIS	-	expression tag	UNP P08684


- Molecule 2 is PROTOPORPHYRIN IX CONTAINING FE (CCD ID: HEM) (formula:

- Molecule 4 is TETRAETHYLENE GLYCOL (CCD ID: PG4) (formula: $C_8H_{18}O_5$) (labeled as "Ligand of Interest" by depositor).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
			Total	C	O		
4	A	1	13	8	5	0	0
4	A	1	26	16	10	0	1

- Molecule 5 is GLYCEROL (CCD ID: GOL) (formula: $C_3H_8O_3$).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	A	1	Total C O 6 3 3	0	0
5	A	1	Total C O 6 3 3	0	0
5	A	1	Total C O 6 3 3	0	0
5	A	1	Total C O 6 3 3	0	0

- Molecule 6 is 1,2-ETHANEDIOL (CCD ID: EDO) (formula: C₂H₆O₂).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	A	1	Total C O 4 2 2	0	0
6	A	1	Total C O 4 2 2	0	0
6	A	1	Total C O 4 2 2	0	0
6	A	1	Total C O 4 2 2	0	0
6	A	1	Total C O 4 2 2	0	0
6	A	1	Total C O 4 2 2	0	0
6	A	1	Total C O 4 2 2	0	0
6	A	1	Total C O 4 2 2	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	A	1	Total C O 4 2 2	0	0
6	A	1	Total C O 4 2 2	0	0
6	A	1	Total C O 4 2 2	0	0
6	A	1	Total C O 4 2 2	0	0
6	A	1	Total C O 4 2 2	0	0
6	A	1	Total C O 4 2 2	0	0


- Molecule 7 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
7	A	214	Total O 214 214	0	0

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Cytochrome P450 3A4



4 Data and refinement statistics

Property	Value	Source
Space group	I 2 2 2	Depositor
Cell constants a, b, c, α , β , γ	77.11Å 99.00Å 133.44Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	40.57 – 1.78 40.57 – 1.78	Depositor EDS
% Data completeness (in resolution range)	99.3 (40.57-1.78) 99.3 (40.57-1.78)	Depositor EDS
R_{merge}	0.04	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.30 (at 1.78Å)	Xtrriage
Refinement program	PHENIX 1.20.1_4487	Depositor
R, R_{free}	0.184 , 0.218 0.184 , 0.218	Depositor DCC
R_{free} test set	2475 reflections (5.03%)	wwPDB-VP
Wilson B-factor (Å ²)	39.5	Xtrriage
Anisotropy	0.057	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.37 , 50.1	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	4184	wwPDB-VP
Average B, all atoms (Å ²)	52.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.41% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: PG4, GOL, HEM, EDO, IMD

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.29	0/3927	0.48	0/5314

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3803	0	3935	20	0
2	A	43	0	30	1	0
3	A	5	0	4	4	0
4	A	39	0	54	9	0
5	A	24	0	31	1	0
6	A	56	0	84	0	0
7	A	214	0	0	0	0
All	All	4184	0	4138	27	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (27) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:275:MET:HE2	1:A:295:LEU:HG	1.78	0.64
1:A:475:LEU:HD11	1:A:485:PRO:HB3	1.78	0.64
1:A:181:MET:HE3	1:A:181:MET:HA	1.86	0.57
1:A:181:MET:HE2	1:A:185:THR:HG23	1.89	0.54
1:A:257:LYS:HG2	1:A:276:ILE:HD11	1.90	0.54
3:A:602:IMD:C4	4:A:603:PG4:H61	2.38	0.54
1:A:379:LYS:O	1:A:381[B]:VAL:HG23	2.11	0.51
3:A:602:IMD:C5	4:A:603:PG4:H61	2.41	0.50
1:A:294:GLU:O	1:A:298:GLN:HG2	2.11	0.50
1:A:324:HIS:HD2	4:A:604[B]:PG4:H41	1.77	0.49
1:A:324:HIS:HA	4:A:604[B]:PG4:H51	1.95	0.48
1:A:469:LYS:H	1:A:469:LYS:HG3	1.41	0.48
1:A:70:LYS:HG3	1:A:71[A]:VAL:HG23	1.95	0.48
3:A:602:IMD:H4	4:A:603:PG4:H61	1.96	0.47
1:A:319:TYR:CZ	1:A:475:LEU:HB2	2.50	0.47
1:A:101:VAL:HG11	1:A:381[B]:VAL:HG21	1.98	0.46
1:A:416:PRO:HD2	4:A:604[B]:PG4:H12	1.99	0.45
1:A:409:THR:HG22	1:A:410:GLU:HG3	1.99	0.44
1:A:55:LYS:HD3	1:A:59:MET:HE1	2.00	0.43
4:A:604[B]:PG4:H42	4:A:604[B]:PG4:H22	1.89	0.43
1:A:329:GLN:HB2	5:A:608:GOL:H2	2.02	0.42
1:A:152:TYR:OH	1:A:192:ASN:OD1	2.32	0.42
1:A:132:LEU:HD13	1:A:290:LEU:HG	2.01	0.42
1:A:71[B]:VAL:HG11	1:A:386[B]:MET:HE1	2.03	0.41
4:A:604[B]:PG4:H52	4:A:604[B]:PG4:H71	1.50	0.41
2:A:601:HEM:HMC2	2:A:601:HEM:HBC2	2.03	0.41
3:A:602:IMD:H5	4:A:603:PG4:H61	2.02	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	474/487 (97%)	463 (98%)	11 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	438/443 (99%)	428 (98%)	10 (2%)	45	26

All (10) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	78	GLN
1	A	111	VAL
1	A	162	ARG
1	A	166	THR
1	A	235	VAL
1	A	243	ARG
1	A	261	LEU
1	A	262	GLU
1	A	406	LYS
1	A	469	LYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	197	ASN
1	A	198	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

23 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
6	EDO	A	618	-	3,3,3	0.41	0	2,2,2	0.40	0
6	EDO	A	610	-	3,3,3	0.55	0	2,2,2	0.25	0
6	EDO	A	609	-	3,3,3	0.54	0	2,2,2	0.45	0
2	HEM	A	601	3,1	42,50,50	1.57	6 (14%)	46,82,82	1.39	5 (10%)
5	GOL	A	606	-	5,5,5	0.74	0	5,5,5	1.21	0
6	EDO	A	622	-	3,3,3	0.47	0	2,2,2	0.37	0
6	EDO	A	611	-	3,3,3	0.63	0	2,2,2	0.24	0
6	EDO	A	613	-	3,3,3	0.44	0	2,2,2	0.43	0
6	EDO	A	616	-	3,3,3	0.47	0	2,2,2	0.50	0
6	EDO	A	620	-	3,3,3	0.44	0	2,2,2	0.46	0
6	EDO	A	619	-	3,3,3	0.47	0	2,2,2	0.29	0
6	EDO	A	617	-	3,3,3	0.44	0	2,2,2	0.38	0
6	EDO	A	612	-	3,3,3	0.44	0	2,2,2	0.49	0
6	EDO	A	615	-	3,3,3	0.43	0	2,2,2	0.38	0
4	PG4	A	604[B]	-	12,12,12	0.68	0	11,11,11	0.37	0
6	EDO	A	614	-	3,3,3	0.46	0	2,2,2	0.41	0
4	PG4	A	603	-	12,12,12	0.63	0	11,11,11	0.59	0
3	IMD	A	602	2	3,5,5	0.62	0	4,5,5	0.50	0
5	GOL	A	608	-	5,5,5	0.95	0	5,5,5	1.02	0
5	GOL	A	607	-	5,5,5	1.33	1 (20%)	5,5,5	0.77	0
5	GOL	A	605	-	5,5,5	1.08	0	5,5,5	1.69	1 (20%)
6	EDO	A	621	-	3,3,3	0.50	0	2,2,2	0.27	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
4	PG4	A	604[A]	-	12,12,12	0.68	0	11,11,11	0.37	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
6	EDO	A	618	-	-	0/1/1/1	-
6	EDO	A	610	-	-	0/1/1/1	-
6	EDO	A	609	-	-	1/1/1/1	-
2	HEM	A	601	3,1	-	1/12/54/54	-
5	GOL	A	606	-	-	2/4/4/4	-
6	EDO	A	622	-	-	0/1/1/1	-
6	EDO	A	611	-	-	0/1/1/1	-
6	EDO	A	613	-	-	1/1/1/1	-
6	EDO	A	616	-	-	1/1/1/1	-
6	EDO	A	620	-	-	0/1/1/1	-
6	EDO	A	619	-	-	0/1/1/1	-
6	EDO	A	617	-	-	0/1/1/1	-
6	EDO	A	612	-	-	1/1/1/1	-
6	EDO	A	615	-	-	0/1/1/1	-
4	PG4	A	604[B]	-	-	6/10/10/10	-
6	EDO	A	614	-	-	1/1/1/1	-
4	PG4	A	603	-	-	4/10/10/10	-
3	IMD	A	602	2	-	-	0/1/1/1
5	GOL	A	608	-	-	4/4/4/4	-
5	GOL	A	607	-	-	0/4/4/4	-
5	GOL	A	605	-	-	2/4/4/4	-
6	EDO	A	621	-	-	0/1/1/1	-
4	PG4	A	604[A]	-	-	6/10/10/10	-

All (7) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	601	HEM	C3C-C2C	-4.07	1.34	1.40
2	A	601	HEM	C3C-CAC	4.03	1.56	1.47
2	A	601	HEM	CAB-C3B	3.38	1.56	1.47
2	A	601	HEM	C3C-C4C	2.69	1.45	1.41
2	A	601	HEM	CMB-C2B	2.33	1.55	1.50
5	A	607	GOL	C1-C2	2.18	1.60	1.51

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	601	HEM	CMD-C2D	2.18	1.55	1.50

All (6) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	601	HEM	C4C-CHD-C1D	2.97	126.47	122.56
5	A	605	GOL	C3-C2-C1	-2.78	101.61	111.80
2	A	601	HEM	C4D-ND-C1D	2.55	108.23	105.21
2	A	601	HEM	CMC-C2C-C3C	2.48	129.63	124.68
2	A	601	HEM	C3B-C2B-C1B	2.24	108.09	106.41
2	A	601	HEM	C4A-C3A-C2A	2.15	108.49	107.00

There are no chirality outliers.

All (30) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
5	A	605	GOL	C1-C2-C3-O3
5	A	605	GOL	O2-C2-C3-O3
5	A	606	GOL	O1-C1-C2-O2
5	A	606	GOL	O1-C1-C2-C3
5	A	608	GOL	O1-C1-C2-C3
5	A	608	GOL	C1-C2-C3-O3
4	A	604[A]	PG4	O3-C5-C6-O4
4	A	603	PG4	O3-C5-C6-O4
4	A	604[B]	PG4	C5-C6-O4-C7
5	A	608	GOL	O2-C2-C3-O3
4	A	604[B]	PG4	O4-C7-C8-O5
4	A	604[A]	PG4	O2-C3-C4-O3
5	A	608	GOL	O1-C1-C2-O2
4	A	604[B]	PG4	C4-C3-O2-C2
4	A	604[B]	PG4	O3-C5-C6-O4
6	A	613	EDO	O1-C1-C2-O2
4	A	604[B]	PG4	C1-C2-O2-C3
4	A	604[A]	PG4	C3-C4-O3-C5
6	A	609	EDO	O1-C1-C2-O2
4	A	603	PG4	C6-C5-O3-C4
4	A	603	PG4	O1-C1-C2-O2
6	A	612	EDO	O1-C1-C2-O2
4	A	604[A]	PG4	O1-C1-C2-O2
4	A	604[B]	PG4	O2-C3-C4-O3
4	A	603	PG4	C3-C4-O3-C5
6	A	614	EDO	O1-C1-C2-O2

Continued on next page...

Continued from previous page...


Mol	Chain	Res	Type	Atoms
6	A	616	EDO	O1-C1-C2-O2
4	A	604[A]	PG4	C6-C5-O3-C4
4	A	604[A]	PG4	C5-C6-O4-C7
2	A	601	HEM	CAD-CBD-CGD-O2D



There are no ring outliers.

5 monomers are involved in 11 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	601	HEM	1	0
4	A	604[B]	PG4	5	0
4	A	603	PG4	4	0
3	A	602	IMD	4	0
5	A	608	GOL	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	466/487 (95%)	0.38	18 (3%) 44 51	19, 48, 86, 126	12 (2%)

All (18) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	107	PRO	3.9
1	A	28	HIS	3.0
1	A	384	ASN	2.9
1	A	166	THR	2.9
1	A	108	PHE	2.7
1	A	164	ALA	2.6
1	A	478	SER	2.6
1	A	265	GLN	2.4
1	A	135	PRO	2.3
1	A	490[A]	VAL	2.2
1	A	281	SER	2.2
1	A	238	ILE	2.1
1	A	267	HIS	2.1
1	A	239	CYS	2.1
1	A	423	ASN	2.1
1	A	289	ALA	2.0
1	A	385	GLY	2.0
1	A	243	ARG	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
6	EDO	A	612	4/4	0.68	0.19	65,68,69,70	0
6	EDO	A	611	4/4	0.81	0.13	49,51,52,59	0
6	EDO	A	621	4/4	0.81	0.14	70,76,78,79	0
6	EDO	A	619	4/4	0.82	0.12	64,70,71,74	0
6	EDO	A	613	4/4	0.82	0.12	80,86,87,93	0
6	EDO	A	616	4/4	0.84	0.16	59,59,68,69	0
4	PG4	A	604[B]	13/13	0.84	0.13	66,68,72,73	13
4	PG4	A	604[A]	13/13	0.84	0.13	64,67,73,74	13
5	GOL	A	606	6/6	0.85	0.10	54,59,65,68	0
5	GOL	A	605	6/6	0.86	0.11	38,44,56,58	0
6	EDO	A	620	4/4	0.87	0.13	66,66,68,70	0
6	EDO	A	617	4/4	0.88	0.12	77,81,84,91	0
5	GOL	A	608	6/6	0.88	0.10	61,63,65,65	0
6	EDO	A	622	4/4	0.88	0.10	44,56,56,59	0
4	PG4	A	603	13/13	0.89	0.12	48,52,59,67	13
6	EDO	A	610	4/4	0.90	0.10	45,58,60,60	0
6	EDO	A	615	4/4	0.90	0.10	63,65,70,71	0
6	EDO	A	609	4/4	0.91	0.09	37,43,43,55	0
3	IMD	A	602	5/5	0.91	0.13	44,45,52,53	0
6	EDO	A	618	4/4	0.92	0.09	57,66,67,67	0
6	EDO	A	614	4/4	0.93	0.09	58,58,62,63	0
5	GOL	A	607	6/6	0.94	0.10	40,40,41,46	0
2	HEM	A	601	43/43	0.97	0.07	34,37,41,48	0




The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around PG4 A 604 (B):

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)




**Electron density around PG4 A 604 (A):**


$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)




Electron density around PG4 A 603:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)







6.5 Other polymers [i](#)

There are no such residues in this entry.